HELMINTHOLOGICAL ABSTRACTS

incorporating

BIBLIOGRAPHY OF HELMINTHOLOGY
COMPILED FROM WORLD LITERATURE OF 1952



Prepared by the

COMMONWEALTH BUREAU OF AGRICULTURAL PARASITOLOGY

(HELMINTHOLOGY)

Published by the

COMMONWEALTH AGRICULTURAL BUREAUX, FARNHAM ROYAL, BUCKS., ENGLAND

O PROTE UP - D

HELMINTHOLOGICAL

THE TO ENTRANCE OF THE AND OF THE PARTY OF T



TOOLOGICATES SERVICIONAL NO DESCRIPTION OF TAXABLE PARTICIONS

DESCRIPTION AND PERSONS NAMED AND POST OFFICE ADDRESS NAMED ASSESSMENT ADDRESS NAMED A

HELMINTHOLOGICAL ABSTRACTS

Vol. 21, Part 6

1952

PRINCIPAL CONTENTS

GENERAL SUBJECTS

Anthelmintics, 887a, 930a, 965a. Bionomics, 880b, 980. Control, 1003. Immunity, 921a, 961c, 961d, 973. Immunology, 918b. Life-histories, 912a, 961e, 961j, 961k, 961m, 970, 982, 991. Morphology, 961s.
Pathology, 884a, 893a, 925b, 961bl, 977.
Physiology & Metabolism, 915a, 930a, 964b, 975.
Technique, 887b. Treatment, 946b, 946c, 946d, 946e, 955a, 990.

HOST DISTRIBUTION

Animals of Economic Importance Horse, 880a, 961bl Cattle, 896a, 985, 988. Sheep, 961bh, 990. Pig, 905a. Poultry, 936a, 961br. Cat, 888a. Laboratory animals, 883a.

Man, 893a, 922a, 977. Other Vertebrate Hosts

Mammals, 961h, 961i, 961v, 961w, 961z, 961bg. Birds, 961n, 961q, 961x, 961y, 961z, 961bp, 961br, 961bt. Fishes, 962a.

Plants, 886a, 921a, 943a, 961bj, 973, 984.

SYSTEMATICS, NEW SPECIES, etc.

Trematoda, 912b, 912c, 961q, 961r, 961u, 961z, 961bn, 981, 982, 1000, 1001. Cestoda, 961f, 961g, 961x, 961bf.
Nematoda, 895a, 912d, 912e, 931a, 961i, 961 l, 9610, 961t, 961ba, 961bb, 961bc, 961bc, 961bi, 961bi, 961bi, 961bi, 962a, 1002.
Acanthocephala, 961z. Hirudinea, 964b.

GEOGRAPHICAL DISTRIBUTION

EUROPE Denmark, 886a. Iceland, 1009.

Russia, 961.

Japan, 905a. Russia, 961.

NORTH AMERICA U.S.A., 888a.

WEST INDIES Puerto Rico, 896a.

SOUTH AMERICA Brazil, 948a.

HELMINTHOLOGICAL ABSTRACTS

Not an Per 6

2001

STREETINGO JANDHIESTES

GEOGRAL MIRRETTS

Anthologischer, 25 pa pilos 2624. Manuscher, 1900; 360. Coursel, 1907.

personally, party, party, party, except ery.

Life-tenorical gross, plans, p

Parketony Man, tops, upp, with yer.
Parketony or blanchism, once, proc. of sp. 1775.
Technique, 1876.

Personal pain, pain, sain, rate, rate, rate,

HOSTUSIATED TROP

Assessed of Sentencies Long-oronous

Homes steen perchi

- THE STREET STREET, S

Contact onto

AND NO. OF LANS. SHOW

THE RESPONDED AND

Manufactures of the other of the other, of the other, othe

STATE OF THE PARTY

STREET, MICH. SPECIES, OF

Toronto with once play plan, when other debte, play only more mink.

State while the party of the pa

Spiriture and spiriture and the spiriture and th

MOLLOSIALISMO TYDDIAN MONTH

APRILAZIO

when paint safe.

ADDRESS OF THE PARTY NAMED IN

Special Specia

AND ADDRESS OF THE PARTY NAMED IN

HELMINTHOLOGICAL ABSTRACTS

INCORPORATING BIBLIOGRAPHY OF HELMINTHOLOGY

FOR THE YEAR 1952 Vol. 21, Part 6

879-Acta Unio Internationalis contra Cancrum. Louvain.

MAKAR, N., 1952.—" A note on the pathogenesis of cancer of the bilharzial bladder." 8 (2), 323-332.

880-Acta Veterinaria. Belgrade.

NEVENIC, V., 1952.—[La diagnose d'oxyurosis par la célophane adhésive.] 2 (2), 129-131.

[In Serbian: French summary p. 131.]
NEVENIC, V., 1952.—[L'influence physique et chimique des facteurs sur le développement

embryonnaire de Parascaris equorum.] 2 (2), 205-208. [In Serbian: French summary p. 208.] c. KOZIĆ, L. & PETROVIĆ, Z., 1952.—[Über eine Invasion des Fohlens mit Strongyloides westeri Ihle 1917.] 2 (2), 221-222. [In Serbian: German summary p. 222.]

(880a) When skin scrapings were made from folds in the anal region of 447 horses from around Belgrade, Oxyuris equi ova were detected in 56.6%, but when the adhesive cellophane technique was applied ova were found in 61.7%. The latter method was very easy to apply.

(880b) Neither water nor dry conditions are favourable to the development of Parascaris equorum ova. They develop rapidly under moderately humid conditions. Indirect sunlight and quartz lamp rays kill the ova and larvae. 37°C. is the most optimum temperature for their development which starts only at 7°C. and ceases at 41°C. Both ova and larvae are very resistant to low temperatures. 5% lysol was the most lethal of the chemicals used. R.T.L.

(880c) It is assumed that a heavy infection with Strongyloides westeri found in a foal on the Government Agricultural Estate at Novi Sad was imported into Yugoslavia with R.T.L. UNRRA horses from the U.S.A.

881-Agricoltura Napoletana.

PALOMBI, A., 1952.—[Parasitic celworms of plants, their biology and control.] 19 (10), 5-8. [In Italian.]

882-Anales de la Facultad de Medicina. Universidad de Lima.

PESCHIERA, C. A. & BAZÁN, J., 1952.—" Las resecciones pulmonares en el tratamiento de los quistes hidáticos del pulmón. Estudio sobre 70 casos operados." 35 (4), 395-410.

883-Archiv für Experimentelle Veterinärmedizin.

KETZ, H. A., 1952.--" Über Trichinenfütterungsversuche beim Syrischen Goldhamster." 6 (4), 262-271.

(883a) Ketz reports that the golden hamster, Mesocricetus auratus, is a suitable experimental animal for Trichinella and will provide specimens of all developmental stages for teaching purposes. Hamsters readily eat infected meat (the optimal dose varied between o·1 and o·3 gm. according to the severity of infection) and are cheaper than guinea-pigs both to buy and to maintain.

^{*} Titles so marked throughout this number have not been seen in the original.

884—Archiv für Klinische Chirurgie vereinigt mit Deutsche Zeitschrift für Chirurgie.

a. DIETHELM, L., HEUCK, F. & KLOOS, K., 1952.—"Die Ascaridiasis und ihre pathogenetische Bedeutung für schwere hämorrhagische Darmveränderungen." 274 (1), 24-61.

(884a) Of 51 cases of severe acute intestinal disturbance with haemorrhage, nine were found to have Ascaris infection. These findings led Diethelm et al. to make a study of the pathogenic significance of ascariasis in such conditions. Injection of whole ascaris extract into sensitized rabbits and guinea-pigs produced changes in the intestines corresponding to those observed in man. This demonstration that an allergic reaction is probably the basis of the intestinal disturbances led to attempts at treatment with antihistamines, to which experimental animals and a single human case responded well.

A.E.F.

885-Arizona Farmer.

*a. DOUGLAS, E., 1952.—"Nematode nuisance becomes a menace; harder and harder to get good cotton stands and yields as eelworm infestations spread." 31 (15), I, 14-15.

886-Arsskrift. Kalmar Läns Södra Hushållningssällskap.

- a. HAEGERMARK, U., 1952.—" Något om nematodangrepp på kulturväxter." Year 1952, pp. 134-139.
- (886a) Following a popular description of the sugar-beet nematode, Heterodera schachtii, and the disease caused by this parasite, most of the paper is devoted to a discussion of stem nematodes, Ditylenchus dipsaci, in onions. This is a dangerous parasite on the island of Öland where onions are grown to a considerable extent. Haegermark found in 1951 that the stem nematode occurred on one-seventh or one-eighth of all the farms which grow onions. To control the parasite clean seed and crop rotation are recommended.

887-Arzneimittel-Forschung. Aulendorf.

- a. SEELKOPF, K. & GRAF, E., 1952.—" Semen Cucurbitae als Bandwurmmittel." 2 (8),
- 352-355.
 BOECKER, H., 1952.—" Die Larven der Oxyuren und ihre Bedeutung für die Therapie.
 Zur Pharmakotherapie der Oxyuriasis. V." 2 (8), 378-381.
- (887a) Although Seelkopf & Graf were able to confirm the findings of earlier workers that extracts of pumpkin seeds were effective in vitro against Tubifex rivulorum and Lumbricus terrestris they were unable to record any success from in vivo tests against Hymenolepis diminuta in rats. Various methods of extraction were tried with seeds from different countries but all were equally ineffective. The authors conclude that pumpkin seed extract cannot be regarded as an efficient anthelmintic.
- (887b) Boecker points out the value of experimental infections with *Passalurus ambiguus* in rabbits as a means of elucidating problems in the bionomics and treatment of Enterobius infection in man. He discusses the relation of bionomics to treatment with special reference to the "intramural" larval stages which, after two ecdyses, settle in the folds of the jejunum and upper ileum and are particularly resistant to anthelmintics. Boecker describes his techniques for maintaining supplies of larvae for experimental purposes and gives details of his methods for testing drugs on rabbits. He emphasizes the need for careful timing in the administration of anthelmintics so that efficacy against the various larval and adult stages can be tested.

888-Auburn Veterinarian. Alabama.

a. BAILEY, W. S. & LOWMAN, C. B., 1952.—" Verminous pneumonia." 9 (1), 36-38.

(888a) The distribution of Aelurostrongylus abstrusus, its known vectors in the U.S.A. and the symptoms as seen in two cats from the region of Newberry, South Carolina are briefly outlined. In the faeces of one case there were approximately 120,000 larvae per gramme.

889-Boletim. Directoria da Produção Animal. Rio Grande do Sul, Brazil.

*a. CORRÊA, O. & GLOSS, R., 1952.—[Use of sodium fluoride in ascariasis of swine.] 8 (12),

17-27. [In Portuguese.] CORRÉA, O. & GLOSS, R., 1952.—[Spontaneous elimination of Metastrongylus sp. by swine.] 8 (14), 28-29. [In Portuguese.]

890-Boletín de la Cámara Oficial Sindical Agraria de Barcelona.

BELTRÁN MONFERRER, J. M., 1952.—"Zoonosis transmisibles a la especie humana: la triquinosis. I." 7, 26-28.

891-Boletín Médico del Hospital Infantil. Mexico.

ANON., 1952.—" Cisticercosis encefálica en un preescolar. Historia clínica condensada." 9 (4), 458-464.

892-Boletín de Sanidad Militar. Mexico.

NETZAHUALCOYOTL DIAZCONTI, A., 1952.—" Eosinofilometria derecta en la oncocerciasis." 5 (51), 211-216.

893-Bulletin de l'Académie Nationale de Médecine. Paris.

TOULANT, P. & BOITHIAS, M., 1952.—" Les lésions du fond d'oeil dans l'onchocercose africaine." 3e Série, 136 (22/23), 378-383.

(893a) Toulant & Boithias describe the various lesions found in the back of the eye in patients with onchocerciasis. They found lesions in the choroid to be the most common. Atrophy of the optic nerve occurred frequently, even when there were very few signs of onchocerciasis, and the authors are of the opinion that the microfilariae are directly responsible for this condition. Treatment is briefly reviewed and the use of tablets of notezine placed in the cul-de-sac of the conjunctiva is suggested.

894-Bulletin de la Chambre d'Agriculture de Casablanca.

*a. DEYRAS, O., 1952.—" La cysticercose bovine." 23 (251), 13-14.

895—Bulletin of the Naniwa University. Series B. Agricultural and Natural Science.

a. NODA, R., 1952.—" On a cat hookworm, Ancylostoma longespiculatum Mönnig, 1938." 2, 11-18.

(895a) Noda collected 17 male and 28 female hookworms from four cats. Although very similar to Ancylostoma caninum they were smaller, the spicules were longer, the ventral teeth were more strongly developed, the pair of triangular lancets were longer, and the cervical papillae longer than they were wide and sharply pointed. The differences are tabulated and illustrated. The specimens appear to be identical with A. caninum var. longespiculata Mönnig, 1938 but the author considers the differences to be sufficient to justify the raising of this variety to specific rank as Ancylostoma longespiculatum.

896-Bulletin. Puerto Rico Agricultural Experiment Station, Rio Piedras.

RIVERA ANAYA, J. D. & MARTÍNEZ DE JESÚS, J., 1952.—"The extent of liver-fluke infestation of cattle in Puerto Rico. (A slaughterhouse survey.)" No. 107, 16 pp.

(896a) Of 46,565 cattle examined in 38 slaughterhouses in Puerto Rico the livers of 3,515, i.e. 7.55%, were found to be infected with Fasciola hepatica: 24.24% were from bulls, 12.26% from bullocks and 63.50% from cows. The incidence by districts, towns and sex and total weight of the condemned livers from each district are tabulated. The weight of condemned livers from the cattle totalled 34,074 lb.

897—Cahiers des Ingénieurs Agronomes.

*a. COMMUN, L., 1952.—" Insecticides, acaricides et nématocides." 7 (special number), pp. 24-28.

898-Československá Dermatologie.

*a. HORAČEK, J., 1952.—" Enterobius, Ascaris a jejich význam v dermatologii." 27 (1/2), 80-85.

899-Československá Farmacie.

a. PETRU, M., 1952.—"Chemotherapie helminthias! a testovac! methody pro anthelmintika."

b. SEKERA, A. & RAHM, J., 1952.—" Přírodní anthelmintika. I. Askaridol." 1 (8), 471-477.
 c. SEKERA, A. & RAHM, J., 1952.—" Přírodní anthelmintika. II. Santonin." 1 (10), 594-598.

900—Chacra. Revista Mensual de Agricultura, Ganadería e Industrías Anexas. Buenos Aires.

*a. GELORMINI, N., 1952.—" Ascaridosis del cerdo." 22 (260), 40.

901-Cirugía y Cirujanos. Mexico.

LARA D., J., 1952.—"Tratamiento de la oclusión intestinal causada por ascárides lumbricoides." 20 (10), 455-458.

902-Deutsche Landwirtschaftliche Presse.

*a. LÜHRS, 1952.—" Bekämpfung des Lungenwurmbefalls beim Rinde." 75, 74.

903-Deutsche Wirtschaftsgeflügelzucht.

*a. SPREHN, C., 1952.—" Die Eingeweidewurmer der Hühner (Hühner-Helminthen)." 4, 530-532.

904-Doelmatige Veehouderij.

*a. HENDRIKSE, D., 1952.—[Lungworm disease of cattle.] 1, 89–92. [In Dutch.] *b. HENDRIKSE, D., 1952.—[Liver fluke disease.] 1, 120–123. [In Dutch.]

905—Experimental Report of Government Experimental Station for Animal Hygiene, Tokyo.

a. SASAKI, N. & ISHITANI, R., 1952.—[Supplementary studies on histopathology of swine kidney worm disease.] No. 25, pp. 121-129. [In Japanese: English summary p. 129.]

(905a) In Japan swine kidney disease known to occur in the Kyushu, Chugoku and Kanto districts has now been found in Hokkaido where the infection reached 65% on the Live Stock Farm of the prefecture. The parasites were most abundant in the fatty tissue along the ureters and were also present in the liver, lungs, lymph glands, muscles and the mesenteric and subpleural tissues. The pathological changes observed were the formation of cysts and tunnels, thrombi and aneurysms, string-like upheavals of the intima of the veins and arteries, and cirrhosis of the liver. The histological changes in tunnels in the liver and mesenteric tissues were haemorrhages, white cell infiltrations and hyperplasia of the connective tissue. The cysts were surrounded by granulation tissue.

906-Feldsher i Akusherka. Moscow.

*a. LUZINA, A. G., 1952.—[Ascariasis and its control.] Year 1952, No. 2, pp. 33-38. [In Russian.]
*b. VASILKOVA, Z. G., 1952.—[Treatment of trichuriasis.] Year 1952, No. 2, pp. 54-56. [In Russian.]

*c. VASILKOVA, Z. G., 1952.—[Treatment of enterobiasis.] Year 1952, No. 2, pp. 56-57. [In Russian.]

*d. VASILKOVA, Z. G., 1952.—[Strongyloides stercoralis.] Year 1952, No. 5, p. 56. [In Russian.]

*e. VASILKOVA, Z. G., 1952.—[Hepatic parasites in man.] Year 1952, No. 7, pp. 54-55. [In Russian.]

*f. VASILKOVA, Z. G., 1952.—[Treatment of helminthiasis.] Year 1952, No. 8, pp. 55-57.

[In Russian.]

907-Flugblatt. Bundesanstalt für Pflanzenschutz. Vienna.

a. BÖHM, O., 1952 .- "Das Chrysanthemenälchen (Aphelenchoides ritzemabosi Schw.)." No. 89, 2 pp.

(907a) Böhm gives a brief, popular account of chrysanthemum eelworm with general recommendations for control including chemical methods.

908—Frontiers. Philadelphia.

BERNSTEIN, J., 1952.—"The lowly leech, bedside companion of the surgeon in the good old days." 17 (1), 12-13, 32.

909-Glas Srpske Akademije Nauka. Odeljenje Medicinskih Nauka.

SIMITCH, T. & PETROVITCH, Z., 1952.—[Study of intestinal parasites in man in Yugoslavia. Part I. Intestinal parasites in children from orphanages in Banat.] 205 (5), 231-242. [In Serbian: French summary.)

910-Groenten en Fruit.

SCHENK, P. J., 1952.—[The root nematode *Heterodera marioni*.] 7, 898. [In Dutch.] KROFT, W. G. VAN DER, 1952.—[A new method of controlling root nematodes on tomatoes. 2.] 7, 970. [In Dutch.]

911-Indian Journal of Child Health.

JUNGALWALLA, A., 1952.—"Ascariasis in children." 1 (8), 389-400.

912-Indian Journal of Helminthology.

THAPAR, G. S. & TANDON, R. S., 1952.—"On the life-history of liver-fluke, Fasciola gigantica Cobbold, 1855 in India." 4 (2), 1–36.

JAIN, S. L., 1952.—"Monogenea of Indian fresh-water fishes. I. Haplocleidus gomtius n.sp. (sub-family: Tetraonchinae) from the gills of Wallagonia attu (Bloch), from Lucknow." 4 (2),

37-42.

JAIN, S. L., 1952.—" Monogenea of Indian fresh-water fishes. II. Thaparocleidus wallagonius n.g., n.sp., (sub-family: Tetraonchinae) from the gills of Wallagonia attu (Bloch), from Lucknow." 4 (2), 43-48.

Lucknow." 4 (2), 43-48.

Lucknow." 4 (2), 43-48.

SANWAL, K. C., 1952.—" On a new avian nematode, Dispharynx ketupae n.sp., (sub-family Acuariinae, Railliet, Henry & Sisoff, 1912) from the brown fish-owl, Ketupa zeylonensis (Gmelin)." 4 (2), 49-53.

SANWAL, K. C., 1952.—"On a new avian nematode, Krusadia indica n.g., n.sp. (sub-fam. Schistorophinae Travassos 1918) from the jungle nightjar (Caprimulgus indicus)." 4 (2), 54-60.

- (912a) Thapar & Tandon give a detailed and illustrated description of the developmental stages of Fasciola gigantica in Limnaea acuminata. The miracidium hatches in 7-20 days. Rediae leave the sporocyst on the 9th day and produce daughter rediae by the 12th day. Only daughter rediae produce cercariae. Cercariae emerge 38 days after infection in April and 65 days after infection in February. The complete cycle from egg to adult varies from 21 to $4\frac{1}{2}$ months. The metacercariae can resist desiccation and can remain alive for a month in water. Only 1.4% to 3.5% of the metacercariae fed experimentally to guinea-pigs and rabbits became infective and the resulting flukes were smaller in size than immature flukes obtained R.T.L. from naturally infected cattle and buffaloes.
- (912b) Haplocleidus gomtius n.sp., from the gills of the fresh-water fish Wallagonia attu near Lucknow, differs from all other species in (i) its larger size (0.90 mm. to 0.96 mm.), (ii) its simple copulatory apparatus, (iii) the presence of additional supporting bars, (iv) the larger size of the dorsal anchor and (v) the presence of a peculiar ventral haptoral bar made up of two recurved pieces.
- (912c) Thaparocleidus wallagonius n.g., n.sp. occurred on the gills of 12 out of 24 Wallagonia attu from Lucknow. The presence of three haptoral bars distinguishes Thaparocleidus from all other genera of the Tetraonchinae except Murraytrema. It is distinguished

from Murraytrema by posteriorly confluent intestinal crura, the dissimilarity in shape and size of the haptoral bars and anchors, the sinistral position of the female genital pore and the position of the ovary which lies behind the testis.

- (912d) Dispharynx ketupae n.sp., of which one male and three females were collected from the oesophagus of Ketupa zeylonensis from Hardoi, is described. The body length is greater than in other species of Dispharynx: that of the male is 13:49 mm. and that of the female 18·15 mm. to 18·40 mm. It is further distinguished by the size of the eggs (0.031 mm. × 0.025 mm.), by possessing a single mid-ventral bell-shaped caudal papilla, by the very narrow caudal alae, by the almost straight anterior cuticular cordons and by the size of the spicules, the left spicule measuring 0.83 mm. and the right 0.225 mm.
- (912e) Krusadia indica n.g., n.sp. in the intestine of Caprimulgus indicus from the Krusadai Islands, superficially resembles Sciadiocara Skryabin, 1916 in the disposition of the cuticular head appendages but possesses six membranous cuticular appendages instead of the four winged appendages, two pointed tooth-like papillae on the dorsal and ventral sides of the lipless mouth, and more than double the number of caudal papillae which are equally spread in two longitudinal rows. The eggs are thin-shelled.

913—Japanese Journal of Plant Protection.

*a. KONDO, T., 1952.—[Root-knot nematode and its control.] 39/40, 352-355. [In Japanese.]

914-Journal of the Albert Einstein Medical Center. Philadelphia.

a. HEINE, W. I. & ROSENTHAL, A., 1952 .- "Trichinosis with cardiac involvement." 1 (1), 24-28.

915—Journal of the College of Arts and Sciences, Chiba University. Natural Science Series.

- YAMAO, Y., 1952.—[Histochemical studies on endoparasites. V. Distributions of the glyceromonophosphatases in the tissues of flukes, Eurytrema coelomaticum, E. pancreaticum, Dicrocoelium lanceatum, and Clonorchis sinensis.] 1 (1), 9-13. [In Japanese: English summary p. 13.]
- (915a) Yamao has shown that in Eurytrema coelomaticum and E. pancreaticum alkaline phosphatase is localized in the body wall and the walls of the excretory canals and gives strongly positive reactions, whereas acid phosphatase is demonstrable in most tissues, although giving only weak reactions. In Dicrocoelium dendriticum the distribution is similar to that in Eurytrema but the acid phosphatase reacts more strongly and is particularly concentrated in the mesenchymatous fibres and their nuclei. The distribution and intensity of acid phosphatase in Clonorchis sinensis are very similar to those in Fasciola hepatica but Yamao was unable to demonstrate the presence of any alkaline phosphatase in this species. S.W.

916—Journal of Sericultural Science. Tokyo.

SANO, T., 1952.-[Studies on the silkworm disease caused by mermithid nema. I.] 21, 298-301. [In Japanese.]

917-Karakulevodstvo i Zverovodstvo.

- PETROV, A. M. & PANISHEVA, L. V., 1952.—[Phenothiazine therapy against Capillaria infections of the stomach and bladder of mink and sable.] 5 (1), 70-71. [In Russian.] SOPELCHENKO, M. I., 1952.—[Dictyocaulus in sheep under conditions of semi-desert steppes of Uzbekistan.] 5 (4), 69-73. [In Russian.] DUBNITSKI, A. A., 1952.—[New cestode from the intestines of Barguzin sables.] 5 (4),

79. [In Russian.] SOPELCHENKO, M. I., 1952.—[Infestation of sheep with cestodes that live as parasites in

918-Klinicheskava Meditsina. Moscow.

DESYATOV, M. M., 1952 .- [Suppurative hepatic echinococcosis with metastases into the

right bile duct.] 30 (E), 88. [In Russian.] LEIKINA, E. S., GAIKO, B. A., CHELISHEVA, K. M. & BOKSHTEIN, M. E., 1952.— [Early immunodiagnosis of ascariasis in man and its clinical and epidemiological significance.] 30 (11), 49-53. [In Russian.]

(918b) Migration of Ascaris larvae frequently produces transitory pulmonary lesions accompanied by general indisposition and eosinophilia in the peripheral blood. That Ascaris does cause this is confirmed by the presence of specific antibodies in the blood and, when the parasites become mature, by the appearance of ova in the faeces. Precipitin reactions are a useful means of diagnosis during the initial stages of infection and also for the determination of the date of infection.

919—Kyushu Agricultural Research.

TANAKA, I., 1952.—[On the effective-bounds of certain fumigants to soil for the root-knot nematode, Heterodera marioni (Cornu) Goodey.] No. 10, pp. 147–148. [In Japanese.] GOTO, S. & HASUKO, E., 1952.—[On the treatment with chloropicrin for the nematic root-rot of sweet potato. 2.] No. 10, pp. 149–150. [In Japanese.]

920-Landwirtschaftsblatt Weser-Ems.

*a. LÜHRS, 1952.—" Die Bekämpfung der Leberegelseuche." 99, 579–580. *b. LÜHRS, 1952.—" Bekämpfung der Lungenwurmseuche der Kälber." 99, 657–658.

921-Lantmannen.

ÅKERBERG, E., 1952.—" Stamförsök med rödklöver och lucern. Några erfarenheter från årets stamförsök vid Sveriges Utsädesförenings Ultunafilial." 36 (30), 589–590.

(921a) In a field experiment stem nematode in red clover reduced the yield by 50%. There are strains with a very good resistance to stem nematode. For central Sweden it is important to obtain resistance and winterhardiness; yield data are given for two new strains, U 036 and U 056, with these characteristics.

922-Medicina y Cirugía. Bogotá.

ALBORNOZ PLATA, A., 1952.—" Estudio epidemiológico y estadístico del parasitismo intestinal en una zona de Bogotá. Período de tres años: 1947-1948-1949." 16 (10), 372-389.

(922a) The incidence of intestinal protozoon and helminth parasites in Bogotá decreased from 71.6% in 1947 to 57.8% in 1949, but that of ascariasis rose from 17.48% to 32.21%. Statistical analyses according to age revealed that in spite of the general decrease, incidence of intestinal parasitic diseases increased in the 15-19 years age group from 6.31 per thousand to 10.0 per thousand, and in the 20-24 years group from 6.67 per thousand to 23.85 per thousand, an increase attributed to the immigration of young people to the capital. M.McK.

923-Medicinski Arhiv. Sarajevo.

PAVIŠIĆ, Z., 1952.—" Operativna terapija intraokularnog cisticerkusa." 6 (1), 23-27. [English summary p. 27.]

924-Medicinski Glasnik. Belgrade.

KRSTIĆ, B. N., 1952.—" Uklanjanje pantljičare atebrinom." [Atebrin therapy of helminthiasis.] 6 (2), 47-51.

925-Mémoires de l'Institut Royal Colonial Belge. Section des Sciences Naturelles et Médicales.

SCHWETZ, J., 1952 .- "Sur la confusion actuelle dans la classification des planorbes centro-

africains et les moyens pour y remédier." Collection in-8°, 21 (4), 30 pp.
b. RODHAIN, J., 1952.—"Les adénolymphocèles du Congo belge." Collection in-8°, 21 (5),

(925b) Rodhain presents a detailed and illustrated account of the lesions of the lymphatic glands caused by filariae in the Belgian Congo, with particular reference to a number of cases of adenolymphocele. Material was obtained from three districts, one of Onchocerca volvulus infection, one of mixed O. volvulus and Wuchereria bancrofti and one of W. bancrofti alone. Three main types of pathological reaction are described. Dipetalonema streptocerca infection was not found. Rodhain's observations confirm the view that not all glandular enlargements of the groin are caused by filarial infection and he was unable to find the cause of the localization S.W. of the lesions in the groin.

926—Memoirs of the Faculty of Agriculture, Hokkaido University.

a. YAMASHITA, I. & TAKAHASHI, S., 1952.—[Epidemiologic survey of parasites of domestic animals in Hokkaido. I. A survey of the suburbs of Sapporo.] 1 (2), 140-144. [In Japanese: English summary p. 144.]

(926a) The infection rates of each helminth species, as revealed by faecal egg counts, in cattle, horses, sheep, pigs and poultry kept in farms in the suburbs of Sapporo are tabulated under each host. R.T.L.

927-Merkblätter über die Hauptsächlichsten Parasiten und Parasitengruppen.

BORCHERT, A., 1952.—" Die Cestoden unserer Haussäugetiere." No. 11, 58 pp. *a.

*b.

BORCHERT, A., 1952.—"Endoparasiten des Geflügels." No. 12, 72 pp.
BORCHERT, A., 1952.—"Knötchen-, Dickdarm-, Haar- und Peitschenwürmer der Säugetiere." No. 13, 35 pp.

928-Neurologie a Psychiatrie Československá.

KARPÍŠEK, J. & VALACH, V., 1952.—" Cysticercus racemosus měkkých plen mozkových s klinickým obrazem apoplexie." 15 (5/6), 179–183. [English & Russian summaries pp. 182–183.]

929-Osterreichische Zeitschrift für Stomatologie.

KALMI, I., 1952.—" Die Wurminvasion in der Ätiologie der paradentalen Erkrankungen der Kinder im Pubertätsalter." 49 (9), 527-528.

930—Okayama Igakkai Zasshi.

MANNAMI, C., 1952.—[Studies on the resistance of Ascaris lumbricoides against repetitive doses of alkylresorcinols.] Supplement, 64, 1271-1288. [In Japanese.]

(930a) Ascaris lumbricoides from pig and man were incubated in solutions of hexylresorcinol, cyclohexylchlororesorcinol and octylchlororesorcinol. Others were expelled by santonin after octylchlororesorcinol had been given orally to the host. Spinicauda japonica were immersed repeatedly in solutions of various concentrations of these drugs. Female Ascaris were exposed to various concentrations in vitro and in vivo and their eggs were then cultivated and the larvae liberated. In no case was any marked increase of resistance to the drugs observed. Clinical tests were also made. One or two weeks after one-third of the standard dose of cyclohexylchlororesorcinol had been administered, a standard dose was given but the effect of the latter was not inferior to that on non-premedicated controls. Other experiments on patients and on puppies are also reported. The conclusion drawn is that the resistance of Ascaris lumbricoides to alkylresorcinols is not increased by repetitive doses. minute repetitive doses of cyclohexylchlororesorcinol or octylchlororesorcinol there was a cumulative effect which suggests a new rational method of administering these drugs. [Based on an abstract in Acta Med. Okayama, 8 (3), p. xxxiii.]

931—Pakistan Journal of Science.

a. AKHTAR, S. A., 1952.—"A redescription of the species of Schistorophus longicornis." [Correspondence. 4 (3), 127-128.

(931a) Akhtar found numerous nematodes under the cuticular lining of the gizzard of a Numenius arquata at Kabul. The identification of them as Schistorophus longicornis was confirmed by Baylis although there were a number of differences from the original description, particularly in the shortness of the head lappets and the shape of the right spicule. The author redescribes and illustrates the species from his specimens.

932-Pediatrické Listy.

HORAČKOVÁ, M., HRUBCOVÁ, M., JÍROVEC, O. & POKORNÝ, M., 1952.—" Parasitologické šetření u školní mládeže na Sušicku v r. 1950." [The investigation of the schoolchildren of Suscice for parasites in the year 1950.] 7 (4), 207-211.

933-Polski Tygodnik Lekarski. Warsaw.

a. RESZKE, H., 1952.—"Rzadki przypadek wągrzycy." [A rare case of cysticerciasis.] 7 (23),

764-765. KOCZOROWSKI, S., KONORSKA, R. & KUBICZ, S., 1952.—"Rzadki przypadek bąblowca płuc." [A rare case of pulmonary echinococcosis.] 7 (37), 1134-1140. [English & Russian

summaries pp. 209*, 211*.]

KUZMICKI, R., 1952.—"Próby leczenia atebryną nosicielstwa tasiemca nieuzbrojonego (Tasnia saginata)." [Attempts at treating the carrier state of Tasnia saginata with atebrine.]

7 (42), 1333–1336. [English & Russian summaries pp. 249*—251*.]

934-Praktický Lékař. Prague.

HLOUCAL, L. & ZOUBEK, V., 1952.—" Nové připady trichinellosy na Strakonicku." [New case of trichinelliasis in Strakonice.] 32 (5), 106–109.

935—Proceedings of the California Fig Institute.

McBETH, C. W., 1952 .- "Control of nematodes on figs." Annual Research Conference (6th), pp. 14-15.

936-Proceedings of the Iowa Academy of Science.

KERR, K. B., 1952.—" Occurrence of *Echinoparyphium recurvatum* (Linstow, 1873) in chickens in Iowa." 59, 467-468.

PALMER, B. M. F., 1952.—" A comparative study of the incidence of intestinal parasites of children in Des Moines, Iowa." 59, 487-491.

(936a) In chickens in Iowa an outbreak of catarrhal enteritis with a 5% mortality was associated with numerous Echinoparyphium recurvatum. Specimens of Physa sp. collected near the brooder house contained a furcocercous cercaria, a stylet cercaria and an ovoid metacercaria. One mature E. recurvatum was recovered from a three-week-old chicken three weeks after it had been fed with the metacercaria.

937—Proceedings of the Nebraska Academy of Science.

JASKOWSKI, B. J. & BONNEMA, J. M., 1952 .- " Detergent effects on ascarid development." [Abstract.] 62, 9.

938-Ptitsevodstvo.

PETRUSHKIN, A. A. & KSENOFONTOVA, M. I., 1952.—[Phenothiazine—an effective measure in controlling ascaridiasis in hens.] 7, 30-31. [In Russian.]

939-Publicaciones del Instituto de Biolgía Aplicada. Barcelona. Serie Zoologica.

GADEA, E., 1952.-" Contribución al estudio de los nematodos libres terrestres y dulceacuicolas de la fauna espanôla." 1, 1-213.

940—Quarterly Bulletin of Northwestern University Medical School. Chicago.

PIHL, H., YEAGER, L. B. & HARDING, H. B., 1952.—"Report of a case of multiple parasitosis including *Trichostrongylus* sp. infestation." 26 (4), 321-324.

941—Quarterly Bulletin of the South Pacific Commission.

KERREST, J., 1952.—"Epidemiological aspects of bancroftian filariosis in New Caledonia."

(941a) [A French version of this paper was published in Méd. trop., 1952, 12, 568-570. For abstract see Helm. Abs., 21, No. 728c.]

942—Report of the Minister for Agriculture. Dublin.

2. ANON., 1952.—"Annual report, 1951-52." 21st (1951-52), 185 pp.+Appendices [87] pp.

(942a) The Republic of Ireland Potato Root Eelworm Order 1951 came into operation on 1st January, 1952 and 3,006 samples of soil from potato-growing areas were tested for eelworm cysts at the laboratory established for this purpose at the Albert Agricultural College. As Heterodera schachtii has not been reported from sugar-beet in Ireland, sugar-beet plants were grown in 146 soil samples from various fields throughout the country in which sugar-beet had been grown for a number of years, but failed to become infected. Other experimental work revealed the occurrence of cruciferous root eelworm. The results of routine examination of faecal samples from domesticated animals are reported by the Veterinary Research Laboratory at Drumcondra and by the Parasitology Section of the Pathological Department of the Veterinary College of Ireland. Immature lungworms and larvae were found post mortem in the bronchioles of a bullock aged one year and nine months which had shown all the symptoms of hoose but no larval lungworms in the faeces.

943—Report of the Science Service. Division of Botany and Plant Pathology, Canada.

a. BAKER, A. D., 1952.—" Notes on some nematode problems in 1952." 32nd (1952), pp. 121-124. [Reprint.]

(943a) Baker reviews briefly the occurrence in Canada of various species of plantparasitic nematodes and indicates their relative importance. One new host record is given, Swiss churd (Beta cicla L.) for Heterodera schachtii.

944—Revista Chilena de Higiene y Medicina Preventiva.

NEGHME R., A., SILVA, R. & ARTIGAS, J., 1952.—" Nuevos aspectos sobre epidemiología de la amebiasis y enteroparasitosis." 14 (4), 243-257.

945-Revista Colombiana de Pediatría y Puericultura.

a. CAMACHO GAMBA, J., 1952.—"La anemia carencial del tropico en el niño. (Necatoriasis - uncinariasis - anemia tropical.)" 12 (2), 90-128.

946-Revista Cubana de Pediatría.

ANIDO, A., VALLEDOR, R., HERNÁNDEZ DE LA BARCA, A., MACHADO ESPARZA, L. F. & MARTÍNEZ CARRERA, N., 1952.—"Parasitismo intestinal. Tratamiento. Ascarid-

L. F. & MARTINEZ CARRERA, N., 1952.— Parasiusmo intestinat. Traumento. Pascartiasis." 24 (11), 653-677.

KOURÍ, P., VALDÉS DÍAZ, R. & SOSA BENS, D., 1952.—"Recientes adquisiciones en la clínica y en la terapeutica de la tricocefaliasis infantil." 24 (12), 717-722.

LEÓN LEÓN, J. M., 1952.—"Tratamiento de la ankilostomiasis." 24 (12), 723-734.

BÉGUEZ CÉSAR, A., CASTELLANOS FONSECA, E. & TEIXIDO VAILLANT, S., 1952.—"Teniasis y hymenolepiasis en niños." 24 (12), 735-745.

PÉREZ HURTADO, F., 1952.—"Parasitismo por Strongyloides stercoralis." 24 (12), 746-758.

(946b) Seventy children with massive infections of Trichuris were cured clinically by an enema of 200 c.c. to 700 c.c. containing Santokin (a hexylresorcinol preparation) diluted 1:300 to 1:400 in tepid water. M.MCK.

- (946c) León León gives the details obtained from 5,600 faecal examinations, including 4,000 from city dwellers, undertaken at different times in Cuba. The incidence of Ascaris, Trichuris, Necator and Enterobius is tabulated. The author outlines the dosage, effectiveness and side effects of hexylresorcinol, chenopodium oil, carbon tetrachloride and tetrachlorethylene, and stresses the great need for hygienic measures, emphasizing particularly the responsibility which the literate populace should, but do not, take towards the poverty-stricken, neglected rural communities.

 M.MCK.
- (946d) In 4,000 children aged from 6 to 12 years from the Colonia Infantil de Oriente in Cuba the helminth incidence was: Trichuris 46·35%, Ascaris 24·08%, Enterobius 2·28%, Necator americanus 2·9%, Taenia saginata 0·025%, Ancylostoma duodenale 0·225% and Heterodera radicicola 0·025%. Twelve children with Taenia infections and three with Hymenolepis nana were treated. Atebrin, in single doses of 1 mg. per lb. body-weight given on an empty stomach and fol. wed by a purge, cured five out of six cases of taeniasis (one not traced) and two out of four with Hymenolepis (one not traced). Eliminol or Phenendiol, organic iodine compounds, administered with the above routine at the rate of 4 cg. per lb. body-weight cured four out of five cases of taeniasis (one not traced).
- (946e) In the region of Norte de Oriente, Cuba, 61·72% of 3,373 faecal examinations were positive for intestinal parasites, with an incidence of 0·43% for Strongyloides stercoralis. Gentian violet was administered at the rate of 0·01 gm. per day for two weeks, with an interval of a week before repeating the treatment. In some cases it produced vomiting and abdominal pains. A list is given of the morphological and biological characters by which the rhabditiform larvae of Necator americanus and S. stercoralis can be distinguished.

 M.MCK.

947-Revista Española de las Enfermedades del Aparato Digestivo y de la Nutrición.

a. GARCÍA MORÁN, J., 1952.—" Quistes hidatídicos con expresión clínica de colecistitis." 11 (3), 767-771.

948-Revista do Serviço Especial de Saúde Pública. Rio de Janeiro.

- a. COSTA, O. R. DA, 1952.—" Contribuição ao conhecimento da esquistossomose na Amazônia." 5 (2), 401-409. [English summary p. 404.]
- (948a) Costa examined 1,682 people of Amazonas State, Brazil, in seven localities on the river Tapajós near Fordlândia, a focus of schistosomiasis. The results, tabulated according to locality and age group, show that Schistosoma mansoni occurred only among immigrants from north-east Brazil. No planorbid vectors were found in the areas. The author recommends eradication of the parasite while the area of infection in Amazonas is still confined to Fordlândia.

M.MCK.

949—Revue Médicale de Nancy.

a. MELNOTTE, P., CANTEGRIT, M. & MICHON, P., 1952.—"Grande éosinophilie sanguine et médullaire chez un colonial. Discussion étiologique. Traitement antifilarien d'épreuve." 77, 603-606.

950-Rivista d'Ostetricia e Ginecologia Pratica.

*2. SASSI, R., 1952.—" Proglottidi di tenia nell'appendice. Due casi." 34 (7), 332-333.

951-Rivista di Patologia e Clinica. Parma.

- *a. FICAI, A. & RICCI, N., 1952.—" Cisti da echinococco della tiroide. Contributo clinico."
- 7 (5), 152-159. *b. LOIZZI, A., 1952.--" Su un caso di ciste idatidea intraepatica simulante un rene mastice." 7 (12), 541-550.

952—Sicilia Sanitaria.

LATTERI, S., 1952.-" Gli interventi demolitori nell'echinococco del polmone." 5 (5).

SAVINO, L., 1952.-" Contributo clinico sulla cisti di echinococco della tiroide." 5 (6), *b. 297-308.

953—Sovetskava Meditsina.

a. MAKHMUDOVA, B. A., 1952.—[Pathogenesis of eosinophilia in the early stages of ascariasis.] Year 1952, No. 8, pp. 23-25. [In Russian.]
b. KAMALOV, N. G. & TSUTSUNAVA, M. N., 1952.—[A clinical symptom of ancylostomiasis.] Year 1952, No. 10, pp. 27-28. [In Russian.]
c. OVNATANYAN, K. T., 1952.—[The problem of ascariasis of the heart and blood vessels.] Year 1952, No. 10, pp. 28-31. [In Russian.]
d. DEMIN, A. A. & SUMAROKOV, A. V., 1952.—[Electrocardiography in echinococcus infection of the heart.] Year 1952, No. 10, pp. 31-35. [In Russian.]

954—Srpski Arhiv za Tselokupno Lekarstvo. Belgrade.

STOJANOVIĆ, V. & ZOGOVIĆ, B., 1952.—[Primary bilocular echinococcosis of the anterior mediastinum.] 80 (1), 71-76. [In Serbian: French summary p. 76.]

955-Svenska Läkartidningen.

LUNDBERG, U., 1952.—" En ny form for behandling av oxyuriasis." 49 (45), 2759-2771.

(955a) Lundberg gives a history of the treatment of enterobiasis from the time of Hippocrates to modern times. It has been shown that the females of Enterobius vermicularis live from 37 to 101 days. In spite of that, treatment during eight to ten days and repeating the treatment after ten days is the best from a practical point of view. Lundberg has obtained good results by giving gentian violet and lignum quassiae as suppositories. The composition of these is given.

956—Terapevticheski Arkhiv.

a. OVNATANYAN, K. T., 1952.—[Problem of pancreatic ascariasis.] 24 (4), 73-78. [In Russian.]

957—Terre Marocaine.

*a. RANOUIL, 1952.—" Le téniasis du chien. Ses dangers pour l'homme et les animaux." 26 (274), 327-330.

958—Tidsskrift for den Norske Laegeforening.

KOLSTAD, P., 1952.—"Ascaris og akutt abdomen. To kasus. Diskusjon." 72 (13/14), 431-432. [English summary p. 432.]

959-Tórax. Montevideo.

a. ARMAND UGÓN, C. V., 1952.—" Tratamiento del quiste hidático del pulmón y de sus complicaciones." 1 (1), 83-104.
b. LARGHERO, P., 1952.—" Equinococosis costal." 1 (2), 199-222.
c. PURRIEL, P., MURAS, O., TOMALINO, D. & MENDOZA, D., 1952.—" Equinococosis cardíaca. Siembra metastática pulmonar." 1 (2), 223-234.
d. RAVERA, J. J., 1952.—" Equinococosis heterotópica pleural con neumotórax espontáneo. Hidátide vagabunda pleural con neumotórax espontáneo hidática (forma seca, sin derrame)." 1 (3), 348-353.

960-Transactions of the American Neurological Association.

a. CABIESES, F. & RAVENS, J. R., 1952.—" Cysticercosis of the central nervous system." 77th Annual Meeting (1952), pp. 76-79.
b. ARANA INIGUEZ, R., RODRÍGUEZ BARRIOS, R. & SAN JULIÁN, J., 1952.—" Surgical

treatment of hydatid cysts of the brain." 77th Annual Meeting (1952), pp. 210-211.
TRELLES, J. O. & ROCCA, E. D., 1952.—" Thirty-five verified cases of cysticercosis of the brain." 77th Annual Meeting (1952), pp. 263-267.

SKRYABIN, K. I., 1952.—[Problems of helminthological science in connection with Stalin's plans to reorganize nature in the SSSR.] 6, 27-35. [In Russian.] PODYAPOLSKAYA, V. P., 1952.—[The inculcation of Pavlov's teachings in the development of helminthology.] 6, 36-51. [In Russian.] SHIKHOBALOVA, N. P. & PRASOLOVA, M. A., 1952.—[Experimental investigation of immunity against Trichinella. I. The development of Trichinella in heavy and light infections in experimental animals.] 6, 25, 26. [In Processor.]

in experimental animals.] 6, 52-59. [In Russian.]
SHIKHOBALOVA, N. P., 1952.—[Experimental investigation of immunity against Trichinella.
II. Immunity acquired as a result of the disease.] 6, 60-71. [In Russian.]
SPASSKI, A. A., 1952.—[An investigation of the life-histories of anisakids (Ascaridata: Anisakidae).] 6, 72-73. [In Russian.]
SPASSKI, A. A., 1952.—[On the nomenclature of the genus *Diorchis* (Cestoda: Hymenolepididae).] 6, 74-75. [In Russian.]
SPASSKI, A. A., 1952.—[On the systematic position of the Hymenolepididae with a double

SPASSKI, A. A., 1952.—[On the systematic position of the Hymenolepididae with a double row of hooks on the scolex.] 6, 76-78. [In Russian.]

- (961c) The growth of Trichinella spiralis in the intestine is not the same in different hosts. In the guinea-pig, the adult is generally larger than in the rat, mouse, rabbit, dog etc. Tabulations of investigations on white mice show (i) the dependence of the percentage of trichinae developing in the intestine on the number of larvae in the infecting dose, (ii) the proportion of larvae in the infective dose to those reaching the muscles, (iii) the ratio of males to females at various periods of infection and (iv) the sizes of the trichinae in the intestine in slight and intense infections. G.I.P.
- (961d) When mice are superinfected with Trichinella spiralis on the 12th, 30th and 60th days after a primary infection, the worms in the intestine are fewer in numbers, smaller in size and leave the host sooner, but no marked shortening of the life of the adults was observed in immunized mice. The number of Trichinella larvae which encyst in the muscles of immunized mice after superinfection is much smaller than in controls and is correlated with the smaller number of worms developing in the intestine. A slight infection with as few as 50 larvae can produce immunity. The immunity resulting from superinfections after 12 or 30 days is more intense than from superinfection on the 60th day.
- (961e) The discovery of anisakid larvae in the subcutaneous tissues of shrews, fish etc. indicates that although some species of Porrocaecum may be acquired by birds from invertebrates, others are acquired from vertebrates. Spasski questions whether these infected shrews should be considered primary intermediate hosts or adventitious reservoir hosts. As he has found over 100 larvae in shrews and other instances have been reported from North America, he is of the opinion that these subcutaneous larvae are not pseudoparasites. G.I.P.
- (961f) The genus Diorchis Clerc, 1903 is now subdivided into two subgenera: (i) Diorchis Clerc, 1903 (syn. Nudiorchis Matevosyan, 1941) with type species D. (D.) acuminata (Clerc, 1902) (syn. Drepanidotaenia acuminata Clerc, 1902 and Diorchis (Nudiorchis) acuminata (Clerc, 1902) Matevosyan, 1941) and (ii) Acanthodiorchis n.subg. (syn. Diorchis Matevosyan, 1941) with type species Diorchis (Acanthodiorchis) skrjabini Udintsev, 1937. G.I.P.
- (961g) Among the avian cestodes, there is a small group with three testes in each segment, with unilateral genital pores and with two rows of hooks on the scolex. They approach the family Hymenolepididae but there are also about 300 species of Hymenolepididae with three testes in each segment and only one row of hooks on the scolex, which in this respect approach the Dilepididae. The combination of the most important characters of these two groups in a number of species listed has introduced discord in the determination of their systematic position. Meggittiella is transferred to the Dilepidinae. G.I.P.

SPASSKI, A. A. & ROMANOVA, N. P., 1952.—[On the morphology of Soboliphyme (Dioctophymata: Soboliphymidae).] 6, 79-84. [In Russian.] SPASSKI, A. A., RIZHIKOV, K. M. & SUDARIKOV, V. E., 1952.—[The helminth fauna

of wild mammals in the region of Lake Baikal.] 6, 85-113. [In Russian.]

j. MOZGOVOI, A. A., 1952.—[The biology of Porrocaecum crassum, nematodes of wild aquatic birds.] 6, 114-125. [In Russian.]
 k. MOZGOVOI, A. A., 1952.—[The biology of ascarids.] 6, 126-130. [In Russian.]

(961h) Soboliphyme baturini Petrov, 1930, common in cats, dogs, foxes etc., is now recorded for the first time from Martes (Lamprogale) flavigula. S. soricis, reported once from Scotland, has been found in Sorex araneus. Spasski & Romanova consider that the sucker-like mouth capsule is an important taxonomic feature of Soboliphyme and they therefore propose the retention of Petrov's Soboliphymidae, suppressed by Baylis & King, in an emended form. A third species, S. hirudiniformis, was described by Kirschenblatt in 1946 from Talpa coeca orientalis.

G.I.P.

(961i) Brief systematic notes are given of the helminths found in 21 species of wild mammals from around Lake Baikal. The parasites are also listed under their respective hosts. An illustrated account is given of a number of the species and of Syphacia sp. from Sciurus vulgaris and of Rictularia baicalensis n.sp. from Mus musculus and Apodemus agrarius. This new species resembles R. coloradensis in that the mouth capsule opens terminally and they both have the same number of combs in a row, but in R. baicalensis the spicules are equal and the egg measures 0.037-0.042 mm. ×0.029-0.033 mm. whereas in R. coloradensis the spicules are unequal and the egg measures 0.033 mm. × 0.022 mm. Both differ from R. harrisi in which the mouth capsule opens dorsally.

G.I.P.

(961j) The eggs of Porrocaecum crassum, a parasite of aquatic birds and particularly of ducks, develop to the infective stage in 5 to 12 days in water. In southern Russia, the infective eggs are taken up by Haematopota larvae. The nematode larvae then hatch, migrate into the blood sinuses and after about 24 hours accumulate in the posterior third of the body. When the infected insect larvae are eaten by ducks or other aquatic birds the nematode larvae first invade the stomach mucosa but afterwards attain sexual maturity in the small intestine.

G.I.P.

(961k) Four types of development are recognized in the Ascaridata, viz., (i) Ascaridioid, typified by Ascaridia galli in which the larvae do not migrate via the blood system of the host, (ii) Ascarioid, typified by Ascaris in which the larvae have a hepato-pulmonary migration, (iii) Toxocaroid, typified by Toxocara and Neoascaris, in which the larvae migrate from the pregnant animal into the foetus via the placenta and become adult shortly after its birth, and (iv) Anisakoid, typified by Contracaecum and Pseudanisakis in which larval development occurs in an intermediate host which is either planktonic or is a predatory fish. As Mozgovoi has recently shown, Porrocaecum crassum also requires an intermediate host. He recalls that Skryabin has divided the Ascaridata into two groups, viz., geohelminthic (Ascaridioid and Ascarioid types) and biohelminthic (Anisakoid types). The Toxocaroid type occupies an intermediate position between Ascaridioid and Ascarioid types on the one hand and Ascaridioid and Anisakoid types on the other. R.T.L.

G.I.P.

RIZHIKOV, K. M., 1952.—[New data on the Syngamidae.] 6, 131–138. [In Russian.] RIZHIKOV, K. M., 1952.—[Reservoirs of *Physocephalus sexalatus* (Molin, 1860)—nematodes of pigs.] 6, 139–141. [In Russian.] RIZHIKOV, K. M., CHERTKOVA, A. N. & VEITSMAN, L. N., 1952.—[On the helminth fauna of domestic guinea-fowl.] 6, 142–151. [In Russian.] SUDARIKOV, V. E. & RIZHIKOV, K. M., 1952.—[Description of a new family of nematodes from fresh-water fish (Spirurate: Haplonearidae a form.)] 6, 162–167. [In Russian.]

from fresh-water fish (Spirurata: Haplonematidae n.fam.).] 6, 152-157. [In Russian.] SUDARIKOV, V. E., 1952.—[Some results of the study of the helminth fauna of vertebrates in Gorkovskoye district (from material collected on the 63rd and 79th Soviet Helminthological Expeditions).] 6, 158-174. [In Russian.]
GUSHANSKAYA, L. K., 1952.—[The helminth fauna of wild birds of the SSSR.] 6, 175-

222. [In Russian.]

- (9611) Rizhikov describes and figures Syngamus (Ornithogamus) arcticus n.sp. from Gavia stellata. The ovum, unlike that of other species of Syngamus, has no operculum. The spicules are only 0.051 mm. long whereas in the other species of the subgenus Ornithogamus they are twice this length. Although Syngamus parvus has similarly short spicules, its other characteristics have placed it in the subgenus Syngamus. Syngamus auris Faust & Tang, 1934 is transferred to Mammomonogamus, and Syngamus hexadontus Chin, 1950 is placed in the R.T.L. subgenus Ornithogamus. G.I.P.
- (961m) It has already been shown that various mammals and birds can be infected, naturally and experimentally, with the infective larvae of Physocephalus sexalatus. It appears that all classes of vertebrates can act as hosts for the infective stage, for Rizhikov has now succeeded in infecting experimentally fish (Carassius carassius and Misgurnus fossilis), amphibians (Rana temporaria and Bombina bombina) and reptiles (Testudo horsfieldii). R.T.L. G.I.P.
- (961n) Guinea-fowl obtained from the zoological gardens in Moscow and from farms in the surrounding country harboured Heterakis gallinae, Capillaria columbae, C. caudinflata, Thominx collaris and Hymenolepis sp. C. columbae and Hymenolepis sp. are new records for this host. As guinea-fowl were also infected experimentally with Syngamus skrjabinomorpha helminth parasites of the guinea-fowl now number 50 species. R.T.L. G.I.P.
- (961 o) Haplonematidae, a new family of Thelazioidea, is created for the genera Haplonema, Ichtyobronema and Cottocomephoronema. Ichtyobronema conoura (Linstow, 1885) Gnedina & Savina, 1930 is renamed I. gnedini nom.nov. and Haplonema hamulatum Moulton, 1931 becomes Cottocomephoronema hamulatum (Moulton, 1931) n.comb. R.T.L. G.I.P.
- (961p) During the 79th Soviet Helminthological Expedition in the Gorkovskoye district 1,324 vertebrates were examined in addition to 1,194 domestic animals in slaughterhouses: 217 helminth species were collected. The numbers of trematodes, cestodes, nematodes and acanthocephalans found are tabulated under the various vertebrate classes. They are also considered from the point of view of their dependence on (i) the type of food of the host, (ii) the host locality, (iii) the age of the host, (iv) seasonal changes and (v) the physiological condition of their hosts. G.I.P.
- (961q) Gushanskaya annotates 101 species of helminths of wild birds in Russia. She also classifies them under their hosts. One new species is described and figured, viz., Corrigia viktori n.sp. in Coturnix coturnix. It differs from Corrigia corrigia in having oral and ventral suckers of equal size and an ovary larger than the testes. In the latter feature it also differs from C. skrjabini. The ovary is round whereas in C. plesiostomum it is lobed. G.I.P.

GUSHANSKAYA, L. K., 1952.—[A new subfamily Gynaecotylinae n.subf. of the family Microphallidae (Trematoda).] 6, 223-224. [In Russian.]
GUSHANSKAYA, L. K. & KROTOV, A. I., 1952.—[The discovery of the male of Schistorophus: skrjabini (Nematoda: Schistorophidae).] 6, 225-228. [In Russian.]
KASIMOV, G. B., 1952.—[A new nematode from a Trans-Caucasian partridge.] 6, 229-231.

KASIMOV, G. B., 1952.—[Skrjabinus popovi n.sp., a new trematode from Tetraogallus: caucasicus.] 6, 232-234. [In Russian.] DELYAMURE, S. L., 1952.—[Zoogeographical characteristics of the helminth fauna of pinnipeds and cetaceans.] 6, 235-250. [In Russian.]

(961r) Gynaecotylinae, a new subfamily of Microphallidae, is created to contain Gynaecotyla Yamaguti, 1939 as type and Diacetabulum Belopolskaya, 1952, and is characterized l R.T.L. particularly by the presence of two ventral suckers. G.I.P. .

(961s) The male of Schistorophus skrjabini (Vasilkova, 1926) Gushanskaya, 1950 is now described and figured for the first time. It is 8.27 mm. to 8.46 mm. in length. The tail carries cuticular alae. There are 22 pairs of pre-anal and six pairs of post-anal papillae. All I the papillae are pedunculate except for two post-anal pairs which are sessile. The spicules; are unequal: the right is 0.572 mm. long, is narrower than the left and has a triangular projection near its distal end; the left spicule is 0.087 mm., of peculiar shape and acts as a l gubernaculum. Its proximal end is 0.042 mm. wide and the distal end is 0.021 mm. wide.

G.I.P.

The male of Lemdana skrjabini n.sp., a filariid worm from Francolinus orientalis: caucasica, is 27.5 mm. to 29.1 mm. and the female 42.9 mm. to 54.2 mm. in length. The cuticle is striated. The head carries one pair of small lateral and two pairs of submedian papillae. The oesophagus is divided into two sections; in the male the length of the anterior is 0.379 mm. and the posterior 0.951 mm.; in the female the anterior is 0.33 mm. to 0.465 mm. and the posterior 1-127 mm. The excretory pore is 0-165 mm. from the anterior end in the male. The spicules are yellowish in colour, unequal in size and shape and ventrally curved. The larger spicule is 0.297 mm. long and is pointed; the smaller spicule is 0.165 mm. and is blunt. There is no gubernaculum. The male tail has two indistinct papillae. In the female: the genital opening is 3.48 mm. to 3.72 mm. from the head end and the tail carries a single papilla at its tip. R.T.L.

G.I.P.

(961u) In Skrjabinus popovi n.sp. from Tetraogallus caucasicus the vitellaria and the testes lie behind the ventral sucker, whereas in S. latus the vitellaria reach the anterior edge: of the ventral sucker and the right testis is on a level with its centre. The vitelline follicles are large; in S. biliosus they are scarcely visible. The ovary lies behind the right testis; in S. lanceatus it is somewhat to the left of the middle line and the vitellaria reach only to the centre of the ventral sucker. In S. similis the testes lie at the sides of the ventral sucker and in S. rarus they are level with its anterior edge. In S. skrjabini the ventral sucker lies between the testes and these are smaller than the ovary, whereas in S. popovi the testes and ovary are about equal in size. R.T.L.

G.I.P.

(961v) Delyamure has investigated the adult and larval helminths of pinnipeds and of cetaceans living in five zoo-geographical regions of the Pacific Ocean, sets out in tabular form the distribution of the 157 species known to be parasitic in these two groups and discusses the underlying general principles.

G.I.P.

G.I.P.

961-Trudi Gelmintologicheskoi Laboratorii. Akademii Nauk SSSR. (cont.)

w. DELYAMURE, S. L., 1952.—[Dependence of the helminth fauna of dolphins of the Black Sea and Sea of Azov on oecological and geographical factors.] 6, 251–258. [In Russian.] KROTOV, A. I., 1952.—[New cestodes (Hymenolepididae and Paruterinidae) from birds.]

6, 259-272. [In Russian.]
KROTOV, A. I., 1952.—[Nematodes of birds from the lower districts of the Amu estuary.]
6, 273-277. [In Russian.]
KROTOV, A. I. & DELYAMURE, S. L., 1952.—[On the helminth fauna of mammals and birds of the SSSR.]
6, 278-292. [In Russian.]

SOBOLEV, A. A., 1952.—[Skrjabinoclava longifuniculata n.sp., a new nematode from birds.] 6, 293-295. [In Russian.]

(961w) Following on his inquiry into the influence of zoo-geographical factors on the distribution of the helminth fauna of marine mammals in the Pacific Ocean (see No. 961v above), Delyamure has made an analysis of the helminths of dolphins, comparing those found in dolphins of the Black Sea with those found in the same host from the Mediterranean and the Atlantic Ocean. He then uses helminthological data in a discussion of the taxonomic status of dolphins in the Black Sea and Sea of Azov (on which mammalogists disagree) and maintains that these data support the views of Tsalkin, Kleinenberg and Kirpichnikov on the taxonomy of the Black Sea Phocaena. G.I.P.

(961x) Of the 37 cestode species recorded by Krotov from birds, three are new. In Dicranotaenia guschanskoi n.sp. from Calidris minuta the ovary is compact and the proboscis has ten hooks which measure 0.089-0.092 mm. A table sets out the different hook sizes by which eight other species of Hymenolepaea can be distinguished. In D. skrjabinissima n.sp. from C. subminuta the proboscis has 68 to 70 hooks, differing from D. polyacantha which has only 60 to 62 hooks and strong circular musculature in the proboscis sheath. In Aploparaksis sachalinensis n.sp. from Capella solitaria japonica the cirrus is bulbous and armed; it is thus differentiated from other species of Aploparaksis.

(961y) Krotov lists and gives brief notes on 14 known and seven unidentified species of nematodes collected from 16 avian host species from the lower Amu estuary region. G.I.P.

(961z) Of 36 helminth species from 27 avian and mammalian hosts two were trematodes, 19 were cestodes, ten were nematodes and five were acanthocephalans. Two new forms are described. In the Campulinae Leucasiella mironovi n.g., n.sp. is described from Delphinapterus leucas. Leucasiella, which has triangular eggs, differs from Synthesium, Orthosplanchnus, Oschmarinella and Odhneriella for the cirrus is unarmed as in Lecithodesmus, Zalophotrema and Campula. But in Lecithodesmus the ovary is lobed and the vitelline follicles form oblong groups, in Zalophotrema the ovary is lobose and the eggs are not triangular in shape, while in Campula the gut has lateral branches and the vitellaria reach the pharynx whereas in Leucasiella they are weakly developed and do not extend forward beyond the middle of the anterior testis. Bolbosoma bobrovoi n.sp. from Callorhinus ursinus curilensis is similar to B. turbinella and B. nipponicum but there are only 14 to 16 longitudinal rows with four to five hooks in each row, the last hook of which is much smaller than the others. R.T.L. G.I.P.

(961ba) In Skrjabinoclava longifuniculata n.sp. from Charadrius sp. the male measures 2.4 mm. and the female 3.468 mm. It differs from S. solonitzini and other species of the genus by the greater prolongation of the cordons on the anterior end of the body. Their length is 160 μ in the male and 180 μ in the female whereas in S. solonitzini they respectively measure 46μ - 63μ and 58μ . Two rows of spines run down the body from below the cordons. The spicules are 0.248 mm. and 0.122 mm. long. G.I.P.

bb. SOBOLEV, A. A., 1952.—[Phylogenetic relationships and systematics of the Camallanata.] | 6, 296-301. [In Russian.] bc. FEDYUSHIN, A. V., 1952.—[Ascaridia skrjabini n.sp., a new nematode from Tetraogallus | himalayensis in Tian-Shan.] 6, 302-304. [In Russian.] bd. SCHULZ, R. S. & DAVTYAN, E. A., 1952.—[Latent helminthiases and their epizootic importance] 6, 205-204. [In Russian.]

tance.] 6, 305-314. [In Russian.] be. MACHULSKI, S. N., 1952.—[Two new genera and species of Spirurata from mammals.] 6, 315-322. [In Russian.]

(961bb) Sobolev agrees with Chitwood & Wehr in bringing Camallanidae and Cucullanidae together into the Camallanoidea Travassos, 1920 and associating this superfamily with Dracunculoidea Cameron, 1934 in Chitwood's Camallanata. Cucullanidae Cobbold, 1864 contains Seuratinae Hall, 1916 and Cucullaninae Yorke & Maplestone, 1926 which is now subdivided into two new tribes named Cucullaninea and Dichelininea [but not defined]. In Dracunculoidea there are now four families, viz., Dracunculidae Leiper, 1912 (Dracunculinae, Avioserpensinae, Philometrinae and Micropleurinae), Tetanonematidae Skryabin & Shikhobalova, 1948, Cystopsidae Skryabin, 1923 and Anguillicolidae Yamaguti, 1935.

(961bc) Ascaridia skrjabini n.sp. from Tetraogallus himalayensis differs from A. galli, A. compar, A. compressa, A. granulosa and A. styphlocerca by the smaller body (male 23 mm.-39 mm., female 35 mm.-60 mm.) and the shorter tail. There are ten genital papillae of which five are pre-anal. The distal end of the slightly unequal spicules is very characteristically swollen, like a pinhead. R.T.L.

G.I.P. (961bd) Schulz & Daytyan discuss the problem of latent helminthiasis in which helminths at various stages of growth remain inactive in the host. From experimental data they conclude that the latent state may extend over 11/2 years during which the larvae remain in the host without undergoing further development. Factors on which the occurrence of this latent stage depends are unfavourable conditions in the hosts, e.g. primary or secondary immunity, specific physiological states and the effect of the external environment. Factors which influence the return of activity are the loss or lowering of the host's immunity and unfavourable feeding and living conditions. The nature of latent helminthiasis explains the impracticability of revealing these infections by examination for ova or larvae. The diagnosis of latent helminthiasis will depend on the utilization of definite immunological reactions and special investigations should be made into the immuno-diagnosis of helminthiases at the different stages of their development, the duration of positive reactions, the degree of specificity of the reactions etc. The recognition of latent helminthiasis may assist in elucidating some mysterious outbreaks of helminthiases and lead to the discovery of new epizootic and epidemiological principles.

> R.T.L. G.I.P.

(961be) Two new genera, Petrowospirura and Skrjabinocercina are added to the Spirocercinae. Petrowospirura lynxi n.g., n.sp. from Lynx lynx and Otocolabus manul has the following characters: there are no lateral alae along the sides of the body, the mouth has six well developed bilobed lips, the chitinized mouth capsule is funnel-shaped and contains six teeth which do not extend beyond its anterior margin (of these five are very small and one is thin and broad); in the female the tail has no terminal papillae and the genital opening is in the anterior third of the body; in the male the gubernaculum is well formed and the spicules are very unequal. In Skrjabinocercina petrowi n.g., n.sp. from Alactaga saltator mongolica there are no lateral alae on the body, the mouth has four well developed lips each separated by a papilla, the funnel-shaped chitinized mouth capsule contains six teeth (of which the two laterals are larger than the subdorsal and subventral), the teeth are supported by ridges arising from the base of the capsule and they do not extend beyond the anterior margin of the mouth; in the male the genital papillae are pedunculated; in the female the genital opening is immediately in front of the anus and the tail has no terminal papillae. R.T.L.

bf. ROMANOV, I. V., 1952.—[New species of helminths from Martes zibellina.] 6, 323-330.

[In Russian.] bg. RUKHLYADEV, D. P., 1952.—[On the helminth fauna of the wild boar.] 6, 331-333. [In

bh. NALETOV, N. A., 1952.—[Pathological and morphological changes in the lungs of sheep caused by Cystocaulus infection.] 6, 334-337. [In Russian.]
bi. PARAMONOV, A. A., 1952.—[On the oecological classification of plant-parasitic nematodes.] 6, 338-369. [In Russian.]

(961bf) Taenia skrjabini n.sp. from Martes zibellina is closely related to T. intermedia Rudolphi, 1809 but has the following characteristics: the strobila is only 8-10 cm. long; the suckers are 0.138-0.168 mm. in diameter; the scolex is 0.634-0.840 mm. in diameter; there are hooks on the scolex, those of the first row measure 0.153-0.155 mm. and those of the second row 0.126-0.129 mm.; the cirrus pouch is 0.160-0.170 mm. in length and 0.060-0.168 mm. in breadth; the uterus has 14-15 lateral branches and the eggs are 0.027 mm. × 0.023 mm. Sobolevingylus petrovi n.g., n.sp. is described and figured from Martes zibellina and a new subfamily of Pseudaliidae is made, the Sobolevingylinae. The new genus has a weakly developed bursa, pre- and post-anal papillae and a dorsal lobe but lacks a gubernaculum. The dorsal ray has four terminals. The spicules are short and broaden at their proximal ends. The genital opening is in front of the anus, near the posterior end. R.T.L.

G.I.P.

(961bg) Rukhlyadev reports the occurrence of the following species from a wild boar in the Caucasian Reserve: Metastrongylus salmi, M. pudendotectus, M. elongatus, Physocephalus sexalatus and Rhabditata gen. et sp. inq. G.I.P.

(961bh) In cystocauliasis in sheep Naletov recognizes two types, viz., uncomplicated cystocauliasis pneumonia and cystocauliasis pneumonia with bacterial complications. He describes the consequent pathological changes in the lungs. R.T.L. G.I.P.

(961bi) Phytonematodes are defined as those which are parasitic in plants or have temporarily settled in their tissues. The Tylenchata are all considered to be parasitic and are divided into (i) pathogenic non-specific ecto- or endo-parasitic phytohelminths and (ii) pathogenic specific phytonematodes, monomorphic and dimorphic. The phytohelminths differ sharply from the phytonematodes in that both in vivo and when fixed they do not stain in cold solutions of methylene blue. The oecological characteristics of each group are compared with their taxonomic characteristics and a rearrangement of the species of Diplogaster is suggested. A new suborder, Diplogasterata, containing a number of new families and genera is proposed whereby the correlation of the oecology and taxonomy underlies their role in the phylogenetic development of natural systematic groups. The order Rhabditida is subdivided into Rhabditata Chitwood, 1933 ex parte, Tylenchata (Thorne, 1949) Chitwood, 1950 and Diplogasterata n.subordo. Diplogasterata contains (i) Diplogasteridae Steiner, 1919, (ii) Diplogasteroididae n.fam. (=Diplogasteroidinae Filipjev & Schuurmans Stekhoven, 1941), (iii) Odontopharingidae Micoletzky, 1922, (iv) Demaniellidae n.fam. (=Demaniellainae Paramonov, 1951), (v) Cylindrocorporidae Goodey, 1939 and (vi) Tylopharingidae n.fam. (=Tylopharinginae Filipjev & Schuurmans Stekhoven, 1941). Diplogasteridae is subdivided into Diplogasterinae Micoletzky, 1922 ex parte and Neodiplogasterinae n.subf. with Neodiplogaster tropica Cobb, 1924 as type. Diplogasterinae contains fourteen genera, viz., Diplogaster Schultze, 1857, type D. micans; Pristionchus Kreis, 1932, type P. lheritieri (Maupas, 1919) n.comb.; Butlerius Goodey, 1929, type B. butlerius; Fuchsia Micoletzky, 1922, type F. butschlii (Fuchs, 1915) and ten new genera: Diplogasteritus n.g., type D. nudicapitatus (Steiner, 1914) n.comb.; Diplogastrellus n.g., type D. gracilis (Bütschli, 1876) n.comb.; Oigolaimella n.g., type O. winchesi (Goodey, 1929) n.comb.; Paroigolaimella n.g., type P. coprophaga (de Man,

bj. SKARBILOVICH, T. S., 1952.—[A study of the species of the genus Hexatylus Goodey, 1926.]

6, 370-375. [In Russian.] bk. POLOZHENTSEV, P. A., 1952.—[New Mermithidae from the sandy soil of pine woods.]

6, 376-382. [In Russian.]
bl. VELICHKIN, P. A., 1952.—[Alfortia causing disease in horses maintained in droves.] [Abstract of thesis.] 6, 383-392. [In Russian.]
bm. LEVASHOV, M. M., 1952.—[The status of the helminth fauna of the SSSR and a study of the characteristics of helminths in oecological and geographical zones.] [Abstract of thesis.]

6, 392-396. [In Russian.] bn. MOROZOV, F. N., 1952.—[Heterophyoidea of man, domestic and wild animals.] [Abstract of thesis.] 6, 397-400. [In Russian.]

1876) n.comb.; Fictor n.g., type F. vorax (Goodey, 1929) n.comb.; Paradiplogaster n.g., type P. similis (Bütschli, 1876) n.comb.; Eudiplogaster n.g., type E. striatus (Bütschli, 1876) n.comb.; Pareudiplogaster n.g., type P. striatulus (Fuchs, 1933) n.comb.; Anchidiplogaster n.g., type A. dubius (Hnatewytsch, 1929) n.comb.; Diplogasterium n.g., type D. micoletzkyi (Hnatewytsch, 1929) n.comb. The formation of these new genera has necessitated the creation of numerous new combinations for species transferred from the original genus Diplogaster. Paroigolaimella anchicoprophaga is proposed for D. coprophagus of Goodey, 1929. G.I.P.

(961bj) Hexatylus vigissi n.sp. is found on the roots, stems and tubers of potatoes and on the weeds Atriplex hortensis, Galium sp., Sonchus arvensis, Plantago major, Polygonum tataricum, Centaurea sp., Equisetum pratense, Cuscuta, Barbaraea, Geranium, Matricaria, Malva and Poa. It differs from H. viviparus in having a smooth cuticle and three basal swellings on the stylet. The anterior portion of the oesophagus is cylindrical. The tail is conical and pointed whereas that of H. coprophagus is thread-like. Hexatylinae n.subf. contains Hexatylus and Deladenus. R.T.L.

G.I.P.

(961bk) Polozhentsev describes three new Mermithinae living in sandy soil of pine woods. The new species are morphologically similar to those of the aquatic genus Bathymermis but, as a follower of the teaching of Michurin and Lysenko, he finds it impossible to unite terrestrial and aquatic species in a single genus and accordingly creates for these new terrestrial species a new genus named Skrjabinomermis. In S. tolski n.g., n.sp. the male is 40·3-43·5 mm. long, the arched spicules are 0.925-1.062 mm. in length, the amphids are 0.013-0.024 mm. × 0.005-0.007 mm. and the excretory pore opens at the level of the posterior end of the oesophagus. In S. sukatschevi n.sp. the male is 47.5-53.4 mm. long, the coiled spicules measure 1.860 mm., the amphids are 0.015 mm. ×0.005 mm. and the excretory pore opens near the beginning of the intestine. Psammomermis busuluk n.sp. occurs in soil pits and is the smallest species of the genus. The male measures 11-13 mm., the female 32-58 mm.; the cuticle is smooth, the spicules are 0.567 mm. long and are twice the length of those of P. korsakowi; the vagina is somewhat curved. The tails in both sexes are longer than in other species.

G.I.P.

(961bl) This thesis concerns the diagnosis of Alfortia disease in horses, the clinical symptoms, pathology and histology of intestinal and peritoneal infections and the influence of Alfortia on the course of salmonelliasis (Salmonella abortus equi) in young horses under experimental and natural conditions. R.T.L.

G.I.P.

(961bn) Morozov's thesis reviews the morphology and geographical distribution of the 116 known species of Heterophyoidea. In a new classification of this superfamily, Euryhelminae n.subf. [this also appeared in a paper by the same author in 1950, for abstract see Helm. Abs., 19, No. 308f] for Euryhelmis Poche, 1925 is added to the five subfamilies of Heterophyidae, and Haplorchoidinae n.subf. to the three subfamilies of Cryptogonimidae Ciurea, 1933.

bo. BUTORIN, F. S., 1952.—[Dynamics of the principal helminthiases and an analysis of measures for their control in farm animals in the Yaroslavl region.] [Abstract of thesis.] 6, 400-403.

[In Russian.] bp. GAGARIN, V. G., 1952.—[Capillariasis of domestic birds.] [Abstract of thesis.] 6, 403-406.

[In Russian.] bq. GEFTER, V. A., 1952.—[Experimental study of the role played by reservoirs in the epidemiology of ascariasis.] [Abstract of thesis.] 6, 406-407. [In Russian.] br. ZASKIND, L. N., 1952.—[Helminthiases of geese and their causal agents.] [Abstract of thesis.]

6, 407-409. [In Russian.]
KANGINA, K. I., 1952.—[Nitrogen metabolism and amino acid composition of the protein of intestinal helminths. (Ascaris suum and Moniezia expansa.)] [Abstract of thesis.] 6, 409-411.

OLIGER, I. M., 1952.—[Parasite fauna of woodcocks of forest zones of the European part of

RSFSR.] [Abstract of thesis.] 6, 411-412. [In Russian.]

Cryptogoniminae Ward, 1917 is subdivided into two new tribes, Cryptogonimea for Cryptogonimus and Caecincola and Neochasmea for Neochasmus and Allacanthochasmus. Siphoderinae Manter, 1934 is also subdivided into two new tribes, Siphoderea for Siphodera, Siphoderina and Centrovarium, and Paracryptogonimea for Paracryptogonimus, Biovarium and Siphoderoides. [The abstract of the thesis does not give any definitions of the new taxonomic groups but the author contributes these to "Trematodes of Animals and Man" R.T.I.. Volume 6. For abstract see No. 1000 below.] G.I.P.

(961bp) This abstract of a thesis states that it deals with the distribution of Capillaria disease in birds in Russia where 29 species of Capillaria are known to occur; their morphology, biology, epizootology, classification and prophylaxis are reviewed. The most widespread and pathogenic are C. columbae and C. caudinflata in poultry, Eucoleus contorta in ducks and C. anseris in geese. A key is provided for the differentiation of the genera Eucoleus, Thominx and Capillaria. G.I.P.

(961bq) This thesis deals with the significance of a large river in a large sewered town, the various stages of pollution with helminth eggs, the process of self purification and the survival time of Ascaris lumbricoides eggs. The relevant literature is analysed. R.T.L.

(961br) This abstract of an unpublished thesis states that Soviet and other literature on the helminths of wild and domestic geese is collated, that it also contains a review of their systematics, biology, diagnosis, therapy and prophylaxis, and original descriptions of several species [not mentioned in the abstract]. Both Capillaria anatis and C. anseris have been found in the domestic goose which is now recorded as a new host for Apatemon gracilis. R.T.L. G.I.P.

(961bt) Oliger gives a description of the parasitic fauna of woodcocks of the forest zones of European RFSSR and discusses their role in the fluctuation of the numbers of birds. In the 531 birds examined there were 38 species of parasites of which six are stated to be new and of these four are new helminths. They are named Corrigia bonasia n.sp., Choanotaenia sylvarum n.sp. and Leucochloridium tetrastae n.sp. from the hazel grouse, and Trichostrongylus medius R.T.L. n.sp. from the woodcock. G.I.P.

962-Trudi Zoologicheskogo Instituta. Akademii Nauk SSSR.

POLYANSKI, Y. I., 1952.—[Some new and little known parasitic nematodes from the intestines of marine fish.] 12, 133-147. [In Russian.]

(962a) From fish from the Berents Sea, Polyanski gives a detailed description of three new species and of the males and females of Ascarophis morrhuae Van Beneden, 1871. A. arctica

n.sp., from the intestine of Zoarces viviparus, differs from A. morrhuae in having less developed cuticular spines, eggs with two polar plugs and numerous filaments, and in the shape of the male spicule which is expanded and indented. A. filiformis n.sp., from the stomach of Gadus morrhua, is differentiated from the two above species by its larger measurements, its smaller male spicule and the absence of cuticular striations. The ovum, like that of A. morrhuae, has one polar plug with two filaments. Capillaria salvelini n.sp., from the intestine of Salvelinus albinus, is separated from all other species of this genus except C. coregoni by its spicule which is proximally thickened and surrounded by a coiled, spring-like structure; C. coregoni is smaller, has smaller eggs and a different relative length of the oesophagus and the postoesophageal part. Polyanski finds great similarity between Ascarophis (misquoted by Nicoll, 1907, as "Ascaropsis") and Capillospirura Skryabin, 1924 and therefore considers Capillospirura to be a synonym of Ascarophis.

963—Türk Veteriner Hekimleri Derneği Dergisi.

TINAZ, A., 1952.—"Kalbin sol atriumu üzerinde ve sol boyun adaleleri arasında Genurus cerebralis." 22 (64/65), 27–30. [English summary p. 30.]

MIMIOGLU, M., 1952.—"Ankara'nın sokak kedileri üzerinde yaptığımız bir araştırmada müşahede ettiğimiz bir Opisthorchis felineus Rivolta, 1885 (karacığer kelebeği) olayı." 22 (66/67),

75-79. [English & German summaries pp. 77-78.]
OYTUN, H. S., 1952.—" Saint-Denis Kimya Fabrikasının phenothiazine ilâciyle, memleketimiz ceșitli evcil hayvanlarında, yaptığımız deneylerin sonuçları." 22 (68/69), 133-139. [English

summary p. 139.] GEDIZ, C., KUMOVA, C. & ACAREL, M., 1952.—"Bir katırın kamera okuli anteryörü hümör aközünde Setaria equina Abildgaard 1789." 22 (70/71), 199–204. [French & German summaries pp. 203-204.]

964-Věstník Ceskoslovenské Zoologické Společnosti.

ERHARDOVÁ, B. & RYŠAVÝ, B., 1952.—" Skrjabinema ovis, Vereščagin, 1926 (Oxyurata, Syphaciinae)." 16 (1/2), 195-196. [German & Russian summaries pp. 195-196.] VAVROUŠKOVÁ, K., 1952.—" Barvoměna pijavky Protoclepsis tesselata (O.F. Müller)." 16 (3/4), 334-353. [German & Russian summaries pp. 348-352.]

(964a) The occurrence of Skrjabinema ovis in goats in the neighbourhood of Roznava is briefly reported.

(964b) The colouration of *Protoclepsis tesselata*, in addition to the colouration of the connective tissue, is due to four types of chromatophore cells with processes in which pigment granules move centrifugally or centripetally. There are smaller chromatophores with redbrown pigment in the subepidermal network. The segment spots are formed by small groups of orange-yellow chromatophores situated somewhat deeper with processes which form a fine subepithelial network. There are larger chromatophores deeper in the connective tissue; these are red-brown in colour and still deeper there are green chromatophores. In the light the leech is usually green. In the dark it is light and transparent. The colour change depends on the concentration and dispersal of the pigment. The dispersal which causes a darkening in colour is probably a primary reaction of the chromatophores acting as photoreceptors. With an increase in light intensity the pigment granules pass from the centre to the processes of the chromatophore cells. Sodium chloride and alkaloids cause the pigment to concentrate. As extracts of lightly coloured or of dark leeches have no effect it is unlikely that the body fluid influences the concentration or dispersal of the pigments.

965-Veteriner Fakültesi Yayınları. Ankara Universitesi.

AKMAN, S., 1952.—" Tek tırnaklı hayvanların bağırsaklarında yaşayan askaritlere karşı en müessir ilâcın araştırılması." 21, Çalişmalar 5, 131 pp. [German summary pp. 128-131.]

(965a) Akman gives his observations on the results of treating small and varying numbers of equines with ten different anthelmintics (including carbon disulphide, carbon tetrachloride, chenopodium oil and santonin) for ascarid infection.

966-Western Fruit Grower.

*a. ALLEN, M. W., 1952.—" Nematode parasites of trees and vines." 6 (2), 33-37.

967-Wissenschaftliche Zeitschrift der Martin-Luther-Universität Halle-Wittenberg.

*a. HARTWICH, G., 1952.—"Vergleichende mikroskopisch-anatomische Untersuchungen über den Kopfbau einiger Ascariden." Mathematisch-naturwissenschaftliche Reihe No. 3, 1 (4), 71-83.

968-Zdravstveni Vestnik.

a. VODOPIVEC, S., 1952.—"Razširjenost in zatiranje trakuljavosti v LRS." [Distribution and control of tapeworm infection in Slovenia.] 21 (11/12), 223-228.

969-Zentralblatt für Chirurgie.

a. ERHART, K. H., 1952.—" Verstopfung einer Duodenalsonde durch einen Spulwurm."

77 (48), 2388-2389. b. ZYLKA, N., 1952.—"Pathologische Veränderungen der Mesenteriallymphknoten bei Askaridiasis." 77 (50), 2442-2450.

NON-PERIODICAL LITERATURE

970—AMELUNG, H., 1952.—"Untersuchungen über die Verteilung der Muskeltrichinellen im Tierkörper sowie über die Invasionsfähigkeit jugendlicher Trichinellen." Dissertation, Hanover, 36 pp.

Amelung's experiments with guinea-pigs show that there is some correlation between the number of Trichinella larvae ingested and the number of larvae found in the muscles. The first larvae were found in the muscles twelve days after infection and larvae were found in the heart up to at least the 25th day. Coiled larvae were present after 18 days; encapsulation was completed after about the 60th day. Larvae can be infective from the 15th day and infection was obtained in 60% of guinea-pigs with larvae not more than 25 days old. In Amelung's view all Trichinella larvae, from the very youngest infective stage, can be identified with certainty by the trichinoscope.

- *971—BENBROOK, E. A., 1952.—" List of parasites of domesticated animals in North America." Minneapolis: Burgess Publishing Co., 2nd edit. (revised), 63 pp.
- *972—BIKHOVSKAYA-PAVLOVSKAYA, I. E., 1952.—[The parasitological study of fishes.]

 Moscow: Izdatelstvo Akademii Nauk SSSR, 62 pp. [In Russian.]
- 973—BINGEFORS, S., 1952.—" Nematode resistance in clover and lucerne." International Grassland Congress (6th), State College, Pennsylvania, 1952. Proceedings, Vol. 2, pp. 1591–1596.

A short review of earlier work on breeding for resistance to Ditylenchus dipsaci in clover and lucerne is given. Breeding has resulted in such strains as Merkur red clover in southern Sweden, F.A.V. San Martin lucerne in Argentina, and Nemastan lucerne in the U.S.A. Investigations in Sweden have shown that it is possible to obtain a rapid improvement in resistance in local adapted strains of red clover by selection, and that crossing with resistant but not adapted strains has also given good results. An artificial infection method has been worked out. The most important cause of the resistance of certain plants seems to be the repression of development of the nematodes in such plants. The inheritance of resistance has also been studied.

- *974—BLAISDELL, K. F., 1952.—" A study of the cat lungworm Aelurostrongylus abstrusus." Thesis, Cornell University, 181 pp.
- *975—CARPENTER, M. F. P., 1952.—"The digestive enzymes of Ascaris lumbricoides var. suis: their properties and distribution in the alimentary canal." Dissertation, University of Michigan, 399 pp.

Carpenter has studied the enzymes in the gut of Ascaris lumbricoides from pigs and has demonstrated the presence of an amylase which is activated by chloride, a maltase, a proteinase and at least four peptidases; lipase and esterase were also demonstrable. All the enzymes showed the greatest activity in the anterior region of the gut. The properties of the Ascaris amylase, maltase and proteinase differed from those of the equivalent enzymes of the pig. In order of activity the enzymes are peptidases, proteinase, maltase, lipase, amylase. [Based on an abstract in Biol. Abstr., 27 (4), No. 11071.]

- *976—DESANLIS, J., 1952.—" Propagation des parasites et des germes infectieux par les pailles et les fourrages secs." Thesis, Alfort, 75 pp.
- 977—DIAS, C. B., 1952.—"A síndrome hépatico-esplênica na esquistossomíase mansônica. Contribuição ao seu estudo, baseada em 22 casos clínicos, dos quais 13 submetidos à esplenectomia." Thesis, Belo Horizonte, 449 pp. [English summary pp. 421-430.]

Dias has investigated 22 cases of schistosomiasis mansoni and deals with the pathological effects of the disease on the liver and spleen and the changes in the blood and in the liver function which followed early splenectomy in 13 cases.

R.T.L.

- *978—DÖLL, W., 1952.—" Die Bedeutung der *Pseudomonas hirudinis* für die Verdauungsphysiologie des medizinischen Blutegels." Dissertation, Marburg.
- *979—FREYTAG, K., 1952.—" Über die Bakterienflora der Haut und der Segmentalorgane von Hirudo officinalis und medicinalis." Dissertation, Marburg.
 - 980—FURCHNER, H. O., 1952.—"Über die Lebensfähigkeit von Askarideneiern in Hauskläranlagen." Dissertation, Munich, 45 pp.

Furchner reports on a series of experiments carried out to determine whether Ascaris ova remain viable and infective in septic tanks and other small sewage plants. Ova of Toxocara canis, Parascaris equorum, Ascaris lumbricoides (human and pig strains), and Neoascaris vitulorum, were used in the experiments but no differences in the behaviour of the varying species were noted. While the number of ova which embryonated after five months in sewage varied between 10% and 80%, practically all ova were destroyed by the 11th month. Furchner also reports on tests made on embryos from the sewage; he was unable to produce experimental infection in mice. He concludes that the use of sludge from septic tanks as garden manure is likely to spread Ascaris infection.

*981—GOODMAN, J. D., 1952.—" Taxonomic studies on the family Ochetosomatidae Leão, 1944, and the life history of *Stomatrema guberleti* Byrd, 1937. Trematoda." Dissertation, University of Michigan, 322 pp.

In this review of the Ochetosomatidae, Goodman replaces the Leptophyllinae with Travirematinae nom.nov. Macroderinae nom.nov. [n.subf.?] is established for Macrodera and Natriodera. N. verlatum Talbot, 1934 is transferred to N. variabilis. The synonymy of species of Lechriorchis, Neorenifer, Ochetosoma and Renifer is discussed. Dasymetra nicolli is transferred to D. natricis n.comb. Ochetosoma lampropelti n.sp., O. rauschi n.sp. and

Pneumatophilus tracheophilus n.sp. are listed [but not described in the abstract]. The first intermediaries of Stomatrema guberleti are Physa spp., the second intermediaries are Amphiuma and Siren and the definitive host is the mud-snake Farancia, the adult worms being found in the mouth and oesophagus. Details of the life-history are given. [Based on an abstract in Biol. Abstr., 27 (5), No. 14465.]

*982—HUNT, J. S., 1952.—" The life-history of Gorgodera vivata n.sp. (Trematoda: Gorgoderidae)." Dissertation, University of Michigan, 149 pp.

Hunt found more than 9% of 14,761 fingernail clams (Sphaerium simile) collected in the vicinity of Ann Arbor to be infected with cercariae of a new trematode, Gorgodera vivata n.sp. The larvae and nymphs of a number of aquatic insects became infected by ingesting the cercariae, and metacercariae developed. When the metacercariae were fed to frogs, Rana clamitans and R. catesbeiana became infected, sexually mature worms developing, but R. pipiens and R. palustris were refractory. [Based on an abstract in Biol. Abstr., 27 (5), No. 14466.] s.w.

983—INTERNATIONAL CONGRESS OF ENTOMOLOGY (9th), Amsterdam, August 17–24, 1951. Transactions.

THÉODORIDÈS, J., 1952.—" Contribution à l'étude écologique des parasites et commensaux de Coléoptères (2e note)." Vol. I, pp. 454-459. [Discussion p. 458.] GORDON, R. M., 1952.—" Problems in the transmission of filariasis." Vol. I, pp. 939-945.

[Discussion p. 945.] BERTRAM, D. S., 1952.—" Factors affecting the efficiency of the mite-vector of cotton rat filariasis." Vol. I, pp. 951-955. [Discussion p. 955.]

(983a) Théodoridès summarizes the different biological and oecological aspects of parasitism and commensalism associated with the Coleoptera and annotates and tabulates the various fungi, protozoa, nematodes, acarines and insects found. The degree of association ranges from simple phoresis to endoparasitism and depends to a large extent on the oecology and ethology of the host.

(983b) As in yellow fever and some rickettsial and spirochaetal diseases, certain forms of filariasis may be partially maintained by arthropods which rarely or never bite man but feed on other animals. Chimpanzees may act as reservoirs of Acanthocheilonema perstans and monkeys of Loa loa. Gordon briefly summarizes recent observations on the uptake of Loa loa by its vector, its development in the vector, the chances of its transfer to the vertebrate host and the deposition of the parasite by the vector. With one exception members of the research unit working in West Africa developed Calabar swellings. Dimethyl phthalate gave protection from Chrysops for two hours. In the discussion, Galliard stated that 50% of the aboriginal tribes (Mois) in Indo-China were infected with Wuchereria bancrofti but showed no apparent symptoms, whereas only 7% of the Indo-Chinese carried infections but clinical cases R.T.L. were numerous.

(983c) The magnitude of the infection and the multiplicity of reinfection of the cottonrat affect the readiness with which Bdellonyssus bacoti becomes infected by Litomosoides carinii. Environmental conditions prevailing during the infecting meal can affect considerably the R.T.L. ultimate infection rate in the mites.

984-KÄMPFE, L., 1952.-" Rüben- und Kartoffelälchen." Leipzig: Neue Brehm-Bücherei, No. 80, 42 pp. /

Kämpfe gives a full, illustrated account of the life-history, development and morphology of the sugar-beet and potato-root nematodes, describing the effects on their hosts. He deals briefly with control measures including chemical and biological control and crop rotation. 985—KELLER, H., 1952.—" Zusammenhang der Lebertuberkulose mit Leberegelbefall." Dissertation, Munich, 44 pp.

The incidence of liver-fluke among cattle slaughtered at Regensburg rose from 8.3% in 1945 to 20.9% in 1949; over the same period tuberculosis of the liver increased from 4.7% to 14.8%. In order to study the relationship between the two infections Keller examined 73 slaughtered cattle during 1951 and 1952. Of 41 animals infected with liver-fluke, seven had tuberculosis of the liver while only one animal out of 32 free from liver-fluke had tuberculosis of the liver. From these figures Keller concludes that liver-fluke infection in cattle favours the development of liver tuberculosis. He recommends further research on this problem.

A.E.F.

- *986—KOESLING, G., 1952.—" Verbreitung parasitärer Tierkrankheiten im Stadt- und Landkreis Braunschweig." Dissertation, Giessen, 35 pp.
- *987—KOLLIGS, A., 1952.—" Verbreitung parasitärer Tierkrankheiten im Kreise Geilenkirchen-Heinsberg." Dissertation, Giessen, 47 pp.
- 988—KRUEDENER, R. von, 1952.—"Über die Ursachen der Aufwärtswanderung von Bacterium coli bei Rindern mit Leberegelbefall." Dissertation, Munich, 56 pp.

Kruedener concludes from his researches that the primary cause of the upward migration of Bacterium coli to the duodenum in cattle infected with liver-fluke is the presence of the liver-fluke, although the symptoms are produced by the migrating bacteria. He considers that in other infections the position is reversed and that it is the presence of B. coli which causes favourable conditions for their development. Control of liver-fluke disease in cattle should reduce the incidence of B. coli infection.

A.E.F.

- *989—MÜNCH, K., 1952.—" Der Kapillarienbefall des Geflügels, insbesondere der Hühner." Dissertation, Giessen, 38 pp.
 - 990—MULZER, B., 1952.—" Über die Technik der intratrachealen Lungenwurmbehandlung beim Schaf unter besonderer Berücksichtigung der Flüssigkeitsverteilung in der Lunge und ein Hinweis auf die Aerosoltherapie." Dissertation, Munich, 37 pp.

After a detailed study of the technique of intratracheal injections against lungworms in sheep, Mulzer concludes that $88\%_{-}90\%$ of large lungworms are reached when injection is made with the sheep lying on its back on a board at an angle of 40° . If this injection is preceded by one in the sitting or standing position up to $95\%_{0}$ of the large lungworms should be reached. Mulzer does not consider intratracheal injections to be efficacious against small lungworms. The paper concludes with a note on the use of aerosols which are thought worthy of further study.

*991—NAJIM, A. T., 1952.—" Life-history of Gigantobilharzia huronensis Najim, 1950, a dermatitisproducing bird blood-fluke (Trematoda—Schistosomatidae)." Dissertation, University of Michigan, 99 pp.

Najim found *Physa gyrina* to be the intermediate host of *Gigantobilharzia huronensis*, and *Spinus tristis tristis* and *Richmondena cardinalis* to be natural definitive hosts. In the laboratory canaries and chicks were readily infected. The eggs are spheroid and spineless and hatch rapidly in water; the miracidium has 22 epidermal plates in four rows, with a wide gap between the second and third rows; the mother sporocyst is elongate and sac-like with a birth pore, and the anterior end of the daughter sporocyst is covered with spines; the cercaria is apharyngeate, spiny, with a bifurcate tail and pigmented eye-spots. Cercariae are shed as early as 24 days after infection of the snails and eggs appear in the faeces of the final host about 31 days after exposure to cercariae. The adult schistosomes are suckerless and filiform with the females considerably longer than the males. [Based on an abstract in *Biol. Abstr.*, 27 (5), No. 14469.]

- *992—PARAMONOV, A. A., 1952.—[Parasitic worms (helminths) of animals and plants and their control.] Moscow: Moskovskoe Obshchestvo Ispitatelei Prirodi, 111 pp. [In Russian.]
- *993—PARRISIUS, G., 1952.—"Der Endoparasitenbefall bei Ziegen in verschiedenen Gegenden unter Berücksichtigung der jeweiligen Umweltfaktoren." Dissertation, Berlin.
- *994—PAVLOV, P., SEMERDZHIEV, P., NIKOV, S. & DIMITROV, N., 1952.—[Diseases of sheep and goats (infectious and parasitic diseases, avitaminoses, and forage poisonings).]
 Sofia: Zemizdat, 543 pp. [In Bulgarian?]
- *995—POGGEL, H., 1952.—" Die Komplementbindung zur Feststellung der Trichinose bei Tieren, insbesondere bei Schweinen und über die Anwendbarkeit der Reaktion mit Schweineseren im allgemeinen." Dissertation, Giessen, 61 pp.
- *996—REICHENOW, E., VOGEL, H. & WEYER, F., 1952.—"Leitfaden zur Untersuchung der tierischen Parasiten des Menschen und der Haustiere." Leipzig: J. A. Barth, 3rd edit., 297 pp.
- *997—RILEY, W. A. & WALLACE, F. G., 1952.—" Introduction to the study of animal parasites and parasitism." Minneapolis: Burgess Publishing Co., 6th edit., 94 pp.
- *998—ROHRBECK, E., 1952.—" Strongylidenbefall und Rennleistung bei Traberpferden." Dissertation, Berlin.
- *999—SCHMIDT, J., 1952.—"Untersuchungen über den Kartoffelnematoden." In: Pflanzenschutztagung in Berlin, March 12–14, 1952. Deutsche Akademie der Landwirtschaftswissenschaften, Berlin. Biologische Zentralanstalt, pp. 49–53.
- 1000—SKRYABIN, K. I., 1952.—[Trematodes of animals and man. Principles of trematodology. Volume VI.] Moscow: Izdatelstvo Akademii Nauk SSSR, 759 pp. [In Russian.]

The sixth volume of Skryabin's "Trematodes of Animals and Man" systematically surveys Aspidogastrea, Heterophyoidea and Microphallidae. Each taxonomic unit is defined. There are a number of keys, numerous sets of illustrations in the text and extensive lists of references. The following new data are incorporated. In the Aspidogastrea, the subgenus Multicalyx Faust & Tang, 1936 is raised to generic rank. In the section on Heterophyoidea (which is contributed by Morozov), Cryptogonimidae is enlarged by the addition of Haplorchoidinae n.subf. (for Haplorchoides) and Acetodextrinae n.subf. (for Acetodextra and Pseudexorchis). In 1952 [for abstract see above No. 961bn] Morozov subdivided Siphoderinae into Siphoderea and Paracryptogonimea. A third tribe, Iheringtremea, is now added for Iheringtrema Travassos, 1947. Siphoderea is enlarged by the inclusion of the genera Exorchis and Metadena. These three tribes are distinguished from one another by the position of the vitellaria in the body. In Paracryptogonimea they occupy a limited space in the anterior or medium third, in Siphoderea they lie in the anterior third and in the Iheringtremea they reach from the pharynx to the posterior portion of the body. The section on Microphallidae is by Belopolskaya. Endocotyle incana n.g., n.sp. in Tringa incana brevipes is added to Microphallinae. Spelotrema oedemia n.sp. from Oedemia fusca deglandi differs from S. pygmaeum in the size of the seminal vesicle which is 0.037 × 0.025 mm.; the uterus occupies the posterior quarter of the body and runs parallel to its posterior edge and the eggs are flattened anteriorly. Monocoecum bariurum is transferred to Microphallus as M. bariurus n.comb.; Spelophallus primas and Paraheterophyes pirum are transferred to Spelotrema as Spelotrema prima n.comb. and S. pirum n.comb. New additions to Maritrematinae are Pseudomaritrema postlecithale n.g., n.sp. in Tringa incana brevipes, Numeniotrema musculosa n.g., n.sp. in Numenius phaeopus variegatus, and Diacetabulum curvicolon n.g., n.sp. in T. incana brevipes.

These three new genera have the following characteristics: in Numeniotrema one loop of the uterus is situated between the seminal vesicle and the intestinal branches; in Pseudospelotrema the vitellaria meet in the middle line, form a loop between the testes and send two follicular branches towards the sides of the body; in Diacetabulum there are two ventral suckers. Maritrema sp. of Afanasev, 1941 is now named M. afanassjewi n.sp. M. magnicirrus n.sp. is recorded from Upupa epops but not differentiated from allied species. [Each of the new genera, new species and new combinations in the Microphallidae is attributed to Belopolskaya, 1952 and in Galactosomatidae Sobolephya oshmarini Morozov, 1952 is entered but it is not clear if these are the first descriptions of them.]

R.T.L.

G.I.P.

1001—SKRYABIN, K. I., 1952.—[Trematodes of animals and man. Principles of trematodology. Volume VII.] Moscow: Izdatelstvo Akademii Nauk SSSR, 762 pp. [In Russian.]

Volume 7 of Skryabin's "Trematodes of Animals and Man" deals with Cephaloporidae, Monodhelmidae, Dicrocoeliidae and Gorgoderidae. The section on Dicrocoeliidae is by Skryabin & Evranova but incorporates a number of new genera, species, subspecies and new combinations proposed by others. Oshmarin is responsible for Pancreatrema disacetabulum n.g., n.sp. (type species) from Eurystomus orientalis, Praeorchitrema praeorchis n.g., n.sp. (type species) from Capella gallinago, Unilaterilecithum beloussi n.g., n.sp. (type species) from Pericrocotus roseus, Brachylecithum capilliformis n.sp. from Turdus dauma, B. coturnixi n.sp. from Coturnix coturnix, B. cuculi n.sp. from Cuculus canorus, B. pici n.sp. from Picus canus, B. praetenuis n.sp. from Apus pacificus, B. attenuatum parinum n.subsp. from Parus palustris, B. lobatum strixi n.subsp. from Strix uralensis, Corrigia separatiorchis n.sp. from Actitis hypoleucos, Dicrocoelium moschiferi n.sp. from Moschus moschiferus, Skrjabinus lanciformis n.sp. from Falco subbuteo and Lyperosomum mosquensis cineli n.subsp. from Circlus pallasi: he also makes the following new combinations, Brachylecithum chivosca (Pratt & Cutress, 1949) for Olssoniella chivosca, B. colorosum (Patwardhan, 1935) for Lyperosomum colorosum, B. donicum (Isaichikov, 1919) for L. donicum, B. eophonae (Yamaguti, 1941) for L. eophonae, and B. lobatum glandarii (Semenov, 1927) for L. lobatum glandarii. Evranova describes Brachylecithum kirghisensis n.sp. from Montifringilla alpicola prosvirowi, and B. uigurica n.sp. from Eremophila penicillata albigua. Kasimov supplies descriptions of Dicrocoelium petrovi n.sp. from Alectoris graeca caucasica, Lyperosomum petrovi n.sp. from Francolinus francolinus, and L. schikhobalovi n.sp. from A. graeca caucasica. Lutztrema sp. of Boyd, 1951 is named Lutztrema sturni n.sp. by Skryabin & Evranova. Pigulevski contributes the section on Gorgoderidae and divides Gorgodera into five subgenera, viz., (i) Gorgodera n.subg., type G. (Gorgodera) cynoides Zeder, 1800, (ii) Antodera n.subg., type G. (Antodera) amplicava Looss, 1899, (iii) Extremodera n.subg., type G. (Extremodera) australiensis Johnston, 1912, (iv) Mediodera n.subg., type G. (Mediodera) pagenstecheri Ssinitzin, 1905, (v) Postodera n.subg., type G. (Postodera) varsoviensis Ssinitzin, 1905. He also divides Petalodistomum S. J. Johnston into two subgenera, Petalodistomum n.subg., type P. (Petalodistomum) polycladum S. J. Johnston, 1913, and Staphylorchis n.subg., type P. (Staphylorchis) cymatodes S. J. Johnston, 1913, and Probolitrema Looss, 1902 into Probolitrema n.subg., type P. (Probolitrema) richiardii Lopez, 1888 and Reduxotrema n.subg., type P. (Reduxotrema) clelandi T. H. Johnston, 1934. Pigulevski also contributes Gorgodera (Mediodera) pawlowskyi n.sp. from Rana sp. and Artyukh is responsible for naming as a new variety Dicrocoelium lanceatum sciuri n.var. from a squirrel. The volume is illustrated by 263 line drawings and has comprehensive lists of references [but lacks a host-parasite list]. R.T.L.

SKRYABIN, K. I., SHIKHOBALOVA, N. P., SCHULZ, R. S., POPOVA, T. I., BOEV, S. N. & DELYAMURE, S. L., 1952.—[Descriptive catalogue of parasitic nematodes. Vol. 3. Strongylata.] Moscow: Izdatelstvo Akademii Nauk SSSR, 890 pp. [In Russian.]

In the third volume of Skryabin's "Descriptive Catalogue of Parasitic Nematodes" the Strongylata are surveyed by Skryabin, Shikhobalova, Schulz, Popova, Boev & Delyamure

s joint authors but various individuals are responsible for different sections and items. Formal definitions and differential keys are provided for the taxonomic groups. The species of each genus are arranged alphabetically with their synonymy, hosts, habitat and geographical Histribution. The main text covers 890 pages including 289 pages of illustrations. There are also indexes of the genera and species mentioned in the volume and a bibliography of 1,433 titles. [Various subfamilies, tribes, genera and species are quoted as named by individuals in 1952 but there is no indication whether they are recorded here for the first time or taken from other publications not given in the list of references.] Popova, 1952, is or was responsible for: Chabertiinae n.subf., Zoniolaiminae n.subf. of Cloacinidae, and Sauricolinae n.subf. of Trichonematidae; the division of Trichonematinae into three tribes, Trichonematea, Cylindropharyngea, Gyalocephalea; the division of Murshidiinae into three tribes, Murshidiea, Pharyngostrongylea and Theilerianea; and for a revised definition of Gelanostrongylus Popova, 1952, type Macropostrongylus macrostoma Davey & Wood, 1938. Skryabin & Shikhobalova have created in 1952 a new family Ollulanidae and five new subfamilies Cheiropteronematinae, Cooperiinae, Filarinematinae, Nochtiinae and Oswaldocruziinae: they also add the following new tribes, viz., Filicapitea to Trichostrongylinae; Microstrongylea to Cheiropteronematinae; Hyostrongylea and Trichohelicea to Cooperiinae; Stunkardionematea to Nematodirinae; and Viannaiea, Viannellea and Pseudoheligmostomatea to an emended Viannaiinae Neveu-Lemaire, 1934. Skryabin & Schulz, 1952 are responsible for a new subfamily Haemonchinae and make Longistriatinae a new subfamily in the Heligmosomatidae with two new tribes Longistriatea and Avellariaea. Skryabin & Shikhobalova add Acanthostrongylea, Heligmonellea, Murielea, Pudicea and Squamastrongylea and Schulz a further tribe Impalaiea. Heligmodentostoma n.g. is proposed by Skryabin & Schulz for a species H. microti described by Schulz in 1929 from Microtus sp. [not traced]. Delyamure adds a new subfamily Halocercinae to the subfamilies of Pseudaliidae. Other apparently new records are: in Halocercinae, Delamurella hyperoodoni n.g., n.sp. Gubanov, 1952 from Hyperoodon ampollatus; in Heligmosomatidae, Mammanidula asperocutis n.g., n.sp. Sadovskaya, 1952 from Sorex sp.; here named for the first time Spiculopteragia alcis n.sp. Schulz, Kadenazii, Evranova & Schaldybin, 1952 in Alces alces alces, A. a. bedfordi and Capreolus capreolus. Skrjabinagia Kasimov, 1942 (a subgenus of Ostertagia) is raised to generic rank by Altaev, 1952 who adds S. dagestanica n.sp. [not described] from Ovis aries. Skryabin & Schulz give Oswaldocruzia gozzei as a new name for O. filiformis (Goeze, 1782) of Travassos, 1917, which they do not consider to be identical with Goeze's original species which is listed separately as O. filiformis (Goeze, 1782) sp. inq. Seven species catalogued under Macropostrongylus are transferred later in the volume to Gelanostrongylus. Heligmosomoides travassosi Schulz, 1926 and Nematospira turgida Walton, 1923 are transferred to Heligmosomum as new combinations. G.I.P.

TAYLOR, E. L., 1952.—" Grassland management and parasitism." In: Report of Proceedings. Conference on Metabolic Disorders and Other Problems Related to Grassland and Fodder Crops and Innovations in Animal Husbandry, London, November 20–21, 1952. B.V.A. Publication No. 23, pp. 112–120. [Discussion pp. 120–126.]

Taylor reviews recent work bearing on grassland management in the control of parasitism. He stresses the importance of subclinical infections and summarizes the work on host-parasite relationships of Gordon and Stewart in Australia, and of Michel and Morgan, Parnell & Rayski in Britain. He describes and discusses a striking instance in which the stock on a hill farm was increased two and three times without any special application of artificial fertilizers or seed mixtures, outlines the recent work on the relation of lung parasitism to fog fever and stresses the peculiar difficulties which the complexity of the oecological field presents to the veterinary practitioner.

- *1004—TINER, J. D., 1952.—" A study of the epidemiology of ascarid nematodes in carnivores and rodents." Thesis, University of Illinois, 130 pp.
- *1005—VOGEL, H. & MINNING, W., 1952.—"Wurmkrankheiten." In: "Handbuch der Innerent Medizin", edited by G. von Bergmann, W. Frey & H. Schwiegk. Berlin: Springer-Verlag,; 4th edit., Vol. I, Pt. II, pp. 784–1008.
- *1006—WACHEK, F., 1952.—" System und Biologie der entoparasitischen Tylenchiden." Dissertation, Erlangen.
- *1007—WEINSHEIMER, P. A., 1952.—" Verbreitung parasitärer Tierkrankheiten im Kreise Daun."

 Dissertation, Giessen, 35 pp.
- *1008—WERCKMEISTER, G., 1952.—" Die Inhalation mit Toluol und die intratracheale Injektion in mit Äther-Toluol bei der Lungenwurmerkrankung des Rindes." Dissertation, Hanover, 44 pp. 1
- Too

 WESENBERG-LUND, E., 1952.—" Acanthocephala." In: Fridriksson, "Zoology of Iceland". Vol. 2, Pt. 16, pp. 1-6.

Acanthocephala have only once been recorded in Iceland. From a search of Icelandic material in the Zoological Museum, Copenhagen and in the Royal Veterinary and Agricultural l College, Copenhagen, Wesenberg-Lund records four species: Polymorphus minutus in Somateria mollissima, Corynosoma strumosum in Phoca vitulina and P. hispida, Bolbosoma porrigens in Megaptera boops, and Echinorhynchus gadi in Gadus callarias. Holmes has recorded finding Polymorphus (Profilicollis) ?botulus in a pond on Grimsey. Bolbosoma turbinella from Balaenoptera borealis was found in material in the Berlin Zoological Museum.

*1010-WIEBE, J., 1952.--" Verbreitung parasitärer Tierkrankheiten im Landkreis Springe." Dissertation, Hanover, 31 pp.

INDEXES

				•••				Page
Index of Authors	•••	•••	***		•••		***	
Index of Periodicals	***	***		600			•••	350
Index of Subjects	•••	***				•••	***	358
Corrigenda		***						398

NOTE

In all indexes the reference is to the serial numbers and not to the pages. Numbers in **bold** type indicate abstracts, and numbers in Roman type refer to title-only entries.

In the Author Index there are no cross-references to show joint authorship, but authors of joint papers are listed individually. Thus, a paper by "Brown, B., Jones, A. & Smith, J." would have three separate entries, "Brown, B.", "Jones, A.", and "Smith, J.".

In the Index of Subjects, alphabetization is under the first word (e.g. "Acer sp." before "Acerina sp."). Under the generic name of a helminth the following order is observed: papers on the genus as such; papers on undefined species; papers on new and defined species, e.g.

Capillaria

-- spp.

- aerophila

— amarali n.sp.

In cross-entries under names of hosts, the specific names of new species of helminths are omitted. Anthelmintics are listed under that word, under the name of the parasite or disease, and under the name of the host. Nematicides for plant eelworms are listed separately under that word and under the name of the parasite.

INDEX OF AUTHORS

(The reference is to the serial number: numbers in bold type indicate abstracts.)

15a, 36c, 67b, 108a, 108b, 109a, 138a, 138c, 155g, 1692, 178a, 190a, 283a, 22 IC, 238a, 255a, 312b, 324a, 386a, 324b, 387b, 358c, 468a, 364a, 520a, 543a, 544a, 510a. 657a, 662a, 673a, 702C, 788a, 794a, 702d, 702e, 788a, 794 811e, 824a, 891a, 942a. Abbott, R. T., 487a. Abdallah, A., 430g. Abdel-Malek, E. T., 230di. 281c. Abdel Shafi, M., 430e. Abdussalam, M., 773b, 773c. Abou-El-Naga, I., 430c. Acarel, M., 963d. Achkar, R., 270e, 270f, 294c. Acikalin, H. C., 482a. Ackert, J. E., 230a, 230 230y, 320a. 230x, Adams, E. G. P., 414a. Adelson, L., 752a. Agosin, M., 2b. Aguiar, H. A. de, 800e. Aguiarre, C. V., 587a. Ahlberg, O., 164b. Aidaros, S. M., 430b. Aigle, G., 367b. Akerberg, E., 921a. Akers, R. P., 230v, 590a. Akhtar, S. A., 931a. Akimoto, T., 396a, 396b, 396c, 396d, 686b, 686d. Akman, S., 965a. Akramovski, M. N., 536h. Alberto, V. G., 522b. Albornoz Plata, A., 922a. Albuquerque Neves, H. de. 800g. Aldrich, D. V., 230bs. Alexander, R. A., 388b. Alicata, J. E., 14a, 428m, 785a. Allen, M. W., 75a, 75b, 126g, 132i, 132j, 261f, 966a. Allen, R. W., 132f, 132g, 230bw. Allenby, G. M., 371a, 371c. Allgén, C., 342a, 555b, 555c, 868a. Allison, J. L., 126b, 216a, 483e. Almeida, A. M. de, 783j, 783 l. Almeida, C. G. L. de, 321a, 624a. Altmann, H. W., 4a. Álvares Corrêa, M. O., 809a. Alvarez, M., 352b. Alves, W. D., 282a.

Alves da Cruz, A., 501a. Alwar, V. S., 221b. Amaral, A. D. F., 399d, 680a. Amato Neto, V., 322a, 399a, 399g, 680c. Amberson, J. M., 334b. Ameel, D. J., 101a. Amelung, H., 970. Amrein, Y. U., 230bj, 428b. Anders, W., 750a. Anderson, C. G., 479c. Anderson, P. J., 127a. Anderson, R. C., 530a. Andeweg, J. M., 242a. Andrássy, I., 56a, 592a. Andrews, J. S., 1321, 206a. Angelis, P. de, 757a. Anido, A., 946a. Ansorena de Rizzo, M. A., 577d. Antaki, H., 430i. Antelava, N. V., 106a. Antiga, P. F., 612b. Aparicio Garrido, J., 111a, 871. Appelget, J. G., 2f. Aprosio, N., 574g. Arafa, M. A., 586 l. Arana Iñiguez, R., 612a, 960b. Armand Ugón, C. V., 701a, 959a. Arnold, H., 617c. Arruda, H. V. de, 636b. Artigas, J., 944a. Arvidsson, U. B., 36a. Arvy, L., 327a, 368a. Asakura, S., 396c. Ascoli, W. F., 230d. Atkins, J. G., 512a. Atkins, jr., J. G., 30m, 257e. Atlas, D. H., 579a. Atmarama Rao, G., 404b. Atsumi, T., 396c. Aubert, P., 768e. Aubry, G., 576e. Audy, J. R., 26d. Aulia, M., 826a. Aulong, J., 715b. Aun, R. A., 503a. Avila Pires, C. D. de, 399d, 680a. Awakawa, S., 827c. Ayad, N., 290a. Ayas, E., 765e. Ayite, E., 362c. Azevedo, R. de, 783m. [Azim, M. A.], 430a. B. B., 132c, Babero, 150b,

Bacigalupo, J., 208b. Baer, J. G., 651b, 819b. Bärner, 119a. Baeza Donoso, F., 266a. Baggi, G., 635b. Baila, A. E., 634a. Bailey, W. S., 230ck, 888a. Baines, R. C., 9a, 30a, 30n, 664a. Baird, D. M., 698a. Baker, A. D., 203a, 943a. Baker, D. W., 777b. Balbin, A., 732c. Baldomir, J. M., 587a, 612b. Baldwin, E., 621a. Ball, L., 20 1. Baltazard, M., 79c. Bamberger, J. W., 20x. Bambridge, B., 176a. Bando, T., 411a. Bangham, R. V., 252a. Banks, A. W., 67a. Barbato, E. C. D., 872. Barbosa, F. S., 783a, 783d, 783h, 783i, 783j, 783k. Barnett, E. E., 69c. Barrett, N. R., 640a. Barretto Neto, M., 614a. Barros, J. de R., 801c.
Barry, P. J., 700a.
Barton, W. L., 85a, 382a.
Barua, D., 403a.
Basnuevo, J. G., 185a, 270c, 270c, 270f, 294c, 811a, 811c, 814a. Bastos, A. F., 399h. Batchelder, R. M., 222a. Bates, B. H., 522b. Batte, E. G., 20h, 154a, 644a, 772a. Battista, L. di, 820c. Bauer, F., 148a. Bauer, O. N., 380b. Baughn, C., 230bd, 230be. Baughn, jr., C. O., 419a. Bauman, P. M., 176h. Baurys, W., 716a. Bazán, J., 882a. Bazán de Segura, C., 351a, 479b. Bazzano, H. C., 186b. Beadle, L. C., 18a. Beaver, P. C., 105b, 124a, 428f, 578b. Beck, J. W., 13a. Beckett, A. H., 709a. Begani, R., 748a. Béguez César, A., 946d. Behrens, H., 148c, 528a, 528c.

Bacigalupo, A. d'A., 208b.

230bq.

Babini, D. S., 672b.

Belding, D. L., 160. Bell, D., 580a, 632a. Bell, E. T., 427a. Bell, jr., S. D., 54a. Bell, W. B., 187b. Belleville, G. I., 634c. Beltran, J. C., 186b. Beltrán Monferrer, J. M., 890a. Benassi, G., 635b. Benbrook, E. A., 971. Bendandi, G., 425a. Bennett, H. J., 262a, 262b. Benoldi, E., 741b. Bentami, 574f. Benzinger, F., 570a. Berg, J. A. G. ten, 379c, 465b. Bergenfreid, A., 190f. Bergeon, P., 273a. Berghe, L. van den, 38k. Bergman, B. H. H., 474a. Bergue, C. E. de, 577d. Bernard, J., 123a. Bernstein, J., 908a. Bertram, D. S., 38b, 152c, 983c. Best, P., 143a. Betourné, C., 768e. Beuchat, A., 586d. Beye, H. K., 176a, 318a. Beytout, 367d. Bhaduri, N. V., 91a, 690a, 690c. Bhatnagar, B. L., 180c. Bickerstaff, E. R., 193a. Bieling, R., 210c. Biester, H. E., 556. Biglane, K. E., 262b. Biguet, J., 58b, 200a, 400a, 607b. Bikhovskaya-Pavlovskaya, I. E., 380a, 972. Binet, L., 768e. Bingefors, S., 164h, 973. Biocca, E., 38h, 58j, 64a, 64b, 275a. Bishop, H. W., 234a. Black, S. L., 171a. Blaisdell, K. F., 974. Blamoutier, P., 829d. Blanc, F., 608a. Blanco Rabassa, E., 270e, 270f, 294c. Bławat, F., 653a. Bloemsma, F. F. S. N., 774a. Blynn, E., 20r. Boardman, H. L., 147a. Bocchetti, G., 748a. Boecker, H., 887b. Böhm, L. K., 832a. Böhm, O., 475b, 477a, 907a. Boev, S. N., 1002. Boichard, R., 576e. Boithias, R., 367f, 893a. Boko, F., 153a, 284a. Bokshtein, M. E., 918b. Bona, F., 610a. Bonilla, J., 611b. Bonnema, J. M., 937a. Bonnet, D. D., 471b.

Bonnet, P. M., 725a. Bonnin, H., 260b. Bonsdorff, B. von, 168a. Boragina, R., 672c. Borai, I., 726a. Borchert, A., 458a, 927a, 927b, 927c. Borg, Å., **533b.** Borges Hernández, F., 185a. Borrotchin, M., 801f. Bosch del Marco, L. M., 612b. Bosher, J. E., 141a, 141b, 277a. Bouchet, R., 506a, 517a, 661a. Bourgeon, R., 574g, 574h. Bourn, N., 394a. Bourns, T. K. R., 230k. Bourrel, M., 233g. Bouvier, G., 276a, 516a. Bowesman, C., 61c. Bowman, A., 580a, 632a. Bowman, G. W., 538a. Boyd, E. M., 230bh. Boyer, R., 594a. Boyle, L. W., 767a. Braams, W. G., 774a. Bradley, R. Esp. den. 46 Brande, J. van den, **447a**, **505a**. Brandt, M., **310a**, **573b**. Braude, G. L., **675h**. Bravenboer, L., 242a. Bravo Hollis, M., 588a, 588b. Brede, H. D., 573a. Brenner, M., 683a. Bretschneider, L. H., 774a. Brian, A., 381a. Briceño Rossi, A. L., 816a. Brimblecome, F. S. W., 485a. British Veterinary Association, Brito Gutterres, J. de, 586x. Brock, N., 210a, 617c. Brock, R. D., 3a. Brock, R. D., 3a.
Brooks, A. N., 125a.
Brooks, C. P., 230cr.
Brounst, G., 482b, 817a.
Brown, H. W., 54a, 232a, 333a.
Brown, R., 723a.
Brown, R. G., 177a.
Brumpt, L. C., 58e, 80a, 829a.
Brunson, R. B., 122a.
Brygoo, E. R., 367b.
Buck, I. S., 413a. Buck, J. S., 413a. Buckley, J. J. C., 38a, 152b, 152i. 152i.
Bueding, E., 415a, 638b.
Büsing, K. H., 43a.
Bullock, W. L., 230bx, 428a.
Burch, G. R., 40b, 286a.
Burckhardt, W., 375a.
Burgis, D. S., 776g.
Burgisser, H., 276a, 516a.
Burnod, M., 367e.
Busch E. 168b. Busch, E., 168b. Bustamante, F. M., 586v, 800c. Butorin, F. S., 961bo. Buttner, A., 12a, 58c.

Byer, M. A., 657b. Byrd, E. E., 230dc, 428s, 713e. Byrd, M. A., 854c. Caballero y C., E., 162, 269c, 436b, 502a, 588a, 588b, 588c, 588d, 588e. Cabieses, F., 960a. Cable, R. M., 29b, 230cc, 230cd, 230dd, 230de. Cabral, A., 501a. Cairns, E. J., 30b, 257f. Calado, O. B., 783j, 783 l. Calero M., C., 230o. Calvo Melendro, J., 268a, 576d. Camacho Gamba, J., 611b, 945a. Camain, R., 362c, 367c, 647a. Camand Zogbe, E., 270b. Cameron, T. W. M., 215a. Campbell, C. H., 230bi. Campos, J. V. M., 801d, 801g. Campos, R., 322a, 399a, 399g, 680c. Canabal, E. J., 587a, 612b. Canet, J., 58g. Cantegrit, M., 949a. Cantoni, O., 778a. Capdehourat, E. L., 797b. Carcassonne, M., 727a. Cardeñosa B., R., 306a. Carlier, C., 607b. Carmichael, jr., F. A., 707a. Carmichael, J., 152k. Carney, B. H., 2d. Carney, D. M., 230cx. Carpenter, M. F. P., 975. Carrera, G. M., 124a. Carrell, J., 313a, 313b. Carroll, J. H., 582a. Carta, A., 38i. Carvalho, J. C., 809a. Casarosa, L., 618d. Casile, M., 7a, 445a. Castellanos Fonseca, E., 946d. Catalano, H., 574g, 574h. Catchpole, B. N., 441a. Catellani, G., 618b. Cattini, S., 274a. Caullery, M., 298. Cauthen, G. E., 230cj, 230cv. Cavett, J. W., 13c. Cavier, R., 79a, 79b, 79d, 526a. Cable, J., 492a. Chabaud, A. G., 58f, 58j, 58k, 581, 79c, 327f. Chaia, J., 705b. Chait, D. C., 20q. Chamberlain, R., 164c. Chamberlam, R., 164c.
Chamblee, D. S., 483e.
Chan, K. F., 52a, 52b, 230z.
Chand, D., 220a.
Chandler, A. C., 230bs, 320c.
Chandy, J., 404a.
Chang, K. M., 704a.
Chaplin, J. F., 30i.

Buxton, J. C., 41a.

Chardome, M., 38k. Charms, B., 415a. Chatterjee, K. D., 873. Chatterji, P. N., 688b. Chaudhuri, K. D., 423d. Chauhan, B. S., 784a. Chavarría C., M., 632b. Chechina, A. S., 675g. Cheek, E., 22a. Chelisheva, K. M., 918b. Chen, T. S., 704a. Cheng, Y. K., 575a. Chérigié, E., 710a. Chertkova, A. N., 961n. Chesterman, C. C., 152i, 729a. Chevrot, C., 233i. Chichijo, K., 696g. Chieffi, G., 566a. Chitwood, B. G., 128a, 335a. Chitwood, M. B., 261h. Choquette, L. P. E., 63a, 63b, 372a. Chowdhury, A. B., 690a. Chowdhury, B., 403a. Christiansen, J., 846a, 846b. Christie, D. G., 312a. Christie, J. R., 125a, 257o, 678b, 776b. Christophers, S. R., 34a. Chu, G. W. T. C., 35a. Chubrik, G. K., 213d, 213g, 553b, 675i. Ciamillo, E., 274a. Ciaudo, D., 829c. Ciauri, G., 340c. Ciccarone, A., 354a. Cicchini, T., 340d, 340e. Ciordia, H., 104a, 713b. Clapp, R. L., 20bg. Clare, N. T., 299. Clark, D. T., 132d. Clarke, M. L., 155b. Clarke, O. F., 9a, 30a. Claude, D., 249a. Clay, T., 166. Clément, P., 392a. Cloake, P. C. P., 193a. Cloyd, W. J., 104a. Čmelik, S., 349a, 398a. Cochet, G., 400a. Cockbill, G. F., 509a. Côdo, V., 802a. Coelho, B., 783g. Coelho, E., 740c. Colak, M., 577d. Colbourne, M. J., 531a. Collier, H. B., 371a, 371c. Colonial Office, 300. Combes, 547b. Combes, F. C., 338b. Commun, L., 897a. Coninck, L. de, 464a. Coun, H. C., 54b. Cooper, M., 168b. Cooper, W. E., 125c, 483b. Cooperrider, D. E., 28b, 40c, 95b, 251e.

Corrêa, O., 889a, 889b. Correia Valério, J., 5860. Cort, W. W., 101a. Costa, A. A., 586n. Costa, O. R. da, 948a. Costa Maia, C. da, 586b, 586y. Costantini, H., 233b, 574b, 574c, 574e. Costerizant, G., 208c, 208d. Couch, jr., A. B., 390a. Coudert, J., 426a, 725a. Courtney, W. D., 126c, 236b. Coutelen, F., 58b, 200a, 400a, 607b. tinho, J. O., 322a, 391a, 399a, 399g, 585a, 680b, Coutinho, J **680c, 680f, 680g, 680i,** 80id, 805b. Cowley, H. S., 707a. Cowley, O., 811c, 814a. Cowley Chávez, O., 270e, 270f, 294c. Cox, H. W., 230bf, 230bg, 419b. Craig, J., 412a. Cralley, E. M., 30c, 341a. Crittenden, H. W., 30d, 30e. Croce, J., 399a, 399g, 680c. Crofton, H. D., 29a. Croll, L. J., 5812. Croll, M., 581a. Cronin, M. T. I., 41b. Cross, R. F., 471b. Crosse, J. E., 331d, 790a. Crowdus, D. H., 13e, 281d. Crusz, H., 78e. Cruz Ferreira, F. S. da, 586r, 624c. Cugnasco, C., 742a. Cunha, I. A. da, 615a. Curtillet, 233b. Cushnie, G. H., 356a. Czyżyk, A., 481a. Dalmat, H. T., 176g, 599a,

Dalmat, H. T., 176g, 599a, 599b.

Damasceno, R. G., 800d.
Damluji, S., 235b.
Danilov, V. P., 825a.
Das, E. N., 784b, 784d.
Dastugue, G., 208c, 208d.
Daugherty, J. W., 230bs, 230co, 385a.
Daulton, R. A. C., 383a.
Davis, D. F., 104a.
Davis, L. R., 538a.
Davtyan, E. A., 961bd.
Dawes, B., 120g.
Dawood, M., 430g.
Deane, L. M., 800b, 800d.
Deblock, St., 58b.
De Bont, A. F., 463a.
De Bont Hers, M. J., 463a.
Deboutteville, C. D., 540a.
De Cuttoli, 727a.
DeGiusti, D. L., 230u.

Dejou, L., 728a, 768c. Delahousse, J., 728b. Delak, M., 850a. Delannoy, E., 607a. Delavault, R., 12b, 62a, 337a,, 368b. Delfim, J., 682b. Delfino, F., 666a. Delyamure, S. L., 961v, 961w,, 961z, 1002. De Meillon, B., 833b. Demin, A. A., 953d. Demirleau, J., 829b. Demke, D. D., 146a. Dennis, W. R., 644a. Dent, J. H., 124a. DePaulo, V. J., 230bn. De Sa, A. E., 219a, 406a. Desanlis, J., 976. Deschiens, R., 326a. Descuns, P., 818a, 836a. Dessaint, A., 607b. Desyatov, M. M., 918a. DeWitt, W. B., 230e, 230cy. Deyras, O., 894a. D'Herde, J., 447a. Diachun, S., 198a. Dias, C. B., 801f, 977. Diaz C., E., 190e. Diaz Ungria, C., 630a. Dierckx, F., 464a. Diernhofer, K., 157b. Dieter, C. E., 1742. Diethelm, L., 440a, 884a. Dietrich, W., 528e. Dighiero, J., 587a, 612b. D'Ignazio, C., 811d. Diker, T., 623a. Dikmans, G., 154b. Dimitrov, N., 994. Dimock, A. W., 127b. Dixit, S. G., 92a. Djaenoedin, R., 395a.
Dobbin, jr., J. E., 783a, 783b, 783h, 783i, 783k, 783m.
Dobrovolski, A. V., 553a. Doby, J. M., 58b.
Döll, W., 978.
Dörken, H., 210b, 237a.
Dolfus, R. P., 58d, 339a, 369d, 389a, 418a. Dombrow, M., 709a. Dominguez Navarro, D. L., 732b. Donckaster R., R., 352e. Dor, J., 727a. Dotzauer, G., 573c. Dougherty, E. C., 132k.
Douglas, E., 885a.
Dowell, F. H., 104a.
Draper, W. L., 36a.
Drechsler, C., 460a.
Dropkin, V. H., 230bb. Drozdova, O. S., 438b. Drudge, J. H., 177b, 723a. Duarte, G. G., 626a, 807b. Dubiel, E., 453d.

Dubnitski, A. A., 675c, 917c.
Dubois, G., 651a, 819b.
Ducassou, J., 724a.
Duke, B. O. L., 97i.
Dumazer, R., 233a.
Dumont, L., 661a.
Duran, F., 496a.
Durand, 233c, 233d.
Durbin, C. G., 230cu, 236a, 261g, 772b.
Durich, J., 504a.
Durie, P. H., 348a.
Dutt, S. C., 29f.

Eads, R. B., 20u.
Eagland, R. D., 449a.
Eden, A., 18b.
Edgar, S. A., 176a, 318a.
Edgerly, R. H., 281e, 529b.
Edmonds, S. J., 841a.
Edwards, G. A., 783b.
Efremova, N. I., 536d.
Egan, T. P., 230cw.
Elkan, E., 488b.
Ellenby, C., 463c.
Ells, H. A., 230bl.
Elsdon-Dew, R., 36b, 521a.
El Tamimi, M., 430j.
Endrejat, E., 83b.
Enigk, K., 247a, 294a, 551a.
Erhardová, B., 866a, 964a.
Erhardt, A., 148d, 210a.
Erhart, K. H., 969a.
Erkilic, S., 623a.
Ershov, V. S., 285b.
Esserman, H. B., 346a.
Estrada, H. R., 307a.
European Plant Protection
Organization, 301.
Euzéby, J., 507a, 756a.
Euzet, L., 73a, 540a, 651d,
819b, 852c.
Evans, A. S., 230cs, 424b.

Fahel, J., 583a.
Fahmy, M. A. M., 20be.
Faiguenbaum A., J., 190c, 190h.
Fain, A., 59a, 59b, 328a, 328b,
328g, 819a.
Fairbairn, D., 416a.
Fairley, N. H., 110a.
Fan, P. C., 2e.
Fanta N., E., 190b, 190f, 352c,
352d, 352e, 803a.
Farkas, E., 233f, 576b, 576c.
Farmer, H. H., 433a.
Farr, M. M., 230da.
Fassuliotis, G., 257g, 479d,
760b.
Fatist, E. C., 2b, 586a, 586w.
Fawzi, M., 235b.
Fedeli Avanzi, C., 597b.
Feder, W. A., 257h, 365a.
Fedtke, H., 735d.

Exner, H., 181a.

Fedyushin, A. V., 961bc. Fedyushin, V. P., 536k. Feld, 363a. Feldmesser, J., 125b, 257i, 479d, 760b. Fellus, B., 576a. Fenwick, D. W., 97a, 164a, 331b. 331b.
Ferguson, M. S., 230bm.
Fernando, E. F. W., 78d.
Fernando, W., 78b, 78c.
Ferrand, J., 233i.
Ferrari, G., 635c, 635e, 635d.
Ferrari, L., 666a.
Ferreira, J. M., 503b.
Ferreira Guedes, L. 586f. Ferreira Guedes, J., 586f. Fetchko, A. M., 582a. Fethers, G., 473a. Ficai, A., 951a. Fidalgo Díaz, J., 451c. Fidalgo Díaz, M. A., 451b. Field, W., 230u. Figueirêdo, A. de, 399i. Filizzola Filho, B., 799b. Filmer, J. F., 493a. Fischer, H., 838a. Fischthal, J. A., 843a. Fischthal, J. H., 101e. Flores Barroeta, L., 269c, 815b. Florez Tascon, F. J., 730a. Florez Tascon, F. J., 730a. Floros, A., 830b. Folse, D. S., 230ck. Ford, C. M., 41a. Forsyth, R. A., 374b. Foster, A. O., 230p, 230q. Fraga de Azevedo, J., 697a. França, O. H. da, 399e. França, O. H. da, 762a. Frank, A., 861a. Frank, J. F., 10d, 11a. Franklin, M. T., 60a, 164f, 164 l. Fraser, L. R., 347a. Fratta, I., 230m, 428n. Freedman, L., 521a.
Freeman, R. S., 20y, 87b.
Freer, P., 38b, 38c.
Freitas, J. A. de, 799e.
Freitas, J. F. Teixeira de, 495a.
French, R. G., 30c.
Freyer, K. G., 30c. Freytag, K., 979. Freytag, R. E., 176f, 428q. Frontini, C. A., 577a. Fros, J., 674a, 674b. Fueki, K., 437a. Fukano, H., 601b. Fukatsu, R., 601a. Fukushima, T., 396b. Furchner, H. O., 980.

Gaafar, S. M., 200, 230y. Gaase, A., 170a, 549a. Gadea, E., 782a, 782b, 939a. Gadgil, R. K., 689a. Gagarin, V. G., 961bp. Gaiko, B. A., 918b. Galal, O., 430f.
Gállego Berenguer, J., 33c, 135a. Galliard, H., 58a, 327f, 622a. Gama Imaginário, J. da, 682a, 818b. Gangstad, E. O., 126f. Garcia-Barón, A., 804b. García Capurro, R., 612d. García Fontes, W., 587b. García Gil, M., 730a. Garcia-Laverde, A., 2b. García Morán, J., 947a. Garnett, K. J., 172a. Gaud, J., 444a, 645a, 739a. Gautier, P., 607a. Gayet, R., 715a. Gayot, G., 63a, 63b. Gediz, C., 963d. Gefter, V. A., 961bq. Gefter, V. A., 961bq. Geisler, G., 617c. Gelfand, M., 69c, 282a. Gelin, O., 523a, 856a. Gelormini, N., 900a. Gerber, J. H., 105a. Gerlach, S. A., 610d. Gervois, M., 400a. Gharpure, P. V., 403b. Ghose, D. C., 690b. Ghosh, K. K., 423b. Giaretta, D., 685a. Giaretta, D., 685a. Gibson, C. L., 2c, 176g, 230d, Gibson, C. L., 2c, 176g, 599b.
Gibson, T. E., 19c.
Gierthmühlen, 718a.
Gilbert, J. C., 770a.
Gilbert, J. L., 842a.
Gilchrist, H. B., 225a.
Gill, A. J., 578c.
Gill, D. L., 31b.
Gillot, F., 576a.
Ginesinskaya, T. A., 675f. 675b. 675f. Giorgi, L., 820b. Giotsas, Z., 830b. Giovannoni, M., 807a. Girgis, B., 38j, 175a. Glocklin, V. C., 416a. Gloss, R., 889a, 889b. Godbille, M., 593a. Goddard, R. F., 232a. Gönnert, R., 4a. Göeters, W., 293a, 552a. Goffart, H., 117b, 336a, 393a, 847a, 870a. Goinard, P., 818a, 836a. Goldberg, A., 20e, 319b. Goldman, M., 20p. Goldsby, A. I., 189a. Gomar Guarner, F., 734a. Gomes da Silva, J., 636a. Gon, S., 90a. Góngora, J., 611b. González, O., 812a. Gonzalez Alvarez, J., 732a.

González Castro, J., 269a. González Mendoza, R., 191a. Gooch, J. M., 471b. Goodall, W. S., 5a. Goodey, J. B., 60b, 97d, 97e, 164k, 331c. Goodey, T., 26b, 164m, 164n. Goodman, J. D., 981. Gordin, R., 168a. Gordon, H. McL., 165f, 546c, 755a. Gordon, R. F., 259a. Gordon, R. M., 38d, 152e, 983b. Gorham, J. R., 678c. Goto, K., 601a. Goto, S., 919b. Gouvéia, E., 586i, 762a. Govinda Rao, A. R., 180a. Grace, H. K., 430b. Graf, E., 887a. Graham, T. W., 30f, 125d, 126e, 483c. Grainger, J., 60c, 194a. Grangaud, P., 233f. Granville, A., 593a. Grau Miró, M., 811b. Greany, W. H., 334c, 334f. Green, H. E., 230cn. Greenberg, B. G., 225a. Greene, H. J., 467a. Greenleaf, W. H., 483a. Greenslit, F. S., 54b. Grégoire, C., 593a. Gregoire, E., 20m. Greig, A. M. W., 493b. Greiner, H., 244a. Gremliza, L., 294d. Grenier, P., 74a. Gresson, R. A. R., 408a. Griffiths, H. J., 678c. Griffiths, R. B., 38d. Grimaldi, J. P., 631a. Grittner, I., 551a. Grocott, R. G., 436b, 588b, 588c, 588e. Groth, A. H., 93a. Groves, H. F., 835a. Gruber, G. B., 863b. Grüninger, U., 735a. Güçhan, S., 694c. Güemes R., M., 190g. Güralp, N., 288a. Gürevin, I., 694a. Guerin, P., 768a. Guevara, R., 307a. Guibal, J., 738a. Guilford, H. G., 843b. Guilhon, J., 492c. Gullion, G. W., 76a. Gunhold, P., 192a. Gunstone, F. D., 223c, 223d. Guntz, M., 574g. Gupta, P. N., 691a. Gushanskaya, L. K., 961q, 961r, 961s. Gustin, 574h. Gutiérrez Arrese, D., 268b.

Haapanen, L., 167a. Hack, M. C., 464a. Hacgermark, U., 886a. Haemmerli, U., 375a. Hagemeyer, J. W., 132j. Hagenunger, W., 735a. Hager, R., 25a.
Halawani, A., 103b, 430g.
Hale, W. M., 775a.
Haley, A. J., 230bx.
Hall, S., 224a.
Hammick, D. L., 223b. Hand, C., 20g. Hansen, M. F., 230x. Hanson, C. H., 144a, 483e. Harada, F., 291a. Harding, H. B., 940a. Hargis, jr., W. J., 55b, 132h, 541a, 713g. Harris, D. G., 234b. Harris, F. C., 289d. Harrison, A. L., 760d. Hartwich, G., 967a. Haseeb, M. A., 69d. Hashimoto, T., 396a. Hastings, R. J., 141a, 141b, 277a. Hasuko, E., 919b. Hausmann, W., 458b. Havis, L., 128a. Hawes, S. C., 641a. Hawking, F., 69a, 152g, 152h, 357a, 358b.

Hawley, W. O., 30q.

Hayashi, S., 409a.

Heeley, W., 205b.

Heggie, R. M., 223c, 223d.

Heidigger, E., 163. Heilig, R., 423a. Heine, W. I., 914a. Heinz, H. J., 143b. Heinze, K., 158a. Held, H., 862a. Helfand, M., 378a. Helg, L., 670a. Hellmeister, O., 343a, 616a. Helo, J. E., 497a, 497b. Hemming, F., 8a, 140a. Hendershott, R. A., 777a. Hendricks, J. R., 176d, 225b, 230bc. Hendrikse, D., 9042, 904b. Hennings, 118d. Henriques, F. F., 586q. Henri-Sarles, 727b. Hepner, W., 116a. Hernández de la Barca, A., 946a. Herrick, C. A., 843b. Hesling, J. J., 97b. Heuck, F., 440a, 884a. Hey, A., 117d. Heydt, R., 23a. Hightower, B. G., 20u. Hightower, J. A., 176h. Higley, J. C., 42a. Hill, W. C. O., 488a, 488e.

Hinchliffe, M. C., 101f. Hingorani, M. K., 218a. Hirschmann, H., 867a. Hitchcock, C. R., 427a. Hitchcock, D. J., 251c. Hicheck, Dr. J., 244. Hloucal, L., 934a. Hobson, A. D., 18a, 18b. Hoekenga, M. T., 176e. Hoeppli, R., 659a, 769a. Hoeven, J. A. van der, 379a. Hoffman, G. L., 230bz. Hoffmann, A., 57a, 325a. Holdeman, Q. L., 125d, 483c. Hollingsworth, K. P., 87d. Holzman, D., 466a. Homeyer, B., 722a. Honain, A., 705b. Honma, M., 719a. Hood, M., 578a. Hopkins, C. A., 13g, 179a, 334e. Horáček, J., 898a. Horáčková, M., 932a. Hornbostel, H., 210b, 237a, 573c. Horrax, G., 250a. Hosaka, Y., 702h. Hoshina, T., 695a, 714a. Houck, K. H., 378a. Houel, J., 233a, 233e. Howard, P., 20bf. Howell, H. B., 126c. Hoy, J. M., 744a. Hrubcová, M., 932a. Hsieh, K. N., 316a. Hsieh, K. N., 310a.

Hsü, H. F., 2e.

Huang, T. C., 2e.

Hubl, H., 159b.

Hueck, O., 552b, 861b.

Hueston, J. T., 619a, 620a.

Hughes, B., 193a.

Hughes, M. H., 289b.

Huisman, C. A., 463d. Huijsman, C. A., 463d. Hulsman, C. A., 465a. Hulshoff, A. A., 465a. Humes, A. G., 230v, 590a. Hunt, J. S., 230ce, 230cf, 982. Hunter, G. C., 422a. Hunter, III, G. W., 13d, 381, 176f, 230s, 428j, 428q, 457a, 696f, 696h. Hunter, W. S., 20q, 101d, 230c. Hupka, E., 528c. Hussey, K. L., 230b. Huston, E. J., 230bh. Hutton, R. F., 228a, 646a. Ichinohe, M., 254a, 265a. Igarashi, T., 719a. Ikoma, H., 649c. Imperati, L., 137a. Imanaga, K., 706a. Innes, J. R. M., 41d, 70a, 221a, 251a, 358a, 569a. Inoue, I., 603b, 864a. International Nematology Symposium, 164.

International Veterinary Congress, 165.
Irtel, H., 623a.
Isaiah, P., 404a.
Ishii, N., 291b.
Ishii, S., 409a.
Ishii, Y., 402a.
Ishiitani, R., 905a.
Ishiizaki, T., 411a.
Islas, C., 272a.
Ito, F., 396e.
Ito, J., 410b, 410e, 696a, 696j.
Ito, S., 649c.
Ivanov, A. V., 296a.
Ivens, V., 230cm, 319a, 839a.
Iyengar, M. O. T., 654a.
Izumi, S., 410a, 410c, 410d.

Jachowski, jr., L. A., 176b.
Jacks, A., 776g.
Jacks, H., 253a, 469a.
Jackson, P. K., 230bw.
Jacobs, T. P., 467b.
Jain, S. L., 912b, 912c.
James, T., 217a.
Janssens, P. G., 328c.
Jaques, W. E., 184a, 184b.
Jarcho, S., 88a.
Jarpa, A., 812a.
Jaskoski, B. J., 87c, 230cw, Jachowski, jr., L. A., 176b. 937a. Jawad, H., 420a. Jelliffe, D. B., 289a, 544d. Jellison, W. L., 122a. Jensen, H. J., 479c. Jepsen, A., 165a. Jettmar, H. M., 181a, 548a, 604a. Jevtić, M., 668a. Jidejian, Y., 229c. Jinnai, D., 425b. Jirovec, O., 43b, 201a, 932a. Johnson, A. W., 205c. ohnson, E. M., 198a. ohnson, R. E., 466a. Johnson, K. E., 400a. Johnson, S., 20p. Johnson, V. M., 2b. Johnston, Jr., J. H., 602a. Johnston, T. H., 841a, 841b. Jonathan, S. R., 151a. Jonathan, S. R., 151a.
Jonchère, H., 362a.
Jones, A. W., 104a, 713a.
Jones, D. J., 206a.
Jones, F. G. W., 164g.
Jones, M. F., 20t.
Jones, M. F., 20t.
Jones, N. L. C., 546d.
Jones, N. S., 546d.
Jones, T. R., 546a.
Jordan, P., 38m.
Jordano, D., 33b.
Joyeux, C., 339b, 651b. oyeux, C., 339b, 651b. ung, R. C., 94a, 289a. Jungalwalla, A., 911a.

Kämpfe, L., 984. Kagan, I. G., 37a, 230cz, 320e, 424a, 424c. Kakizaki, C., 686a. Kalinin, I. V., **536f.** Kalmi, I., 929a. Kamalov, N. G., 953b. Kamenear, H., 579a. Kangina, K. I., 961bs. Kano, R., 409a. Karpišek, J., 928a. Kartman, L., 230dk. Kasai, G., 686d. Kasimov, G. B., 961t, 961u. Kates, K. C., 230p, 230q. Kaufman, jr., E. H., 13d. Keech, M., 485a. Keleher, J. J., 421a. Keller, H., 211a, 985. Keller, J. R., 127b. Kennedy, C. B., 132g. Kennedy, P. C., 81a. Kerr, K. B., 13c, 129b, 230cn, 936a. Kerrest, J., 728c, 941a. Kershaw, W. E., 152d, 152e, 334d, 334g, 639a. Kessel, J. F., 176a. Ketz, H. A., 883a. Killingsworth, W. P., **147a.** Kim, C. W., **230bc.** Kincaid, R. R., **202a, 776i.** Kinnear, A. A., 143a. Kips, R. H., 447a. Kiryanova, E. S., 296b. Kitamura, K., 338a. Kivits, M., 328d. Klimars, G., 292a. Klimars, G., 292a.
Kloos, K., 440a, 884a.
Kniskern, V. B., 281a, 529a.
Kobayashi, Y., 411a.
Kobulei, T., 726b.
Koczorowski, S., 933b.
Köhler, H., 119b.
König, E., 863b.
Konigstein, R. P. 552c. Koenigstein, R. P., 552c. Koesling, G., 986. Kolligs, A., 987. Kolstad, P., 958a. Komine, I., 409a. Korniya, Y., 696a, 696d, 696e, 696g, 696i, 696k. Kondo, T., 649b, 913a. Kono, I., 642a. Konorska, R., 933b. Korovaev, N. M., 849c. Kosheva, A. F., 553d. Kostić, D., 668a. Kothare, S. N., 219a. Kotlán, A., 165d, 309a. Kotthoff, P., 684a. Kouri, J., 270a. Kouri, P., 136a, 270a, 270d, 946b. Kourias, B., 660a, 768b, 768d. Koutz, F. R., 32a, 835a.

Kozar, Z., 653a.
Kozić, L., 850c, 880c.
Krahnert, R., 118b.
Kraljevic, R., 499a.
Kraneveld, F. C., 395a.
Krastin, N. I., 213b, 536c.
Krawczyk, H. J., 230bt.
Kreis, H. A., 516b, 851a.
Krieger-Königsbert, 735b.
Krishnaswamy, A. K., 405b.
Krispien, H., 148b.
Kroft, W. G. van der, 242a, 910b.
Krotov, A. I., 961s, 961x, 961y, 961z.
Krstić, B. N., 924a.
Kruedener, R. von, 988.
Kruidenier, F. J., 230cg.
Krull, W. H., 81c, 81d, 209a, 230ch, 374a, 374c.
Ksenofontova, M. I., 938a.
Kubicz, S., 933b.
Kubo, S., 396a.
Kudenko, I. D., 145b.
Kufs, H., 853a.
Kudenko, I. D., 145b.
Kufs, H., 853a.
Kuhns, E., 122a.
Kulik, T. N., 675b.
Kumagai, R., 702g.
Kume, S., 702b, 702i.
Kumova, C., 963d.
Kuntz, R. E., 20b, 37b, 132b, 229b, 261i.
Kurowski, A. C., 639a.
Kusminsky, N., 801g.
Kuźmicki, R., 933c.
Kyles, P. M., 132f.

Laan, P. A. van der, 120c.
Lackany, A., 103a.
Lacour, G., 672c.
Lafferty, J. W., 124a.
Lagrange, E., 59a, 59b, 327c, 328b, 328i.
Laha, N. K., 405a.
Lal, M. B., 120a.
Lamy, H., 852b.
Lamy, H., 852b.
Lamy, L., 852b.
Landtman, B., 470a.
Langer, L., 672b.
Lanternier, J., 426b.
Laquière, 233b.
Larbe, F., 426b.
Larghero, P., 959b.
Larsh, jr., J. E., 225a, 230t, 230bc, 230bc, 230bi.
Latham, W. J., 833a.
Latteri, S., 952a.
Lauret, G., 606a.
Laurie, W., 789a, 791a.
Lawrence, D., 38e.

Kovaleva, T. S., 285a.

Leader, G. H., 41b. Leão de Moura, S. A., 809b. Lear, B., 125b, 257n, 257t, 373a, 667a. Lee, C. L., 230cz, 424c. Lee, Y. C., 575a. Leedy, W., 20bf. Lefevere, S., 464a. Lefrançois, R., 200a. Le Génissel, 233e. Leigh, W. H., 230df. Leikina, E. S., 918b. Leiper, J. W. G., 155f. Lenschow, 118d. Leon, G. V. de, 307a. León León, J. M., 946c. Le Riche, H., 143a. leRoux, P. L., 38h. Levashov, M. M., 961bm. Levine, N. D., 230cm, 319a, 839a. Levrat, M., 725a. Le Zotte, jr., L. A., 230cb. Li, S. Y., 2e, 20s. Liaras, 574d. Lienert, E., 157c, 157d. Lightwood, R., 485a. Lima, L. I. de A., 783c. Lin, S., 457a. Lindhardt, K., 527a, 681a. Lindquist, W. D., 20k. Linford, M. B., 387a. Ling, L., 677a. Lino de Sousa, 501a. Lipparoni, E., 514e. Lippi, M., 340a. Lobato Paraense, W., 798b. Lobo, M. B., 399f. Locker, B., 21a. Logachev, E. D., 213a, 213e, 675a, 675d. Logé, G., 492c. Loitsyanskaya, M. S., 455a. Loizzi, A., 951b. Lopes de Faria, J., 874. López-Neyra, C. R., 33a, 269b, 498a, 627a. Lordello, L. G. E., 636a, 796a. Losee, R. E., 122a. Lotmar, F., 515a. Loubeyre, J., 233f, 576b, 576c. Loughnane, J. B., 700a. Loures, J. de C., 399h. Lowman, C. B., 888a. Lownsbery, B. F., 30g, 257c, 257j, 478c. Lownsbery, jr., B. F., 30q. Lownsbery, J. W., 30g, 478c. Lucker, J. M., 369b. Lucker, J. T., 20i. Lührs, 902a, 920a, 920b. Lumbroso, P., 363a. Lundberg, U., 955a. Lurie, H. I., 833b. Luttermoser, G. W., 230cp. Luzina, A. G., 906a. Lynd, F. T., 95a.

McBeth, C. W., 257p, 935a. McCraw, B. M., 204b, 372b. McFadden, I. M., 147a. McFadzean, A. J. S., 531b. McFadzean, J., 38f, 69b. Macfarlane, W. V., 110b, 428c. McGaughey, C. A., 41c. McGregor, I. A., 152h, 152 l, 358ь. McGuire, D. C., 770a. Machado Esparza, L. F., 946a. Machado Ferraz, D., 800a, 800f. Machado de Mendonça, J., 495a. Machulski, S. N., 961be. Maciel, H., 355a. McIntosh, A., 230da, 369a. McKay, R., 700a. Mac Keith, R., 605a. Mackie, A., 195a, 195b, 223a, 638a. MacLagan, D. S., 89a. McLeod, J. A., 305. McMahon, J., 120f. McMullen, D. B., 176c. McPherson, S. E., 461a. McQuay, jr., R. M., 13f. Macy, R. W., 428i, 746a. Madden, F. J., 546b. Madsen, H., 669a. Magalhães Neto, B., 783b, 783e, 783 1. Magath, T. B., 2g. Magaudda-Borzi, L., 513a. Magid, A., 38j. Mai, W. F., 30o, 125b, 257c, 257d, 257k, 257n, 257t, 518a, 667a, 760c. Makar, N., 879a. Makhmudova, B. A., 953a. Makkas, M., 652a. Malheiro, D. de M., 807a. Malheiros Santos, J., 798b. Maligin, S. A., 536g. Malkani, P. G., 155a. Mallarmé, J., 826b. Mamo, A., 705a. Mañas Montalvo, J., 269a. Mandarino, E., 637a. Manier, J. F., 361a. Mann, K. H., 488d. Mann, P. H., 230 230m, 251b, 251d, 428n. Mannami, C., 930a. Manni, C., 591a. Manson-Bahr, P., 84a, 152a, 235a, 379b, 531d, 557.

Manter, H. W., 369c.

Mapes, C. R., 20n, 81c, 81d, 209a, 230ch, 374a, 374c.

Marangos, G., 660a. Marchionatto, J. B., 494a. Marini, E., 747a. Markell, E. K., 703a. Marković, D., 570a. Markowski, S., 72a, 422b. Markwalder, J., 828a.

Marogna, L., 780a. Marques, R. J., 58i. Marsh, H., 10e. Martin, H. M., 261c. Martin, W. E., 20x, 230i. Martin Vaquero, B., 534a. Martinez Báez, M., 450d, 731a. Martinez Carrera, N., 946a. Martinez de Jesús, J., 17a, 896a. Martinez Pedroza, M., 815a. Martini, D., 665a. Martini, J., 812a. Maruyama, M., 396b. Maskar, U., 694b. Massi, O., 275a. Massonnat, J., 576e. Mastrandrea, G., 340b. Masusaki, M., 831a. Mathieu, J., 328f. Mathois, H., 157c, 157d. Matilla, V., 589a. Matsuno, K., 686c. Matsuyama, T., 719b. Mauzé, J., 367a. Mawson, P. M., 841b. May, J. M., 16a. Mayan, L., 608a. Mayaudon T., H., 813b. Mayer, T. C., 713a. Mayhew, R. L., 230ci, 428r. Mazzotti, L., 243d, 243e, 766a. Medoc, J., 706b. Meeser, M. J. N., 434a. Mehlman, B., 87a, 230cq. Meier, P., 375a. Meira, J. A., 503b. Meira e Cruz, J. H., 625a. Meisel, H., 735c. Meleney, H. E., 2d, 13b, 20j, 230cx. Melki, L., 482b, 705b. Mello Balthazar, A. de, 616c. Melnotte, P., 949a. Melvin, D. M., 230h. Méndez Pinilla, A., 270b. Mendheim, H., 294b. Mendoza, D., 959c. Menezes, H., 801a. Menon, T. C., 92c. Mérab, A., 482b, 705b. Mercer, H. H., 322b. Mergold, D. P., 439a. Mesquita, B. de, 586k, 586m. Mestdagh, M., 328f. Meyer, P. R., 147a. Mezzetti, A., 692a. Michel, J. F., 120b, 120d, 120e, 155c. Michon, P., 949a. Mihajlović, S., 850e. Mikačić, D., 850a. Mikami, S., 686d. Mille, R., 176a, 318a. Milleman, R. E., 230bk. Miller, L. I., 257 1.

Miller, R. B., 643a. Miller, W. S., 6a.

Mimioğlu, M., 963b. Mini, M., 741c. Ministry of Agriculture Fisheries; 44, 45, 46. Minning, W., 1005. Minou, A., 79c. Miretski, O. Y., 213c. Mirsa, A., 816b. Mirsa, M., 816b. Misra, B. G., 248a, 405b. Mitrovich, M., 137b. Mitsuno, T., 706a. Miyasaka, S., 636a. Miyazaki, I., 402a, 402b, 402c. Modesti, G., 635a. Moignoux, J. B., 50a, 208a, Moiroud, P., 727b, 727c. Moise, R., 330a. Mol, L. van, 447a. Molese, A., 566a. Moller Meirelles, M., 399c. Momose, T., 831a. Mondon, E., 735a. Mongelli Sciannameo, N., 598b, 635a, 820a. Monnerot-Dumaine, M., 260c. Montanari, A., 820d. Monte, T. de, 139b, 808a. Moore, D. V., 2d, 13b, 20j, 230cx. Moore, J. P., 472a, 749a. Moraes, J. G. de, 783j, 783 l. Mora Torres, J., 732a. Moreira, M., 399f. Moreira da Fonseca, J., 260a. Morel, P., 725a. Moretti, G. F., 260b. Morgan, D. O., 142a.
Morhouse, C. H., 358e.
Morini, E. G., 490a.
Morisani, S. R., 761a.
Morley, J. S., 417a, 417b.
Morozov, F. N., 961bn. Mosera, R., 587b.
Most, H., 2d.
Most, H., 2d.
Moychan, N. A., 455a.
Moyle, V., 621a.
Mozgovoi, A. A., 213f, 849b,
961j, 961k.
Mozley A 875 Mozley, A., 875. Mühlbauer, H., 459a. Müller, G., 554a. Münch, K., 989. Muggleton, W. J., 152a. Mukasa, K., 265a. Mules, M. W., 67d. Mulhearn, C. R., 787a. Mullet, S., 400a. Mullin, R. S., 30h. Mulzer, B., 990. Muñiz González, J., 804a. Muñoz G., A., 190e. Muñoz Rivas, G., 497c. Munro, D. C., 223b. Munro, I. B., 41a.

Muras, O., 959c.

Murase, K., 696d. Murgatroyd, F., 38e. Murphy, M. F., 320d. Murray, N. A., 2b. Murray, R. W., 488b.

Naerland, G., 165e. Naffah, N., 817a. Nagano, K., 38 1, 457a. Najarian, H. H., 20bb, 230dg, 230dh, 230di. Najim, A. T., 991. Nakamura, S., 410a. Naletov, N. A., 961bh. Nani, A., 500a. Nani, A., 500a.
Náquira, F., 352a.
Narumi, H., 396d.
Naude, T. J., 388c.
Naumann, P., 671b.
Navarranne, P., 362c.
Neethling, J. L., 138b.
Neethling, L. J., 511a.
Neethling, A. 352a, 352 Neghme, A., 352a, 352b. Neghme R., A., 190d, 944a. Neiland, K. A., 230g, 428d, Nettel F., R., 243b, 450c. Netzahualcoyotl Diazconti, A., 892a. Neuhaus, W., 158b, 159a. Nevenić, N., 850b, 850f, 880a, 880b. Neveu, P., 594a. Neveu-Lemaire, M., 876. Newson, I. E., 302.
Newson, W., 141a, 141b.
Newton, W. L., 20t, 230j.
Neylans, jr., O. S., 713d.
Nicholas, W. L., 152e, 334e.
Nickel, E. A., 528d.
Nicolai 574f Nicolai, 574f. Nieć, L., **245e.** Nigon, V., **71a, 79e, 337a, 359a.** Nikolskaya, N. P., **380b.** Nikov, S., 994. Niño, F. L., 765a, 765b. Nishimiya, H., 696g, 696i. Nishimura, T., 753a. Nishizawa, T., 721a. Nix, H., 442a. Noda, R., 895a. Noger, C., 327b. Nolte, H. W., 117c, 119b, 376a. Norcross, J. W., 250a. Nor El Din, G., 430j. Noronha, H. de, 801e. North, A. D., 96b. North, A. D., 96b.
Norton, D. C., 663a.
Nosik, A. F., 865a.
Nouvel, J., 199a.
Novi, S., 24a.
Nozawa, Y., 827a, 827b.
Nunes, P., 586h.
Nusbaum, C. J., 30i.
Nyberg, W., 567a.
Nyholm, K. G., 869a.

Obez, A., 200a. Ockerse, T., 143a. Odlaug, T. O., 529c. Ogawa, Y., 291b. Ogino, C., 714a. Ogren, R. E., 428p. Oike, K., 831a. Oka, Y., 702g. Okabe, K., 696c. Olds, R. J., 67c. Oliger, I. M., 961bt. Oliveira, J. E. de, 399f. Oliveira Ferreira, M. de, 800a, 800f. Oliveira Lecuona, M., 586r, 624c. 024c.
Oliver, W. T., 77a.
Oliver-González, J., 98b, 350b.
Olivier, L., 1a, 20a, 176d, 230cq.
Olsen, L. S., 491a, 564a.
Olsen, O. W., 101c, 150d.
Olson, L. J., 230x.
Onabamiro, S. D., 61a, 488c, 544b. Ono, Y., 702a. Oostenbrink, M., 117a, 149a, 164e. Orekhov, M. D., 536b. Ormerod, W. E., 357a. Orris, L., 338b. Ortiz, I., 816b. Osche, G., 297a, 550a. Oshio, Y., 649a. Osimani, J. J., 612c. O'Sullivan, P. J., 65a. Oteifa, B. A., 30j, 257a, 261e, 335a. Otori, Y., 428j. Otto, G. F., 54a, 176b, 187a. Ottolina, C., 799a. Ovnatanyan, K. T., 953c, 956a. Oytun, H. S., 963c. Ozaki, H., 432b. Ozaki, Y., 432a, 432b, 603a.

Palma, E. C., 186a, 186b, 611a. Palmer, B. M. F., 936b. Palombi, A., 610b, 881a. Pan, C., 13d, 38 l, 176f, 230s, 696£. Panebianco, F., 618c. Panisheva, L. V., 917a. Papadopoulos, J., 763a. Pape, H., 117e. Papi, R., 806a. Paramonov, A. A., 961bi, 992. Pardi, M. C., 626a, 807b. Parent, M., 328e. Paris, P., 362d. Parmalee, P. W., 840a. Parmell, I. W., 142a, 155d, 638a. Parra, B. E., 713c. Parreira, F., 586i, 762a. Parrisius, G., 993. Parsons, H. E., 183a. Pasqualino, A., 340f.

Pate, J. B., 126f. Patel, J. C., 219b. Patrón Hernández, M., 810a. Patten, J. A., 20bd.
Paul, K., 861a.
Pavišić, Z., 923a.
Pavlov, P., 994.
Pavlovski, E. N., 553c. Payet, M., 367c. Payne, W. J. A., 314a. Pedemonte, P. V., 612d. Peel, E., 328f. Peers, J. H., 230bt. Pegreffi, G., 207a. Pellegrini, D., 514c, 598a, 618e, 781a. Pellissier, A., 200b, 200c. Pelot, G., 715b. Peluffo, É., 186b. Peña, R. T. de la, 450d. Pereira, M. B., 616a. Pereira, O. A., 614a, 799d. Pérez Hurtado, F., 946e. Perlowagora-Szumlewicz, 800e. Perret, J. L., 651c. Perry, V. G., 125a, 479e, 776c. Perumal Pillai, C., 41d, 70a. Peschiera, C. A., 882a. Pesigan, T. P., 429a. Pessôa, S. B., 680d, 680e, 680f, 680g, 680h, 805b. Pester, F. R. N., 97j. Peters, B. G., 97f, 164d, 205a, 331a, 792a, 793a. Peterson, L. C., 518a. Petitdidier, 492b. Petrasch, 170a. Petrochenko, V. I., 296c. Petrov, A. M., 917a. Petrović, Z., 880c, 850b. Petrovitch, Z., 909a. Petru, M., 899a. Petrushkin, A. A., 938a. Pfister, R., 7b, 362a, 362b, 877. Phipps, F., 366a. Piaggio Blanco, R. A., 587a, 587b. Piana, G., 618a, 823a. Pichon, G., 173a. Pierotti, P., 597a. Piétri, 233c, 233d. Pietri, H., 574g, 574h. Piette, M., 363a. Pihl, H., 940a. Pilleri, G., 139b, 808a. Pimps, E., 736a. Pinotti, M., 34b. Pinto Coelho, D., 586c. Piquer, G., 360a. Pitcher, R. S., 331d, 790a. Pitchford, R. J., 522a. Pittman, H. A. J., 96a. Plane, 574i. Podyapolskaya, V. P., 961b. Poggel, H., 995 Poirier, M., 326a.

Pokorný, M., 932a. Polozhentsev, P. A., 961bk. Polyanski, Y. I., 962a. Polzin, H., 458a. Pommier, G., 878. Poncelet, P., 768a.
Pontes, J. F., 801d, 801g.
Poole, J. B., 204a.
Popova, T. I., 1002.
Popova-Batueva, L. V., 849d. Porges, M., 570a. Porter, A., 488a, 488f. Porter, D. A., 538a. Pototschnig, H., 279a. Potts, D. E., 176f. Pou, M. C., 628a. Poul, J., 63a. Prasolova, M. A., 961c. Pratt, I., 20m. Prévôt, R., 237a. Priebe, M. D., 529d. Prieto Lorenzo, A., 111a, 451a, 871. Progulske, D. R., 854b. Prügelhof, F., 545a. Pujatti, D., 139a, 344a, 676a, 821a. Pumarola Busquets, A., 135a. Purcallas, J., 587a. Purriel, P., 959c. Pustówka, T., 245c.

Qazilbash, N. A., 102a. Quenouille, M. H., 422a. Quesada, A., 207a.

Rachou, R. G., 800b. Radice, J. C., 765c. Raeburn, J., 195a, 195b, 223a. Ragab, M. M., 430d. Raghavan, N. G. S., 120h, 248a, 405b. Rahm, J., 899b, 899c. Rahn, E., 735d. Raigorodetskaya, E. A., 145b. Ramalingam, K., 784c. Ramanujachari, G., 221b. Ramírez, A. A., 758a. Ramírez, L. A., 758a. Ramírez Ramírez, E., 804a. Ramos, A., 758a. Ramos e Silva, J., 687a. Ranouil, 957a. Rao, K. A., 423c. Raski, D. J., 31a, 75b, 126g, 257q, 261b, 261d, 760a. Raspa, G., 514d. Rausch, R., 20ba, 21a, 132c, 428e, 613a. Raust, R., 197a. Ravens, J. R., 960a. Ravera, J. J., 959d. Rawson, D., 332a. Raymont, J. E. G., 97g. Rayski, C., 142a, 165b.

Read, C. P., 20c, 101b, 230bj. 230bk, 230bl, 230bu, 385b, 428b. Read, W. H., 431a. Reboud, E., 727d. Rebrassier, R. E., 28a, 835a. Rees, C. W., 230by. Rees, G., 29c, 29d. Reeves, P. A., 711a. Reichenbach-Klinke, H., H. 859a. Reichenow, E., 996. Reid, E., 97a. Reinertson, J. W., 385e. Reinhardt, F., 456a. Reisinger, E., 717a. Renjifo, S., 50b. Repetto, R. L., 672e. Reszke, H., 933a. Rey, L., 805a. Reynoldson, T. B., 519a. Ribeiro, H. M., 616b. Ribeiro, V. de M., 733a. Ricci, M., 34c, 275b, 514a, 785a, 785b, 785c, 785d, 785e. Ricci, N., 951a. Riedel, H., 720a. Rick, R. F., 65a, 348a. Rick, R. F., 65a, 548a. Riemens, J. M., 242a. Riley, W. A., 997. Rindi, G., 635c, 635d, 635c. Rinjard, J., 199a. Riordan, D., 143a. Ritchie, L. S., 38 l, 176f, 230r, 230s, 428j, 428q, 457a, 696f, 696h. Ritter, M., 256a. Rivera-Anaya, J. D., 17a, 896a. Rivero, E., 765d. Rizhikov, K. M., 9611, 9611, 961m, 961n, 961o. Rizzotti, G., 514e. Robert, M., 71a. Roberts, F. H. S., 65a, 345a. Roberts, J. H., 262a. Roberts, S. J., 81a. Robertson, G. H., 745a. Robinson, H. W., 230n. Rocca, E. D., 960c. Rocha, F. J. da, 798a. Rocha, U. F., 626a, 807b. Rodhain, J., 586s, 586u, 925b. Rodinò, D., 820c. Rodiriyues da Silva, L. 700c. Rodrigues da Silva, J., 799c, 801b, 801f. Rodríguez, C., 271a, 813c. Rodríguez Barrios, R., 6128, 960b. Rodríguez-Molina, R., 350a, 350b. Roe, R. J., 82a. Roedel, G., 798a. Rösler, O. A., 857a. Rogers, K. B., 699a. Rogers, R. T., 467d. Rogers, W. P., 66a.

Rohrbeck, E., 998.

Rojales, H., 758a. Roman, E., 359a, 426b. Romanov, I. V., 961bf. Romanov, N. P., 961b. Romić, S., 537a. Rosen, E., 53a. Rosenthal, A., 914a. Ross, G. L., 250a. Ross, I. C., 786a. Ross, I. C., 786a.
Ross, J. A., 639a.
Ross, W. M., 371b.
Roth, H., 165a.
Rothschild, M., 166.
Rousselot, R., 200b, 200c.
Rousset, P., 647b.
Row, P., 448a.
Roy, R., 248a, 405b.
Rubin, R., 95c.
Ruiz, J. M., 454a, 454b, 454c,
454d, 740a, 740b, 740c.
Ruiz Reves, F., 243a. Ruiz Reyes, F., 243a. Rukhlyadev, D. P., 961bg. Russell, C. M., 851a. Russell, J. C., 776e. Rutowitsch, M., 584a. Ryazantsev, V. F., 536j. Ryšavý, B., 866a, 964a. Saakova, E. O., 675f. Saavedra, J., 812a. Sabadini, L., 724a. Saccharin, H., 7a.

Sacco, G., 741a. Sachs, H., 397a. Sacks, M., 281b. Sadun, E. H., 230bn, 230bo,

230bp. Saggese, P., 822a. Saif, M., 430g. Saito, S., 702f. Sakaguchi, S., 107a. Sakoda, A., 753d. Saleh, A., 430c. Salvestroni, P., 629a. Samuel, M., 484a. Sanches, A., 399b. Sánchez Medal, L., 810a. Sandosham, A. A., 19a, 131a. San Julián, J., 612a, 960b. Sanmartin, C., 50b.

Sanmarin, C., 300.
Sano, T., 916a.
San Román, G. D., 577b.
Santos, A. G. dos, 586j.
Sanwal, K. C., 912d, 912e.
Sanyal, P. K., 690c.
Sar Dessai, M. S. N. P. R., 586t.
Sarkies, I. W. R., 68a, 152m. Sarkies, J. W. R., 68a, 152m, 334a.

Sarmento, A., 586e. Sarwar, M. M., 29e, 773a, 773b. Sasa, M., 409a. Sasaki, N., 905a. Sasser, J. N., 30k, 126d, 257b,

257m, 479a. Sassi, R., 950a. Sastry, S. A., 92b.

Sato, K., 396c, 409a. Sato, N., 827c. Satoo, T., 425b. Savary, A., 508a, 508b, 508c. Savaton-Pillet, J., 526a. Savel, J., 79a, 79b, 79d. Savino, L., 952b. Sawada, I., 864f. Sayad, W. Y., 2b. Scaff, L. M., 800g. Schack, J. A., 315a. Scheibe, G., 112a. Scheid, G., 294b, 737a. Schell, S. C., 281f, 428h. Schenk, P. J., 910a. Schieppati, E., 672c. Schiller, E. L., 20d, 21b, 99a, 150c, 428u.

Schlieper, C., 442a. Schmidt, J., 294b, 999. Schmidt, P., 327b. Schneider, P. A., 276a, 516a. Schnelle, G. B., 154d, 287a. Schobinger von Schowingen, R.,

308a. Schönberg, F., 23b. Scholten, C., 547a. Schreier, O., 475a.

Schroeder, A. H., 706b. Schubert, R., 838a. Schützler, G., 118c. Schulz, R. S., 961bd, 1002. Schulze, W., 118a, 118e. Schuster, M. L., 771a, 771b. Schuurmans-Stekhoven, jr., J.

H., 551b, 572a, 572b. Schwarte, L. H., 556. Schwartz, B., 20w, 239a, 678a. Schwarz, E., 334b.

Schwetz, J., 59c, 74b, 327e, 328h, 596a, 596b, 650a, 925a.

925a.
Schwöbel, W., 489a.
Sciacchitano, I., 595a, 610c.
Scofield, G. F., 230dc, 428s.
Scotza, J. V., 565a.
Scott, J. A., 20r, 20v, 230ba.
Scott, R. A., 466a.
Seale, C. C., 126f.
Sedgwick, C. E., 837a.
Seelkopf, K., 617a, 617b, 887a.
Seinhorst, J. W., 164i, 280a,

Seinhorst, J. W., 164i, 280a,

446a. Sekera, A., 899b, 899c. Selkon, J. M., 833a. Sell, O. E., 698a. Semenova, N. S., 145a. Semerdzhiev, P., 994. Semikin, V. I., **536e.** Senet, A., 27a, 121a. Senger, C. M., 428d, 428i. Senk, O., 535a. Serman, M., 797a. Serra García, E., 730a.

Sesnic, R., 499a. Sette, H., 799f.

Shah, K. S., 180d.

Shah, S. N., 689a. Shakir, M. H., 430h. Shalimov, A. A., 106b. Sharma, K. P., 180b. Shaw, J. N., 471a.

Shcherbovich, I. A., 536a. Shepherd, C. J., 463e. Sher, S. A., 261b.

Shibue, H., 696c. Shibuya, M., 246a. Shichijo, K., 696i.

Shikhobalova, N. P., 961d, 1002.

Shoho, C., 41d, 70a, 196a, 358a, 569a.

Shookhoff, H. B., 230b. Shope, R. E., 777e. Shorb, D. A., 230cl.

Short, G. E., 95a.

Short, G. E., 55a, 230db, 320b. Short, R. B., 55a, 230db, 320b. Siang, W. N., 480b. Sibalić, S., 850b. Siems, G., 61cc.

Siguier, F., 363a. Silva, G. M. da, 783d, 783f. Silva, N. N. de, 78a.

Silva, R., 352b, 944a. Silva, V. da, 586g. Silva C., R., 190d.

Silva Horta, J. da, 682b. Simitch, T., 909a. Simões, A., 624b.

Simonet, P., 182a. Sinclair, L. R., 230p, 230q.

Singh, J., 405b. Singh, K. S., 688a. Sioufi, H., 482b. Sirot, L., 574i.

Sirot, L., 574.
Sivec, S., 570b.
Skarbilovich, T. S., 961bj.
Skoglund, W. C., 858a.
Skryabin, K. I., 118f, 961a,
1000, 1001, 1002.
Sleeth, B., 133a, 779a.
Slobodskaya, R. A., 145b.

Smiles, J., 38f. Smirnova, K. A., 438a.

Smit, R., 143a.

Smit, R., 143a.
Smith, A. L., 578c.
Smith, D. A., 152h, 152 l, 358b.
Smith, H. W., 563a.
Smith, R. J., 230ce.
Smith, W. A., 263a.
Smith, W. N., 261k.
Smith, W. T., 193a.
Smyth, J. D., 226a.
Snow, D. 441a.

Snow, D., 441a. Snyder, C. H., 124a. Sobolev, A. A., 961ba, 961bb. Sodeman, W. A., 94a, 105b. Soguri, M., 695a.

Sohier, H. M. L., 362d. Sokolov, V. M., 849d. Soliman, K. N., 196b, 196c,

287ь. Solomon, C., 467c. Soltys, A., 329a.

Soo-Hoo, G., 134a, 385d. Soots, 607a. Sopelchenko, M. I., 917b, 917d. Soprunov, F. F., 1142. Soprunova, N. Y., 1142. Soria, M. F., 130a. Sosa Bens, D., 136a, 946b. Sotolongo, F., 270e, 270f, 294c. Sousa Santos, J. do C. de, 586p. Southey, J. F., 258a. Southwell, B. L., 206a. Southwick, M. D., 488a. Spasski, A. A., 961e, 961f, 961g, 961h, 961i. Specht, A. W., 128a. Spedding, C. R. W., 97c, 214a, 539a. Spencer, E. L., 776g. Sprau, F., 476a. Sprehn, C., 903a. Sprent, J. F. A., 98d, 132e, 152f, 4636 Spruyt, F. J., 125b. Srepel, B., 570a. Srivastava, H. D., 29f. Stabler, R. M., 20z. Stalfoort, T. G. J., 774a. Stammer, H. J., 555a, 848b. Standaert, A., 464a. Standen, J. H., 480a. Standen, O. D., 19b, 38g, 61b, 152j. Staniland, L. N., 26a, 164j. Stauffer, H., 670b. Steele, J. H., 600a. Steiner, G., 257r, 776a. Stephenson, W., 18a, 18b. Sterman, M. M., 230b. Stevens, N. E., 558. Stevens, R. B., 558. Stewart, T. B., 698a. Stijns, J., 58h. Stirewalt, M. A., 20a, 230cs, 230ct, 424b. Stoddard, E. M., 478c. Stöckli, A., 860a. Stoffberg, N., 833b. Stojanović, V., 954a. Stokes, W. M., 383a. Stone, L. E. W., 26a. Stone, W. M., 129a. Stone, W. S., 2a. Stoner, R. D., 775a. Stones, P. B., 544c. Stoppa, 574d. Stoyanovitch, V., 701b. Strassen, O. zur, 848a. Strenzke, K., 859b. Stryzecka-Kstupska, A., 525a. Stunkard, H. W., 101f. Sudarikov, V. E., 961i, 961o, 961p. Suguro, T., 696g, 696i. Sullivan, J. F., 471a. Sultan, Z. M., 452a. Sumarokov, A. V., 953d.

Supperer, R., 157a, 832a.
Surrey, A. R., 413a.
Susman, M. P., 188a.
Susman, O., 777a.
Suter, C. M., 413a.
Suzacq, C. V., 587a, 612b.
Suzuki, R., 719a.
Swales, W. E., 10a, 10b, 10c.
Swank, jr., G., 776d.
Swanson, L. E., 20h, 154a, 644a, 772a.
Sweatman, G. K., 656a.
Szafiarski, J., 453d.
Szewczak, J. T., 696f.
Szidat, L., 130a, 500a.
Szkutnik, Z., 453a.

Tagle V., I., 352a, 352f. Takahashi, S., 926a. Takeyama, O., 753b, 753c. Takeya-Sikô, 706a. Taleb, N., 705b. Tamargo I., A., 352c. Tanabe, H., 38 l. Tanaka, I., 919a. Tandon, R. S., 912a. Tarjan, A. C., 301, 30p, 30q, 261a, 478a, 478b, 795a. Taufflieb, R., 74a.
Taylor, A. L., 126a, 257m,
257s, 311a, 479d, 678b.
Taylor, E. L., 51a, 120b, 165g, 1003. Tedder, P. W., 154c. Teixeira, E., 799f. Teixidó Vaillant, S., 946d. Tesseraux, H., 863a. Tetley, J. H., 264a.
Thames, jr., W. H., 126f, 776h.
Thapar, G. S., 912a. Théodorides, J., 327d, 514b, 852a, 983a. Thetford, N. D., 54a. Thillet, C. J., 230cx. Thomas, D., 640a. Thomas, L. J., 230br, 261]. Thompson, F. A., 531c.
Thompson, F. A., 531c.
Thornton, G. D., 776f.
Thornton, H. Thornton, H., 303. Threlkeld, W. L., 187b. Thurston, J. P., 152g, 357a. Tigertt, W. D., 696h. Tillier, H., 233h, 576e. Timmons, H. F., 20f. Tinaz, A., 963a. Tiner, J. D., 230ca, 461a, 1004. Tisdale, W. B., 126h. Tjallingii, F., 242a. Tocheport, G., 327b.
Todd, A. C., 13e, 40a, 87d, 129a, 150a, 177a, 198b.

Tolentino, P., 743a.
Tolosa, E., 496a.
Tomalino, D., 959c.
Tomažič, G., 570b.
Tomberg, V., 328i.
Tompkins, V., 467d.
Tonelli, L., 609a.
Toulant, P., 367f, 893a.
Tournoux, P., 327b.
Toxopeus, H. J., 463d.
Tracey, M. V., 559.
Traub, R., 26d.
Tredre, R. F., 435a.
Trelles, J. O., 960c.
Tricoire, J., 576a.
Trincão, C., 586i, 762a.
Trincão, C., 586i, 762a.
Trivedi, B. P., 403a.
Tromba, F. G., 261k, 428o.
Trottmann, M., 759a.
Tsuruta, K., 396b.
Tsutsunava, M. N., 953b.
Tugwell, R. L., 230a.
Turk, R. D., 240a.
Turner, J. H., 230p, 230q.
Turpin, R., 526a.
Tuzet, O., 361a.
Twente, G. E., 602a.
Tymniak, M., 113b, 453b.

Üner, R., 845a, 845b.
Ullrich, K., 528b.
Ulmer, M. J., 323a, 693a, 754a.,
Umanzio, C. B., 708a.
Ungar, H., 317a.
Ungar, R., 317a.
U.S. Department of Agriculture,
560, 561, 562.
U.S. Livestock Sanitary Association, 486a, 777c.
Urbain, A., 199a.
Uspenskaya, A. V., 675e.
Ussing, H., 679a.
Uteshev, A. I., 536k.
Uzmann, J. R., 20bc.

Vaidyanathan, S. N., 407a.
Vajrasthira, S., 230bn, 230bo, 230bp,
Valach, V., 928a.
Valdés Diaz, R., 136a, 270d, 946b.
Valladolid del Val, A., 732a.
Valle, D. del, 634b.
Valleau, W. D., 86a, 198a.
Valledor, R., 946a.
Vanbreuseghem, R., 596c.
Van Burkalow, A., 88a.
Van Cleave, H. J., 20f, 87e, 132a, 230bv, 524a.
Vandenberghe, J., 59d.
Van der Woude, A., 101a.
Vannfält, K. A., 571a.
Vanni, V., 765c.
Vargas, L., 243c, 450a, 450b.
Vasilev, A. A., 536i.

Vasilkova, Z. G., 906b, 906c, 900d, 906e, 906f. Vasudeva, R. S., 218a. Vaughan, A. W., 155h. Vaughn, C. M., 176d. Vaughn, J., 231a. Vavroušková, K., 964b. Vecht, J. van der, 474a. Vega, J., 401a. Vega Fernández, S. de, 496b. Vegors, H. H., 698a. Veitsman, L. N., 961n. Velichkin, P. A., 961bl. Velimirović, S., 535b. Venturiello, J. J., 672d. Vera, V. R. de, 577c. Verbruggen, J., 328e. Vercruysse, R., 542a, 855a. Vergapi F. 620c. Vergani, F., 630a. Vergoz, 574a, 574f, 768d. Vermeil, C., 327b. Verna, F., 48a. Vernberg, W. B., 101d, 230f. Vernier, J., 362c. Verspyck, R., 710a. Vesell, H., 315a. Viallet, M., 233i. Viallet, P., 233i. Viana Martins, A., 799b. Viehmann, H., 863a. Vieira, A. E., 783h. Vilela, M. P., 343a. Villalpando del Valle, E., 353a. Villaret, B., 830a.
Villaret, B., 830a.
Villaret, O. T. de, 388a.
Vodopivec, S., 570b, 968a.
Voge, M., 20g, 39a, 428g, 529e.
Vogel, K., 671a.
Vogel, Tyber, M. 361a. Vogeli, R., 9/16. Vogelis Zuber, M., 361a. Vogelsang, E. G., 813a, 813b, 813c, 813d. Volk, G. M., 202a. Von Brand, T., 87a, 230cq, 304. Von Mechow, J., 257k.

Wachek, F., 555a, 1006. Wade, L. L., 177c. Wagner, F., 476b. Wagner, O., 148a. Wahlin, B., 533a. Wakayama, S., 827c.

Vuyadinovitch, B., 701b.

Walker, A. R. P., 36a, 143a. Wallace, F. G., 997. Walton, A. C., 230bj. Wang, C. K., 2e. Wanson, M., 596c. Ward, H. L., 104a, 104b, 150e, Wardle, R. A., 305. Warren, B. H., 428 1. Watson, J. M., 105c, 227a, 229a, 235b, 605a. Webber, W. A. F., 357a. Weber, A. P., 120c. Weber, J. E., 582a. Weber, J. R., 764a. Wehr, E. E., 230w. Weinbach, E. C., 87a. Weinsheimer, P. A., 1007. Weinstein, P. P., 2f, 87a, 230bt, 385c. Weissman, F. M., 378a. Wells, H. D., 126b. Wells, H. S., 98a, 98c. Wells, J. C., 126b. Welti, J. J., 363a. Werckmeister, G., 1008. Wertejuk, M., 113a. Wesenberg-Lund, E., 1009. Wetzel, R., 83a, 156a, 295a, 377a. Weyer, F., 996. Whelan, J. P. F., 22a. White, E. G., 356a. Whitehead, H., 462a. Whitlock, J. H., 81a, 81b, 209b. Whitten, L. K., 165c. Willers, L. R., 100c.
Wieser, W., 100a.
Wilser, W., 100a.
Wilhelm G., O., 633a.
Wilkinson, F., 672e.
Willers, E. H., 471b.
Williams T. H., 655a, 844a. Williams T. H., 655a, 844a.
Williamson, J., 152d, 334d, 334g.
Willmott, S., 97h, 97j.
Wilmanns, H., 210a.
Wilson, A. L., 155e.
Wilson, J. F., 278a.
Wilson, J. K., 703a.
Winckel, W. E. F., 674a, 674b.
Winter, H. A. 150e. Winter, H. A., 150e. Winterhalter, M., 850d. Wisniowski, J., 453d. Wisseman, jr., C. L., 26d. Witter, W. F., 433a.

Wong, C. C., 531b. Woodruff, A. W., 289c, 358d. Worley, D. E., 252a. Wu, C. L., 659a. Wyant, Z. N., 13e. Wykoff, D. E., 230r. Wysocka, F., 653a.

Yakhnina, R. Z., 834a.
Yamaguti, S., 49a, 568a.
Yamamoto, S., 721a.
Yamane, S., 425b.
Yamao, Y., 864b, 864c, 864d,
864e, 864g, 915a.
Yamaoka, K., 696g, 696i.
Yamashita, J., 926a.
Yarinsky, A., 104c.
Yasuda, M., 396b.
Yasuda, S., 396e.
Yasudaya, M., 702j.
Yeager, L. B., 940a.
Yokogawa, M., 381, 176f, 696b,
696f, 696g, 696i, 6961.
Yokoyama, S., 601b.
Yosufzai, H. K., 26c, 658a.
Young, R. T., 230 1, 436a.

Zach, O., 115a.
Zadura, J., 245e.
Zaffino, C., 514d.
Zahawi, S. el, 648a.
Zahra, A., 289e.
Zaiman, H., 20bf.
Zaky, H. A., 212a.
Zaleski, J., 245a, 245b, 245d, 453c.
Zamith, A. P. L., 796a.
Zarapico Romero, M., 267a.
Żarnowski, E., 113a.
Zaskind, L. N., 961br.
Zaverdinos, A., 83ob.
Zerecero y D., M. C., 588c.
Zettl, J., 751a.
Zhuk, A., 849a.
Zimmerman, L. E., 706a.
Zlokas, D., 668a.
Zobrist, L., 506a, 517a, 828a.
Zogović, B., 954a.
Zoubek, V., 934a.
Zulueta, J. de, 50b.
Zunino, A., 672a.
Zwillenberg, L. O., 120c.
Zylka, N., 969b.

INDEX OF PERIODICALS

(The reference is to the serial number, not to the page.)

Abstracts of Doctoral Dissertations. Ohio State University, 563.
Abstracts of Doctoral Dissertations. University Anais do Instituto de Medicina Tropical. Lisbon, 321, 586. Anais Paulistas de Medicina e Cirurgia, 322. Anales de la Facultad de Medicina. of Nebraska, 564. Acta Agronomica. Palmira, 306. sidad de Lima, 882. Anales de la Facultad de Medicina de Monte-Acta Anatomica. Basle, 47. video, 587. Anales del Instituto de Biología. Mexico, 588. Acta Biologica Venezuelica, 565. Acta Medica Italica di Malattie Infettive e Parassitarie, 48, 566. Acta Medica Philippina, 307. Anales de la Real Academia Nacional de Medicina. Madrid, 589. Anatomical Record, 323, 590. Acta Medica Scandinavica, 167. Anesthésie et Analgésie. Paris, 591. Animal Health Leaflet. Ministry of Agriculture Acta Medica Scandinavica. Supplementum, 168, and Fisheries. London, 178, 324. Acta Medicinae Okayama, 49, 568. Acta Neurologica et Psychiatrica Belgica, 569. Acta Pharmaceutica Jugoslavica, 570. Acta Societatis Medicorum Upsaliensis, 571. Annales Historico-Naturales Musei Nationalis Hungarici, 56, 592. Annales de l'Institut National de la Recherche Acta Tropica. Basle, 50, 308. Agronomique. Série C. Annales des Épiphy-Acta Unio Internationalis contra Cancrum. ties, 57, 325. Annales de l'Institut Pasteur. Paris, 326. Louvain, 879. Acta Veterinaria. Belgrade, 880. Acta Veterinaria. Budapest, 309. Acta Zoologica Lilloana. Tucumán, 572. Annales de Médecine Vétérinaire, 593. Annales Médico-Psychologiques, 594.

Annales du Musée Royal du Congo Belge.

Série in 8vo, Sciences Zoologiques, 595. Advisory Leaflet. Department of Agriculture, Scotland, 169. Annales de Parasitologie Humaine et Comparée, 58, 327. Arztliche Wochenschrift. Berlin, 170, 310, 573. Annales de la Société Belge de Médecine Tropi-Afrique Française Chirurgicale, 574. Agricoltura Napoletana, 881. cale, 59, 328, 596. Annales Universitatis Mariae Curie-Skłodowska, Agricultural Chemicals. Baltimore, 311. Agricultural Gazette of New South Wales, 171, Lublin, 329 Annali della Facoltà di Medicina Veterinaria. Pisa, 597. Annali di Medicina Navale e Tropicale, 330. Agricultural Ireland, 313.
Agricultural Journal. Department of Agriculture, Fiji, 172, 314. Agricultural Research. Taiwan, 575. Annali della Sanità Pubblica, 598. Annals of Applied Biology, 60, 331. Annals of the Entomological Society of America, Agriculture. London, 51. Agriculture Pratique. Paris, 173. Annals of Internal Medicine, 600 Algérie Médicale, 576. Almanaque del Ministerio de Agricultura y Annals and Magazine of Natural History, 332. Ganadería. Buenos Aires, 577. Annals of the New York Academy of Sciences, American Fruit Grower, 174. American Heart Journal, 175, 315.

American Journal of Clinical Pathology, 578.

American Journal of Hygiene, 1, 52, 316.

American Journal of the Medical Sciences, 53. Annals of the Phytopathological Society of Japan, 601. Annals of Surgery, 602. Annals of Tropical Medicine and Parasitology. American Journal of Medicine, 579. 61, 179, 334. Annotationes Zoologicae Japonenses, 603. American Journal of Nursing, 580. American Journal of Ophthalmology, 581. Annual Review of Microbiology, 335. American Journal of Pathology, 317. Antiseptic. Madras, 180. American Journal of Surgery, 582. American Journal of Tropical Medicine and Anzeiger für Schädlingskunde, 336. Archiv für Experimentelle Veterinärmedizin, 883. Archiv für Hygiene und Bakteriologie, 181, 604.
Archiv für Klinische Chirurgie vereinigt mit
Deutsche Zeitschrift für Chirurgie, 884. Hygiene, 2, 54, 176, 318. American Journal of Veterinary Research, 177, 319. American Midland Naturalist, 55, 320. Anais da Academia Brasileira de Ciencias, 583. ives d'Anatomie Microscopique et de Morphologie Expérimentale. Paris, 62. Anais Brasileiros de Dermatologia e Sifilografia, Archives de Biologie. Paris, 337. Archives of Dermatology and Syphilology, 338.

Archives of Disease in Childhood, 605.

Archives Françaises de Pédiatrie, 606.

584.

Anais da Faculdade de Medicina da Univer-

sidade de São Paulo, 585.

Archives de l'Institut Pasteur d'Algérie, 63, 182. Archives de l'Institut Pasteur du Maroc, 339. Archives des Maladies de l'Appareil Digestif et

des Maladies de la Nutrition, 607. Archives de Médecine Générale et Tropicale, 608.

Archives of Ophthalmology, 183.

Archives of Pathology, 184.
Archivio "De Vecchi" per l'Anatomia Patologica
e la Medicina Clinica, 609.

Archivio Italiano di Scienze Mediche Tropicali e di Parassitologia, 340.

Archivio Zoologico Italiano, 610. Archivos de Medicina Infantil, 185.

Archivos de Pediatría del Uruguay, 186, 611. Archivos Uruguayos de Medicina, Cirugia y

Especialidades, 612. Arctic. Montreal, 613. Arizona Farmer, 885. Arkansas Farm Research, 341.

Arkiv för Zoologi, 342. Arquivos de Biologia. São Paulo, 343. Arquivos Brasileiros de Medicina, 614.

Arquivos Brasileiros de Medicina Naval, 615. Arquivos Médicos Municipais. São Paulo, 616. Arsskrift. Kalmar Läns Södra Hushållningssällskap, 886.

Arzneimittel-Forschung. Aulendorf, 617, 887. Atti della Accademia Nazionale dei Lincei. Rendiconti. Classe di Scienze Fisiche, Matematiche e Naturali. Rome, 64. Atti della Società Italiana di Scienze Naturali

e del Museo Civico di Storia Naturale in

Milano, 344. Atti della Società Italiana delle Scienze Veterin-

arie, 618. Auburn Veterinarian. Alabama, 187, 888. Australasian Annals of Medicine, 619.

Australian Journal of Agricultural Research, 3,

Australian Journal of Science, 345. Australian Journal of Scientific Research. Series B, Biological Sciences, 66, 346.

Australian and New Zealand Journal of Surgery, 188, 620.

Australian Plant Disease Recorder, 347. Australian Veterinary Journal, 67, 348.

Beiträge zur Pathologischen Anatomie und zur Allgemeinen Pathologie, 4.

Bimonthly Bulletin. North Dakota Agricultural

Experiment Station, 189. Biochemical Journal, 621. Biochemische Zeitschrift, 349. Biologie Médicale, 622.

Bitki Koruma Bülteni, 623.

Boletim Cultural da Guiné Portuguesa, 624. Boletim. Directoria da Produção Animal. Rio Grande do Sul, Brazil, 889.

Boletim da Sociedade de Estudos de Moçam-

bique, 625. Boletim da Sociedade Paulista de Medicina Veterinária, 626.

Boletín de la Asociación Médica de Puerto Rico, 350.

Boletín de la Cámara Oficial Sindical Agraria de Barcelona, 890.

Boletín. Centro Nacional de Investigación y Experimentación Agrícola de La Molina. Lima, 351.

Boletin de Información. Consejo General de Colegios Veterinarios de España, 627. Boletín de Informaciones Parasitarias Chilenas,

Boletín Informativo. Chinchiná, Colombia, 191. Boletín Informativo. Minsterio de Ganadería y Agricultura, Uruguay, 628.

Boletin del Instituto de Ciencias Naturales. Quito, 629.

Boletin del Instituto de Investigaciones Veterinaris. Caracas, 630.

Boletín Médico del Hospital Infantil. Mexico,

353, 891. Boletín Mensual. Dirección de Ganadería, Montevideo, 631.

Boletín de la Oficina Sanitaria Panamericana,

Boletín de Sanidad Militar. Mexico, 892. Boletin de la Sociedad de Biologia de Concepción (Chile), 633.

Boletines y Trabajos. Academia Argentina de Cirugía, 634.

Bollettino della Società Italiana di Biologia Sperimentale, 635.

Bollettino della Stazione di Patologia Vegetale.

Rome, 354. Bonner Zoologische Beiträge, 192.

Bragantia. Campinas, 636.

Brain, 193. Brasil-Medico, 355, 637.

British Agricultural Bulletin, 194.

British Farmer, 5.
British Journal of Nutrition, 356. British Journal of Ophthalmology, 68.

British Journal of Pharmacology and Chemotherapy, 195, 357, 638. British Journal of Radiology, 639.

British Journal of Surgery, 640. British Journal of Tuberculosis and Diseases of the Chest, 641.

British Medical Journal, 6, 69, 358. British Veterinary Journal, 70, 196.

Bulletin de l'Académie Nationale de Médecine.

Paris, 893. Bulletin de l'Académie Vétérinaire de France, 197.

Bulletin Biologique de la France et de la Belgique, 71, 359.

Bulletin of the British Museum (Natural History).

Zoology, 72. Bulletin de la Chambre d'Agriculture de Casa-

blanca, 894. Bulletin of the Faculty of Agriculture, Kagoshima

University, 642. Bulletin. Fisheries Research Board of Canada,

Florida Agricultural Experiment Bulletin. Stations, 202, 644.

Bulletin Horticole. Liége, 360.

Bulletin de l'Institut Français d'Afrique Noire,

Bulletin de l'Institut d'Hygiène du Maroc, 645. Bulletin de l'Institut Océanographique. Monaco, Bulletin. Kentucky Agricultural Experiment Station, 198. Bulletin of Marine Science of the Gulf and

Caribbean, 646.

Bulletin Médical de l'Afrique Occidentale Française, 362, 647. Bulletins et Mémoires de la Société Médicale

des Hôpitaux de Paris, 363, 648.

Bulletin of the Ministry of Agriculture and

Fisheries. London, 364.
Bulletin du Muséum National d'Histoire
Naturelle. Paris, 199.
Bulletin of the Naniwa University. Series B.
Agricultural and Natural Science, 895. Bulletin of the National Institute of Agricultural Sciences. Chiba. Series G. Animal Hus-

bandry, 649. Bulletin. New York State Flower Growers Incorporated, 365.
Bulletin, Puerto Rico Agricultural Experiment

Station, Rio Piedras, 896. Bulletin des Séances. Institut Royal Colonial

Belge, 650.

Bulletin de la Société Française de Dermatologie

et de Syphiligraphie, 366. Bulletin de la Société Neuchâteloise des Sciences

Naturelles, 651. Bulletin de la Société de Pathologie Exotique,

7, 74, 200, 367. Bulletin de la Société Vétérinaire Hellénique, 652. Bulletin de la Société Zoologique de France, 368.

Bulletin of the State Institute of Marine and Tropical Medicine in Gdańsk, Poland, 201,

Bulletin of the World Health Organization, 654. Bulletin of Zoological Nomenclature, 8, 369.

Cahiers des Ingénieurs Agronomes, 897. California Agriculture, 9, 75. California Fish and Game, 76. Canadian Field-Naturalist, 370. Canadian Insect Pest Review, 203. Canadian Journal of Comparative Medicine and Veterinary Science, 10, 77, 204.
Canadian Journal of Medical Sciences, 371.
Canadian Journal of Medical Technology, 655.
Canadian Journal of Public Health, 11, 656. Canadian Journal of Zoology, 372. Caribbean Medical Journal, 657. Cellule, 658. Československá Dermatologie, 898. Československá Farmacie, 899. Ceylon Journal of Science. Section B. Zoology, Chacra. Revista Mensual de Agricultura, Ganadería e Industrías Anexas. Buenos Aires,

Chemistry and Industry. London, 205. Chinese Medical Journal, 659.

Chirurg, 660.

Chrysanthème, 661.

Circular. Georgia Coastal Plain Experiment Station, 206.

Illinois Agricultural Experiment Station, 662.

Circular. Texas A. & M. College Extension Service, 663.

Cirugia y Cirujanos. Mexico, 901.

Citrus Leaves, 664.

Clínica y Laboratorio, 665. Clinica Veterinaria. Milan, 207, 666.

Comptes Rendus des Séances de l'Académie des

Sciences. Paris, 12, 79. Comptes Rendus des Séances de la Société de

Biologie. Paris, 80, 208. Cornell Extension Bulletin. New York State: College of Agriculture, 373, 667.

Cornell Veterinarian, 81, 209, 374 Courrier de l'Institut d'Hygiène. Belgrade, 668. Cyprus Medical Journal, 82.

Danish Review of Game Biology, 669.

Dermatologica. Basle, 375, 670. Deutsche Gesundheitswesen (Das), 671.

Deutsche Landwirtschaft, 376.

Deutsche Landwirtschaftliche Presse, 902.

Deutsche Medizinische Wochenschrift, 210.

Deutsche Pelztierzüchter (Der), 211.

Deutsche Tierärztliche Wochenschrift, 83, 377. Deutsche Wirtschaftsgeflügelzucht, 903.

Día Médico. Buenos Aires, 672.

Diseases of the Chest. Chicago, 212.

Diseases of the Nervous System, 378, 673.

Documenta de Medicina Geographica et Tropica.

Amsterdam, 84, 379, 674.

Doelmatige Veehouderij, 904.

Dokladi Akademii Nauk SSSR, 213, 380, 675.

Doriana, Genoa. [Supplement to Annali del Museo Civico di Storia Naturale "G. Doria".], 381, 676.

East African Medical Journal, 85, 382. Economic Botany. Lancaster, Pa. 86.

Empire Journal of Experimental Agriculture, 214, 383.

Endeavour. London, 215.

Ergebnisse der Wissenschaftlichen Untersuchung

des Schweizerischen Nationalparks, 384. Experimental Parasitology. New York, 13, 87, 385.

Experimental Report of Government Experimental Station for Animal Hygiene, Tokyo,

Extension Bulletin. Hawaii College of Agriculture, 14.

FAO Development Paper. Rome, 677. FAO Food and Agricultural Legislation. Rome,

FAO Plant Protection Bulletin. Rome, 216, 387.

Farmer and Stock-Breeder, 15.
Farmers' Bulletin. U.S. Department of Agriculture, 678.

Farming in South Africa, 388. Feldsher i Akusherka. Moscow, 906. Feuille des Naturalistes. Paris, 389. Field and Laboratory. Dallas, Texas, 390.

Flora og Fauna, 679.

INDEX OF PERIODICALS

Flugblatt. Bundesanstalt für Pflanzenschutz. Vienna, 907. Folia Clinica et Biologica, 391, 680. Frontiers. Philadelphia, 908. Fruits d'Outre Mer. Paris, 392.

Gartner-Tidende, 681.
Gazeta Médica Portuguesa, 682.
Geburtshilfe und Frauenheilkunde, 683.
Geographical Review. New York, 16, 88.
Gesunde Pflanzen. Frankfurt, 393, 684.
Giornale di Scienze Mediche, 685.
Glas Srpske Akademije Nauka. Odeljenje
Medicinskih Nauka, 909.
Glasgow Medical Journal, 217.
Glasgow Naturalist, 89.
Groenten en Fruit, 910.
Grower. London, 394.

Harefuah, 90.
Hemera Zoa. Buitenzorg, 395.
Hirosaki Medical Journal, 396, 686.
Höfchen-Briefe für Wissenschaft und Praxis, 397.
Hoppe-Seyler's Zeitschrift für Physiologische Chemie, 398.
Hospital. Rio de Janeiro, 399, 687.

Hygiène et Médecine Scolaires. Paris, 400.

Idia. Buenos Aires, 401.
Igaku Kenkyuu. Kyushu University, 402.
Indian Farming, 218.
Indian Journal of Child Health, 911.
Indian Journal of Helminthology, 688, 912.
Indian Journal of Medical Sciences, 219, 403, 689.
Indian Journal of Surgery, 404.
Indian Medical Gazette, 91, 220, 405, 690.
Indian Medical Record, 691.
Indian Physician, 406.
Indian Veterinary Journal, 92, 221, 407.
Informatore Fitopatologico. Bologna, 692.
Iowa State College of Science, 693.
Iowa Veterinarian, 93.
Irish Naturalists' Journal, 408.
Istanbul Universitesi Tip Fakültesi Mecmuasi,

Japanese Journal of Experimental Medicine, 409. Japanese Journal of Ichthyology, 695. Japanese Journal of Medical Science and Biology, 410, 696. Japanese Journal of Pharmacology, 411.

Japanese Journal of Plant Protection, 913.

Jornal do Médico. Oporto, 697.

Journal of Agriculture of Western Australia,

Journal of Agriculture of Western Australia, 412. Journal of the Albert Einstein Medical Center.

Philadelphia, 914.

Journal of the American Chemical Society, 413.

Journal of the American Medical Association, 94.

Journal of the American Veterinary Medical

Association, 17, 95, 222.

Journal of Animal Science, 698.

Journal of the Bengal Natural History Society, 414.

Journal of Biological Chemistry, 415.

Journal of Cellular and Comparative Physiology, 416.

Journal of the Chemical Society. London, 223, 417.

Journal of the Christian Medical Association of India, Pakistan, Burma and Ceylon, 224.

Journal of Clinical Pathology. London, 699.

Journal of the College of Arts and Sciences, Chiba University. Natural Science Series, 915.

Journal du Conseil Permanent International pour l'Exploration de la Mer. Copenhagen, 418.

Journal du Conseil Permanent International pour l'Exploration de la Mer. Copenhagen, 418. Journal of the Department of Agriculture. Dublin, 700.

Journal of the Department of Agriculture.

Victoria, 96.

Journal of the Elisha Mitchell Scientific Society, 225, 419. Journal of Experimental Biology, 18, 226.

Journal of Experimental Biology, 18, 220.

Journal of the Faculty of Medicine of Baghdad,

Iraq, 227, 420.

Journal of the Fisheries Research Board of Canada, 421.

Journal of Helminthology, 19, 97, 422.

Journal of the Indian Medical Association, 423.

Journal of Infectious Diseases, 98, 424.

Journal International de Chirurgie, 701.

Journal of the International College of Surgeons,

425.
Journal of the Japanese Veterinary Medical
Association, 702.
Journal of Laboratory and Clinical Medicine, 703.

Journal of Mammalogy, 99. Journal of the Marine Biological Association of

the United Kingdom, 100, 228.

Journal de Médecine de Lyon, 426.

Journal of the Medical Association of Formosa, 704. Journal Médical Libanais, 229, 705.

Journal of the National Cancer Institute, 427.
Journal of Nervous and Mental Disease, 706.
Journal of Neurosurgery. Springfield, Illinois,

Journal of Osteopathy, 708. Journal of Parasitology, 20, 101, 230, 428. Journal of Pathology and Bacteriology, 231. Journal of Pediatrics, 232.

Journal of Pediatrics, 232.

Journal of Pediatrics, 232.

London,

Journal of the Philippine Medical Association, 429.

Journal de Radiologie et d'Electrologie, 233, 710. Journal of the Royal Army Veterinary Corps, 234, 711. Journal of the Royal Egyptian Medical Associa-

tion, 103, 430.

Journal of the Royal Welsh Agricultural Society,

Journal of the Science of Food and Agriculture.
London, 431.

Journal of Science of the Hiroshima University. Series B, Division I. (Zoology), 432. Journal of Sericultural Science. Tokyo, 916. Journal of Small Animal Medicine. Inglewood, Calif., 433.

Journal of the South African Veterinary Medical

Association, 434.

Journal of the Tennessee Academy of Science,

104, 713. Journal of the Tokyo University of Fisheries, 714. Journal of Tropical Medicine and Hygiene, 105,

235, 435. Journal d'Urologie Médicale et Chirurgicale,

715. Journal of Urology, 716.

Journal of the Washington Academy of Sciences, 21, 236, 436.

Kärtner Bauer, 717 Karakulevodstvo i Zverovodstvo, 917. Keio Journal of Medicine, 437. Khirurgiya. Moscow, 106, 438. Kinderärztliche Praxis, 718. Kitakanto Medical Journal, 719. Kitasato Archives of Experimental Medicine, Klinicheskaya Meditsina. Moscow, 439, 918. Klinische Wochenschrift, 237, 440, 720. Kyushu Agricultural Research, 721, 919.

Lancet, 22, 108, 441. Landwirtschaftliche Forschung, 442. Landwirtschaftliches Wochenblatt für Westfalen und Lippe, 722. Landwirtshaftsblatt Weser-Ems, 920. Lantmannen, 921. Leaflet. Kentucky College of Agriculture, 723. Ministry of Agriculture, Northern Ireland, 109, 238. Leaflet. United States Department of Agriculture, 239. Lebensmitteltierarzt. [Supplement to Deutsche Tierärztliche Wochenschrift.], 23. Lunds Universitets Arsskrift, 443. Lyon Chirurgical, 724. Lyon Médical, 725.

M.S.C. Veterinarian. Michigan State College, Magyar Állatorvosok Lapja, 726. Maroc Médical, 444. Marseille Chirurgical, 727. Médecine Tropicale, 241, 445, 728. Mededelingen. Directeur van de Tuinbouw. 's-Gravenhage, 242, 446. Mededelingen van de Landbouwhogeschool en de Opzoekingsstations van de Staat te Gent, 447. Medical Journal of Australia, 24, 110, 448. Medical Journal of Malaya, 449. Medical Press, 729. Medicamenta. Madrid, 730. Medicina. Revista Mexicana, 243, 450, 731. Medicina y Cirugía. Bogotá, 922.

Medizinische. Stuttgart, 735. Medizinische Klinik, 112, 244, 736. Medizinische Monatsschrift. Stuttgart, 737. Medycyna Weterynaryjna, 113, 245, 453. Mémoires de l'Académie de Chirurgie. Paris, Mémoires de l'Institut Royal Colonial Belge .: Section des Sciences Naturelles et Médicales, Mémoires de l'Institut Scientifique de Madagascar. Série A. Biologie Animale, 739 Memoirs of the Faculty of Agriculture, Hokkaido University, 926. Memoirs of the Faculty of Agriculture, Kago-shima University, 246. Memorias do Instituto Butantan, 454, 740. Merkblätter über die Hauptsächlichsten Parasiten und Parasitengruppen, 927. Mikrobiologiya. Moscow, 114, 455. Mikrokosmos, 25, 115, 456. Mikroskopie. Vienna, 116. Military Surgeon, 457. Minerva Chirurgica. Turin, 741. Minerva Medica, 742. Minerva Pediatrica. Turin, 743. Mitteilungen aus der Biologischen Zentralanstalt für Land- und Forstwirtschaft, 117. Monatshefte für Praktische Tierheilkunde, 247. Monatshefte für Veterinärmedizin, 118, 458. Münchener Medizinische Wochenschrift, 459. Mycologia, 460. Nachrichtenblatt für den Deutschen Pflanzenschutzdienst. Berlin, 119. Natural History Miscellanea. Chicago, 461. Naturalist. London, 462. Nature. London, 26, 120, 248, 463. Nature. Paris, 27, 121, 249. Natuurwetenschappelijk Tijdschrift, 464. Nederlandsch Tijdschrift voor Geneeskunde, 465. Neurologie a Psychiatrie Československá, 928. New England Journal of Medicine, 250, 466. New York State Journal of Medicine, 467. New Zealand Entomologist, 744. New Zealand Journal of Agriculture, 468. New Zealand Journal of Science and Technology-B. General Research Section, 469. New Zealand Medical Journal, 745. Nordisk Medicin, 470. North American Veterinarian, 28, 251, 471. Northwest Medicine, 746. Northwest Science, 122.

Notiziario sulle Malattie delle Piante. Milan, 747.

Notulae Naturae of the Academy of Natural

Nuovi Annali d'Igiene e Microbiologia. Rome,

Sciences of Philadelphia, 472.

Medicina y Cirugia de Guerra. Madrid, 732.

Medicina Colonial. Madrid, 111, 451.

Medicinski Arhiv. Sarajevo, 923. Medicinski Glasnik. Belgrade, 924.

Medicina Española, 734.

Medicina, Cirurgia, Farmácia. Rio de Janeiro,

Medicine and Laboratory Progress. Cairo, 452.:

748.

INDEX OF PERIODICALS

Occasional Papers of Bernice P. Bishop Museum, Honolulu, 749. Offentliche Gesundheitsdienst (Der), 750. Österreichische Kleintierzüchter, 751. Österreichische Zeitschrift für Stomatologie, 929. Ohio Journal of Science, 252. Ohio State Medical Journal, 752. Okayama Igakkai Zasshi, 930. Orchardist of New Zealand, 253. Osaka Daigaku Igaku Zassi, 753. Oyo-Dobutsugaku-Zasshi. (Magazine of Applied Zoology.) Tokyo, 254.

Pakistan Journal of Science, 931. Pamphlet. Department of Health, Southern Rhodesia, 255.
Papers from the Michigan Academy of Science,

Arts and Letters, 754.

Parasitica. Gembloux, 123.

Parasitology, 29. Pastoral Review. Melbourne, 473, 755. Pediatria. Naples, 757.

Pediatrické Listy, 932.

Pediatrics. Springfield, Ill., 124.

Pemberitaan Balai Besar Penjelidikan Pertanian,

Bogor, 474. Pflanzenarzt. Vienna, 475. Pflanzenschutz. Munich, 476. Pflanzenschutzberichte. Vienna, 477. Philippine Journal of Surgery, 758. Phytoma. Paris, 256, 759.

Phytopathology, 30, 125, 257, 478.
Plant Disease Reporter, 31, 126, 479, 760.
Plant Disease Reporter. Supplement, 127, 480.
Plant Pathology. London, 258.
Plant and Soil. The Hague, 128.

Policlinico (Sezione Pratica). Rome, 761. Polski Tygodnik Lekarski. Warsaw, 481, 933.

Portugal Médico, 762. Poultry Field. Preston, 259.

Poultry Science, 32, 129. Poumon. Paris, 763. Praktický Lékar. Prague, 934.

Praxis. Berne, 764. Prensa Médica Argentina, 130, 765.

Prensa Médica Mexicana, 766. Press Bulletin. Georgia Agricultural Experiment

Station, 767.

Presse Médicale, 260, 482, 768.

Proceedings of the Alumni Association of the King Edward VII College of Medicine,

Singapore, 131, 769.
Proceedings of the American Society for Horti-

cultural Science, 770.
Proceedings. American Society of Sugar Beet Technologists, General Meeting, 771.
Proceedings of the American Veterinary Medical

Association, 772.
Proceedings. Association of Southern Agricul-

tural Workers, 483. Proceedings of the California Fig Institute, 935. Proceedings of the Helminthological Society of

Washington, 132, 261.
Proceedings of the Indian Academy of Sciences.

Section B, 484.

Proceedings of the Iowa Academy of Science, 936. Proceedings of the Louisiana Academy of Science,

Proceedings of the Nebraska Academy of Science,

937.
Proceedings of the Pakistan Science Conference,

Proceedings of the Rio Grande Horticultural Institute, Weslaco, Texas, 133.

Proceedings of the Royal Academy of Sciences,

Amsterdam, 774.
Proceedings of the Royal Society of Medicine,

485.

Proceedings of the Society for Experimental Biology and Medicine, 134, 77

Proceedings. Soil Science Society of Florida, 776. Proceedings. United States Livestock Sanitary

Association, 486, 777.
Proceedings of the United States National Museum, 487.
Proceedings of the Zoological Society of London,

Profilassi, 778.

Progress Report. Texas Agricultural Experiment

Station, 779.
Progresso Medico. Naples, 780.
Progresso Veterinario. Turin, 781.

Protoplasma, 489.

Ptitsevodstvo, 938. Publicación. Escuela de Veterinaria. Universidad de Buenos Aires, 490.

Publicaciones del Instituto de Biología Aplicada.

Barcelona, 782. Publicaciones del Instituto de Biología Aplicada. Barcelona. Serie Zoologica, 939. Publicações Avulsas do Instituto Aggeu Magal-

hães. Recife, 783.

Publications of the Institute of Marine Science,
University of Texas, 491.

Quarterly Bulletin of Northwestern University Medical School. Chicago, 940. Quarterly Bulletin of the South Pacific Commission, 941.

Oueensland Agricultural Journal, 263.

Records of the Indian Museum, 784.

Recueil de Médecine Vétérinaire, 492. Rendiconti. Istituto Superiore di Sanità. Rome,

Report of the Commonwealth Scientific and Industrial Research Organization, Australia, 786.

Department of Agriculture, New Report. Zealand, 493.

Report of the Department of Agriculture and Stock, Queensland, 787.
Report. Department of Scientific and Industrial

Research, New Zealand, 264. Report. Department of Veterinary Services,

Swaziland, 788. Report. East African Medical Survey, 789.

Report of the East Malling Research Station,

Report. Filariasis Research Unit, East Africa, 791.

Report of the Minister for Agriculture. Dublin,

Reports on the Progress of Applied Chemistry,

Report of the Rothamsted Experimental Station,

793. Report of the Science Service. Department of

Agriculture, Canada, 794. Report of the Science Service. Division of Botany and Plant Pathology, Canada, 943. Research Bulletin of the Hokkaido National

Agricultural Experiment Station, 265. Research. Rhode Island Agricultural Experiment

Station, 795. Revista de Agricultura. São Paulo, 796.

Revista Argentina de Agronomía, 494. Revista de la Asociación Médica Argentina, 797. Revista da Associação Médica de Minas Gerais,

Revista Brasileira de Biologia, 495.

Revista Brasileira de Gastroenterologia, 799. Revista Brasileira de Malariologia e Doneças

Tropicais, 800. Revista Brasileira de Medicina, 801. Revista Ceres. Minas Gerais, 802.

Revista Chilena de Higene y Medicina Preventiva,

944. Revista Chilena de Pediatría, 266, 803. Revista Clínica Española, 267, 496, 804. Revista Clínica de São Paulo, 805.

Revista Colombiana de Pediatría y Puericultura,

945. Revista Cubana de Pediatria, 946. Revista Española de las Enfermedades del Aparato Digestivo y de la Nutrición, 268, 947.

Revista Española de Oto-Neuro-Oftalmología y Neurocirugia, 806.

Revista da Faculdade de Medicina Veterinária. São Paulo, 807.

Revista de la Facultad de Medicina. Bogotá, 497. Revista Ibérica de Parasitología, 33, 135, 269, 498, 808.

Revista do Instituto Adolfo Lutz. São Paulo,

Revista de Investigación Clínica. Mexico, 810. Revista Kuba de Medicina Tropical y Parasitología, 136, 270, 811.

Revista Médica de Chile, 499, 812. Revista Médica de Córdoba, 271.

Revista de Medicina Veterinaria. Buenos Aires, 500.

Revista de Medicina Veterinaria. Lisbon, 501. Revista de Medicina Veterinaria y Parasitología. Caracas, 502, 813.

Revista Médico-Quirúrgica de Oriente. Santiago de Cuba, 814.

Revista de Paludismo y Medicina Tropical.

Mexico, 272, 815.
Revista Paulista de Medicina, 503.
Revista de Sanidad y Asistencia Social, 816. Revista de Sanidad e Higiene Pública. Madrid,

504.

Revista do Serviço Especial de Saúde Pública. Rio de Janeiro, 948.

Revue de l'Agriculture. Brussels, 505.

Revue de Chirurgie, 137. Revue d'Élevage et de Médecine Vétérinaire des

Pays Tropicaux, 273. Revue Horticole Suisse, 506. Revue de Médecine Vétérinaire, 507.

Revue Médicale du Moyen-Orient, 817.

Revue Médicale de Nancy, 949. Revue Neurologique, 818.

Revue de Pathologie Comparée et d'Hygiène Générale, 756.

Revue Romande d'Agriculture, de Viticulture et d'Arboriculture, 508.

Revue Suisse de Zoologie, 819. Rhodesia Agricultural Journal, 509. Rhodesian Farmer, 138, 510. Rhodesian Tobacco Journal, 511. Rice Journal. New Orleans, 512. Riforma Medica, 274, 820. Rivista di Biologia Coloniale, 821.

Rivista Italiana d'Igiene, 513, 822. Rivista d'Ostetricia e Ginecologia Pratica, 950.

Rivista di Parassitologia, 34, 139, 275, 514. Rivista di Patologia e Clinica. Parma, 951.

Rivista di Zootecnia, 823. Rural Research in C.S.I.R.O. Melbourne, 824.

Sad i Ogorod. Moscow, 825.

Sang, 826.

Sapporo Medical Journal, 827.

Schweizer Archiv für Neurologie und Psychiatrie,

515. Schweizer Archiv für Tierheilkunde, 276, 516. Schweizerische Gartenbaublatt, 828.

Schweizerische Gärtnerzeitung, 517. Science. Lancaster, Pa, 35, 140, 518.

Scientific Agriculture, 141, 277.

Scottish Agriculture, 142. Scottish Naturalist, 519. Semaine des Hôpitaux de Paris, 829.

Semaine Médicale. [Supplement to Semaine des Hôpitaux de Paris.], 830.

Seminar International. Hoddesdon, Herts, 520. Shikoku Acta Medica, 831.

Sicilia Sanitaria, 952. Sitzungsberichte der Österreichischen Akademie der Wissenschaften. Abteilung I. Mathematisch-Naturwissenschaftliche Klasse, 832.

South African Journal of Clinical Science, 521. South African Medical Journal, 36, 143, 522, 833.

Southern Medical Journal, 278. Southern Seedsman, 144.

Sovetskaya Meditsina, 145, 834, 953.

Speculum, 835. Srpski Arhiv za Tselokupno Lekarstvo. Belgrade,

954. Stain Technology, 146. Strasbourg Médical, 836. Surgical Clinics of North America, 837.

Svensk Frötidning, 523. Svenska Läkartidningen, 955.

Systematic Zoology. Washington, D.C., 524.

INDEX OF PERIODICALS

Terapevticheski Arkhiv, 956. Terre Marocaine, 957.
Texas Reports on Biology and Medicine, 525. Texas State Journal of Medicine, 147. Thérapie. Paris, 526. Therapie der Gegenwart, 279, 838.
Tidsskrift for den Norske Laegeforening, 958.
Tidsskrift for Planteavl, 527.
Tierärztliche Umschau, 148, 528. Tijdschrift over Plantenziekten, 149, 280. Tórax. Montevideo, 959. Transactions of the American Microscopical

Society, 37, 150, 281, 529.
Transactions of the American Neurological

Association, 960.
Transactions of the Illinois State Academy of Science, 839.

Transactions of the North American Wildlife Conference, 840.

Transactions of the Royal Canadian Institute, 530. Transactions of the Royal Society of New Zealand, 151.

Transactions of the Royal Society of South Australia, 841.

Transactions of the Royal Society of Tropical Medicine and Hygiene, 38, 152, 282, 531. Transactions of the Suffolk Naturalists' Society,

842. Transactions of the Wisconsin Academy of Sciences, Arts and Letters, 843. Treatment Services Bulletin. Ottawa, 844. Tropical Medicine and Hygiene News, 283.

Trudi Gelmintologicheskoi Laboratorii. Akademii Nauk SSSR, 961.

Trudi Zoologicheskogo Instituta. Nauk SSSR, 962. Türk Ijiyen ve Tecrübî Biyoloji Dergisi, 845.

Türk Veteriner Hekimleri Derneği Dergisi, 963.

Ugeskrift for Laeger, 846. Umschau, 847. Universitetet i Bergen. Årbok, 532. University of California Publications in Zoology,

Växtskyddsnotiser, 533. Verhandlungen der Deutschen Zoologischen Gesellschaft. (Zoologischer Anzeiger, Supplementband.), 848. Věstník Československé Zoologické Společnosti, 964. Veterinaria. Madrid, 534. Veterinaria. Sarajevo, 153, 284, 535.

Veterinariya, 285, 536, 849. Veterinarski Arhiv, 537. Veterinarski Glasnik. Belgrade, 850. Veterinary Medicine, 40, 154, 286, 538. Veterinary Record, 41, 155, 287, 539.

Veteriner Fakültesi Yayınları. Ankara Üniversitesi, 288, 965. Videnskabelige Meddelelser fra Dansk Natur-

historisk Forening i København, 851.

Vie et Milieu. Paris, 540, 852. Virchow's Archiv für Pathologische Anatomie und Physiologie und für Klinische Medizin,

853. Virginia Journal of Science, 541, 854.

Vlaams Diergeneeskundig Tijdschrift, 156, 542,

Vlugschrift. Plantenziektenkundige Wageningen, 543.

WHO Newsletter. (World Health Organization.) Geneva, 290.

Weibulls Illustrerade Arsbok, 856. West African Medical Journal, 289, 544.

Western Fruit Grower, 966. Wiener Klinische Wochenschrift, 857. Wiener Tierärztliche Monatsschrift, 157, 545. Wissenschaftliche Zeitschrift der Martin-Luther-Universität Halle-Wittenberg, 967.

World Crops, 42. World's Poultry Science Journal, 858.

Year Book. Institute of Inspectors of Stock of New South Wales, 546. Yokohama Medical Bulletin, 291.

Zdravstveni Vestnik, 968. Zeitschrift für Ärztliche Fortbildung, 292, 547. Zeitschrift für Hygiene und Infektionskrankheiten, 293, 548.

Zeitschrift für Immunitätsforschung und Experimentelle Therapie, 549.

Zeitschrift für Morphologie und Ökologie der Tiere, 550. Zeitschrift für Parsitenkunde, 158, 551, 859.

Zeitschrift für Pflanzenernährung, Düngung, Bodenkunde, 860.

Zeitschrift für Tropenmedizin und Parasitologie, 294, 552, 861. Zeitschrift für Urologie, 862.

Zentralblatt für Allgemeine Pathologie und Pathologische Anatomie, 863.

Zentralblatt für Bakteriologie, Parasitenkunde, und Infektionskrankheiten Hygiene.

Abteilung I. Originale, 43, 295.
Zentralblatt für Chirurgie, 969.
Zoological Magazine. Tokyo, 864.
Zoologicheski Zhurnal, 296, 553, 865.
Zoologické a Entomologické Listy, 866.

Zoologische Garten (Der). Leipzig, 554. Zoologische Jahrbücher. Abteilung für Syste-matik, Ökologie und Geographie der Tiere,

297, 867. Zoologischer Anzeiger, 159, 555, 868. Zoologiska Bidrag från Uppsala, 869. Zucker, 870.

(The reference is to the serial number: numbers in bold type indicate abstracts.)

Acanthias vulgaris, Trilocularia acanthiae-vulgaris Alaska, see also U.S.A. -, helminths in rodents 428e. -, trematodes in canines 132c.

Alces alces, Spiculopteragia n.sp. in 1002. Acanthobothrium gracile n.sp. in Narke japonica 49a. Alectoris graeca, Dicrocoelium n.sp. in 1001. - grandiceps n.sp. in Dasybatus zugei 49a. -, Lyperosomum n.sp. in 1001. - latum n.sp. in Dasybatus akajei 49a. - micracantha n.sp. in Dasybatus akajei 492. Aleutian Islands, helminths in Rattus norvegicus - triacis n.sp. in Triacis scyllium 49a. Acanthocephala in animals in Australia 841a. , Trichinella spiralis 😢 9a. Alfalfa, see Lucerne. - birds 529d. -, chemistry of glands 230bx. Algeria, helminths in dog 63a. -, evolution & speciation 524a. Schistosoma haematobium 182a. - in mammals in North America 230bv. Allgéniella obliqua n.sp. 610d. Alopias vulpinus, Marsupiobothrium n.g., n.sp. in -, morphology & host-parasite relationships 87e. -, nomenclature 132a. 49a. -, Phyllobothrium n.sp. in 49a. - reviewed 104b. America, see also North America. -, systematic position 296c. Acanthocephalans in animals in Iceland 1009. -, onchocerciasis 243a. Amidostomum anseris in goose in Canada 77a, 204b. Acanthocheilonema, see also Dipetalonema, Filaria, - -- , carbon tetrachloride 528d. Microfilaria. perstans in Cameroons, transmission by Culicoides spp. 179a. - raillieti in Fulica americana 76a. Ammospiza maritima, Leucochloridium n.sp. in -, development in Culicoides austeni 334e. 101d. - in man in French West Africa 7b. Amnicola dupotetiana, intermediary for Ratzia — — —, hetrazan 321a. joyeuxi 12a, 58c. - streptocerca in man in Belgian Congo 38k. Amoeboid organism attacking Heterodera rosto-chiensis 120c, 249a. Acanthocolpidae revised 162, 502a. Acanthostrongylea n.tribe 1002. Amoebotaenia emended 33a. Amphicotyle quinquarii n.sp. in Quinquarius 49a. Acetodextrinae n.subf. 1000. Achromadora inermis n.sp. 384a. Amphimerus caudalitestis n.sp. in Chironectes - ruricola macroprocta n.subsp. 384a. panamensis 588c. guayaquilensis n.comb. for Opisthorchis guaya-— semiarmata n.sp. 384a. quilensis 588c. Acridotheres tristis, Diorchis n.sp. in 688a. Amphiuma, 2nd intermediary for Stomatrema guberleti 981. , Hymenolepis n.sp. in 688a. Actinolaimus frostae n.sp. 342a. Actitis hypoleucos, Corrigia n.sp. in 1001. Anchidiplogaster n.g. for Diplogaster dubius 961bi. Acuaria, see also Dispharynx, Synhimantus.
— (Echinuria) uncinata in duck in Britain 41a. Ancylostoma, see also Hookworm. - in man in France 607b. Acutogordius n.g. for Gordius feae, G. doriae & -- braziliense & A. ceylanicum morphologically distinct 64a. G. incertus 158a. Aëdes togoi transmitting Wuchereria malayi - caninum, copulation 230bm. 409a. -, development inhibited by Roussin's salts Aelurostrongylus, see also Lungworms. 513a. -, phototropism of larvae 291a. - in rat, experimental 20k. Aepyceros melampus, Gaigeria pachyscelis in - Sigmodon hispidus, experimental 20k. -, water balance regulation 385c. -, helminths in 434a. - var. longespiculata to A. longespiculatum Africa, see also South Africa, West Africa. 895a. -, nematodes in mammals 586x. - duodenale extract, effect on blood 635a. -, schistosomiasis 290a. - - larvae, galvanotaxis not shown 753d. -, Schistosoma mansoni intermediaries 596a. - in man in Angola 586e. —, Taenia saginata 152k. ---, tetrachlorethylene 423d. Agamascaris septentrionalis n.sp. in Pleuronectes -- in Turkey 694a. limanda 851a. - -, therapeutic use of 58e. Agrostis tenuis, Anguina agrostis in 126c. - longespiculatum for A. caninum var. longespi-

culata 895a.

other felines 64b.

in cat, morphology 895a.

- paraduodenale in Felis (Leptailurus) serval &

Alactaga saltator, Skrjabinocercina n.g., n.sp. in

Alaeuris yumanae n.sp. in Dipsosarus dorsalis

529b.

```
Animals, domestic, helminthiasis in 453c.
Ancylostomiasis, see also Creeping eruption,
                                                            -, -, helminths in 40c, 67a, 138c, 172a, 245b, 245d, 388b, 629a, 662a, 926a, 961bo, 1003.
    Hookworm disease.
  control 632a.
                                                            , -, hydatid in 82a, 275a, 577c, 628a.
  in leopard 642a.
                                                            -, --, hydatidosis in 153a.
    - man 340c, 465a, 482a, 580a, 611b, 657b,
                                                            -, --, liver-fluke in 904b, 920a.
     725a, 810a, 820d.
                                                            -, --, lungworms in 238a.
       - in Angola 586g.
                                                            -, --, oxyurids in 104a.
      — — Cape Verde Islands 586f.
                                                           -, -, parasites in 971, 996.
     -, carbon tetrachloride 761a.
                                                           -, -, phenothiazine 963c.
    - in Chile, control 633a.
                                                            _, _, _ photosensitization 299.
    ---, chloroform 598b.
                                                           -, laboratory, Ancylostoma caninum in 20k.
   -, -, Ascaris in 440a.
  — — in India 220a, 423a.
— — — Italy 820a.
                                                            -, -, Schistosoma mansoni in 783g.
                                                           —, —, Strongyloides in 247a.
—, —, Trichinella spiralis in 961c.
  — —, pathology 260a.
— — in Puerto Rico 350b.
                                                           -, zoo, helminths in 488e, 488f.
   -- - Somaliland 330a.
                                                           Anomotaenia emended 33a.
   ----, symptoms 340e.
                                                            - filovata n.sp. in Charadrius vociferus 132d.
     — —, treatment 589a.
— — in Trinidad 657a.
                                                           Anonchotaenia gaugi n.sp. in Turdoides somervillei
                                                                688a.
  - - U.S.A., control 320a.
                                                           Anser albifrons, Hymenolepis n.sp. in 20d.
Andrya arctica n.sp. in Dicrostonyx groenlandicus
                                                             - -, - creplini in 20d.
                                                           Antarctic, Cercaria n.sp. 230i.
Angola, Ancylostoma duodenale 586e.
                                                           -, cestodes in seals 72a.
-, ancylostomiasis 586g.
                                                           Anthelmintics, acridine 742a.
-, Schistosoma mansoni 586k.
                                                            -, - derivatives 765a.
   , schistosomiasis haematobia 586m.
 Anguilla spp., Stegodexamene anguillae in 428c.
                                                           -, ACTH 467c.
                                                           -, alkyl resorcinols 930a.
 -, Telogaster opisthorchis in 428c.
                                                           -, aluminium compounds 292a.
  - anguilla, Triaenophorus lucii in 332a.
                                                           -, amino acridine compounds 413a.
 - japonica, Nybelinia n.sp. in 49a.
                                                            -, 2-amino-5-nitrothiazole 10b.
 Anguillula, see also Turbatrix.

— silusiae, effect of acridine dyes on growth & reproduction 464a.
                                                           -, anthiomaline 624b.
                                                           —, antibiotics 98a, 98c, 333a.
                                                           -, - & sulfa drugs 52b, 230z.
 Anguillulina, see also Anguina, Ditylenchus,
                                                            -, antimony compounds 430h, 702a, 702e.
      Pratylenchus, Radopholus, Rotylenchus, Tylen-
                                                            -, antrypol 68a, 152m, 586p.
                                                            -, arsenamide 54a, 177b.
   - (Ditylenchus) dipsacoidea n.sp. 56a.
                                                            -, arsenical compounds 134a, 702f.
   - tritici in wheat in Ireland 700a.
                                                            -, N-(p-arsenoso-benzyl)-glycine amide hydro-
 Anguina, see also Anguillulina.
   - sp. causing galls in grasses 26b.
                                                                chloride 385d.
                                                            -, Arsobal 624a.
 - agrostis in Agrostis tenuis, life-history &
                                                            __, ascaridole 709a, 827c, 899b.
      control 126c.
                                                            -, atebrin 94a, 105b, 210b, 352e, 637a, 924a,
  - poophila n.sp. in Poa sp. 296b.
 - tritici with Corynebacterium tritici causing
                                                                933c.
                                                            -, Atrimon 210a, 718a.
 disease in wheat 218a.
Animals, Acanthocephala in 841a.
                                                            -, bilberries 470a.
                                                            -, Butynorate 129b.
 -, acanthocephalans in 1009.
                                                            -, carbon tetrachloride 285b, 528d, 534a, 536e,
 -, chenopodium oil 148d.
                                                                546b, 586y, 761a, 850f.
 -, Diphyllobothrium sp. in 230bq.
                                                             -, -- with turpentine oil 492b.
 -, Echinococcus in 613a.
                                                            -, caricide, see also Anthelmintics, hetrazan.
 —, helminthiasis in 546c.
—, helminths in 245a, 345a, 471b, 516a, 516b,
                                                            —, — 41c.
                                                            -, cashew nut-shell oil 690c.
      726b, 813c, 942a, 992, 961p.
                                                            -, chemicals tested 157c, 157d, 638a.
  -, hydatid in 656a, 823a.
                                                            -, chenopodium oil 148d, 528b.
 -, iso-amyl ester of mandelic acid 148d.
                                                            -, chloroform 598b.
   -, new nematodes in 572a.
                                                              -, chloromycetin 499a.
  —, parasites in 1007, 1010.

—, — & disease in 986, 987.

—, Trichinella spiralis in 99a, 230u, 995.
                                                              -, Contaverm, see also Anthelminucs, pheno-
                                                                 thiazine.
                                                            —, — 735a.
  —, trichinelliasis in 785a, 890a.
   -, domestic, cerebrospinal filariasis in 221c, 251a, 283a, 358a, 569a.
                                                             -, cortisone 53a.
                                                            -, crystal violet 617c.
                                                            -, detergents 230cm, 839a.
    -, --, cestodes in 865a, 927a.
                                                            -, diamino-diphenyl-suphone 690a.
    -, -, cysticerciasis in 138a.
                                                            -, dichlorphenarsine hydrochloride 702i.
```

-, diethylcarbamazine 80a.

, 1-diethylcarbamyl-4-methylpiperazine

citrate 196a.

Anthelmintics, sodium antimony tri-gluconate Anthelmintics, 1-diethylcarbamyl-4-methylpiperazine citrate 196a. , - fluoride, see also Anthelmintics, Sus-s -, 2:3-dihydro-3-ketobenzo-1:4-thiazine 223a. caridin. -, Diospyros mollis berries 230bp. -, — — 157b, 545a, 575a, 850a, 889a. -, diphenyl methane compounds 230cn. -, solustibosan, see also Anthelmintics, stibanose.e —, Egressin 200a, 232a. —, emetine 271a, 797a. -, stibanose, see also Anthelmintics, solusti-i -, - hydrochloride 230cu. bosan. —, — — & sulpha drugs 696i. —, filix mas 244a, 617a. —, fouadin 170a, 230cp. -, - 278a. -, Supatonin 409a. -, Suscaridin, see also Anthelmintics, sodium -, gammexane 687a. -, gentian violet 514f, 946e. fluoride. -, glucantime 566a. -, - 118a. -, technique for testing 887b. —, n-heterocyclic compounds tested 319a. -, Terit 148a, 528e. -, hetrazan, see also Anthelmintics, caricide, -, terramycin 467a. notezine. --, -- 2g, 69a, 152h, 155h, 176e, 235a, 307a, 308a, 321a, 353a, 358b, 379a, 379c, 399g, 405b, 407a, 430j, 450d, 465b, 531a, 531c, 544c, 674a, 711a, 731a. -, 6-tertiary-butyl-m-cresol 261c. -, tetrachlorethylene, see also Anthelmintics, Vermalin. -, - 423d. -, toluene 177a. -, hexachlorethane 773a. -, Trichostrongylin 458a. -, hexamethylenetetramine 720a. -, triphenylmethane dyes 547a. -, hexylresorcinol 136a, 185a, 270e, 270f, 294c, -, various 93a, 118e, 147a, 180a, 247a, 292a, 430g, 702j, 965a. 649b, 811a, 811c, 814a, 946b. -, - & tetrachlorethylene 270c. -, iodine 492a. -, -, tested 316a, 385e, 617b, 899a. -, Vermalin, see also Anthelmintics, tetra--, in vitro & in vivo testing 155f. - - testing 13c. chlorethylene. -, iso-amyl ester of mandelic acid 148d. - 718a. -, Vermella 735c. -, Karlsbad salt 857a. -, vermiplex 251c. -, Magenwurmtabletten (Marienfelde) 458a. , volatile oils 827a, 827b. —, magnesium sulphate 573c. Anthobothrium bifidum n.sp. in Dasybatus akajei -, miracil-D, see also Anthelmintics, nilodin. - 69d, 223b, 235b, 289d, 328b, 424c, 801b. -, MSb 152d, 334d, 334g. - pteroplateae n.sp. in Pteroplatea japonica 49a. -, Nematolyt 764a. - rajae n.sp. in Raja kenojei 49a. -, nilodin, see also Anthelmintics, miracil-D. Antirrhinum, Meloidogyne spp. in 478b. — majus, Meloidogyne spp. in 478a. Apatemon gracilis of Caballero & Vogelsang, 1949 -, - 586n, 586o, 586r, 624c, 625a. -, notezine, see also Anthelmintics, hetrazan. -, - 647b. to Strigea vaginata Brandes 651a. -, organic antimonials 424c. Aphanurus caesionis n.sp. in Caesio kuning 568a. -, penicillin 129a. Aphelenchoides & Corynebacterium causing disease —, pheniodol 36c, 612c. in strawberry 331d. -, phenosafranine compounds 357a. — sp. in chrysanthemum, control 127b. -, phenothiazine, see also Anthelmintics, Conta-- spp., control by E.605 forte 397a. - in ornamental plants, control 543a. besseyi, see also A. oryzae.
redescribed 261f. - 10a, 40a, 113a, 148c, 165c, 230q, 230ci, 230cj, 230ck, 230cv, 299, 312a, 314a, 340b, 428r, 458a, 473a, 528a, 528c, 534a, 573a, - in rice, control 601b. 917a, 938a, 9630 - blastophthorus n.sp. in scabious 60a. -, - derivatives 207a. - fragariae & A. ritzema-bosi compared 164 l. -, - & derivatives, effect on enzyme systems - in strawberry in Denmark 527a. 371a, 371c. - -, new host records 527a. -, -, mode of action 346a. – — in plants in Australia 347a. -, β-phenyl-β-carboxy-(3,5-diiodo-4-hydroxy - redescribed 261f. phenyl) ethane 36c. - oryzae, see also A. besseyi. -, phloroglucinol 617a. -, culture in vitro 30m. -, piperazine analogues 417a, 417b. - in rice 483d, 601a. -, piperazine diphenylacetate 526a. ----, control 30c, 257e, 341a. -, potassium antimony tartrate 38j. - ritzema-bosi & A. fragariae compared 1641. -, Protuvlasan 850b. - in chrysanthemum, control 96a, 360a, -, pumpkin seed extract 887a. 475b. -, reviewed 165f. --- in France 325a. —, santonin 102a, 223c, 223d, 570a, 570b, 899c. —, sodium antimony tartrate 38j, 85a, 382a, ----, hot-water treatment 394a.

586q, 801f.

----, parathion 506a, 517a.

---, unusual symptoms 258a.

Ascarids of animals in man 294b. Aphelenchoides ritzema-bosi in dahlia 57a. - in carnivores & rodents, epidemiology 1004. -, new host records 527a. - in cat, vermiplex 251c. - - redescribed 261f. -, effect of detergents on development 937a. - subtenuis redescribed 261f. -, life-history 961k. Aploparaksis sachalinensis n.sp. in Capella solitaria - in mouse, larval migration 98d. -, structure of head 967a. tandani n.sp. in Tringa hypoleucos 688a. Ascaris antigens 403b. Apogon lineatus, Bothriocephalus n.sp. in 49a. - & appendicitis in man 958a. Aponurus carangis n.sp. in Caranx sp. 568a. - in bile-duct, effect on gastric juice 686c. Aporina percnopteri n.sp. in Neophron percnop----, --- liver 686a. terus 688a. - blocking duodenal sound 969a. Appendicitis & Enterobius in man 112a. causing vomiting 686d.
extract, bactericidal activity 548a. Aprocta colaptidis n.sp. in Colaptes campestroides - in heart of man 752a. - ptiloscelidis n.sp. in Ptiloscelis resplendens - & intestinal occlusion in man 6852, 7102, 741b, 572a. - pyrrhurae n.sp. in Pyrrhura cruentata 495a. Apus pacificus, Brachylecithum n.sp. in 1001. - in laboratory animals, pathology 440a. - larvae in brain of man 515a. Arctic, Echinococcus 613a. Arctostaphylos, Sphaeronema n.g., n.sp. in 261b. Ardeola grayi, Dilepis n.sp. in 688a. - & linitis plastica in man 735d. — in liver of man 310a. — man 106b, 405a, 423b, 544d, 576d, 608a, Argentina, Diphyllobothrium latum 130a. —, Diplostomulum n.sp. 500a.

—, Hedruris n.sp. in 551b.

—, new nematodes in animals 572a.

—, Trichonema (Cylicocyclus) radiatum 490a.

—, Tylodelphys n.sp. 500a. 682b, 845b, 861a. — — —, ascaridole 827c. — — — causing liver abscesses 266a. ———, control 719b. ———, hetrazan 405b. Arhythmorhynchus macracanthus n.sp. in Umbrina — —, nervous & allergic symptoms 743a. roncador 150e. — — , — symptoms 757a.
— ova, permeability of shell 410c. Ariolimax columbianus, intermediary for Brachylaemus virginiana 20g. ----, survival in sewage 980. Arius sp., Monodhelmis n.sp. in 568a. -, technique for culturing 410a. — , Prosogonarium n.g., n.sp. in 568a.

Artemia salina, Hymenolepis californicus cysti-- on vegetables, control 442a. - in pig 900a. cercus in 436a. ———, eosinophilia 953a. ———, sodium fluoride **575a**, 889a. Ascariasis in animals, chenopodium oil 148d. ---, iso-amyl ester of mandelic acid 148d. __ ___, Suscaridin 118a. - man 190c, 406a, 438a, 834a, 953c, 956a, - devosi n.sp. in Martes spp. 132e. - lumbricoides, carbohydrate metabolism 79d. 969b. — —, control 906a. — —, diagnosis & immunology 918b. --, changes in cell content during oogenesis 410d. ____, general account 911a, 946a. - -, cytology 774a. ---, hetrazan 176e, 531a, 531c. — —, digestive enzymes 975. — —, effect of alkyl resorcinols 930a. — — , immunology 845a. — — , nervous symptoms 594a. --, -- hetrazan in vitro 307a. - pig 240a. — extract & eosinophilia in guinea-pig 231a. - - , sodium fluoride 157b. -- & intestinal occlusion 270b. Ascarid larvae in mouse, experimental 230ca. — —, in vitro culture 79a, 79b. - ova, chemicals tested against 181a. — in man 107a, 190h. Ascaridia in fowl, carbon tetrachloride 536e. _____, cashew nut-shell oil 690c. ____ in Japan 396a, 396b, 396c, 396d. — —, phenothiazine 938a. — galli, anthelmintics tested against 13c. - - of man & pig, dentigerous ridges 152f. -, effect of media on development 13e. ____, form of denticles 463b. --, -- mineral deficiency 230y. --, mode of locomotion 411a, 686b. ---, --- temperature on development 13e. -, - X-rays on development 150b. --, -- spread 437a. ---, nitrogen metabolism 961bs. — in fowl, carbon tetrachloride 534a. -- ova, distortion 699a. ____ in Egypt 20o. ____, life-history 281d. ———, effect of Duponol 80 230cw.
———, effect of light on development 213c. _ _ _ _, penicillin 129a. ———, effect of urine 753b, 753c. __ _ __, phenothiazine 534a. - - -, function of protein coat 87c. - -, survival in different soils 428f. - - -, - dust 753a. _____, resistance 150a. — —, physiology 864b, 864c. — — in pig, sodium fluoride 850a. - skrjabini n.sp. in Tetraogallus himalayensis - - from pig, volatile fatty acids 6212.

38k. -, filariasis 328f.

- -, Hirudinea 595a.

-, - bovis 59a.

Begonia, Pratylenchus sp. in 126g.

- —, — & adenolymphocele 925b.

--, - rodhaini 58h, 327e, 328h, 596b.

Belonolaimus gracilis on crops in U.S.A. 125a.

- --, schistosome intermediaries 328a.

---, Onchocerca volvulus 586u.

- -, schistosomiasis 328e, 650a.

Belgium, Ditylenchus dipsaci 123a.

-- peanut, control 257 1. - - & wilt in cotton 125d, 483c. Beta cicla, Heterodera schachtii in 943a.

Bibliography, Azim, M. A. 430a. - & obituary, Brumpt, E., 58a.

Biology, Thiara granifera 487a.

—, — — ova 753a.

Bionomics, Ascaris lumbricoides 428f.

-, Fasciola hepatica 542a, 593a.

- —, Tetrapetalonema berghei 152i.

-, Schistosoma mansoni 59c.

Belgian Congo, Acanthocheilonema streptocerca

Ascaris lumbricoides, technique for preparing extracts 845a. -, volatile oils tested against 827a. — var. suum, physiology 18a, 18b. Ascarophis for Capillospirura 962a. - arctica n.sp. in Zoarces viviparus 962a. - filiformis n.sp. in Gadus morrhua 962a. - morrhuae redescribed 962a. - sebastodis n.sp. in Sebastodes caurinus 491a. Ascarops strongylina in cattle 132 1. Asparagus officinalis, Meloidogyne incognita var. acrita in 30e. Aspiculuris tetraptera in mouse, anthelmintics tested against 316a, 385e. - —, antibiotics 98a, 98c. Asthenes modesta, Diplotriaena n.sp. in 572a. Ateles geoffroyi, Capillaria hepatica in 588e. Australia, Acanthocephala 841a. -, Aphelenchoides fragariae 347a. -, Dicrocoelium dendriticum 546a. -, Fasciola hepatica 171a. -, Gongylonema pulchrum 546d. -, haemonchiasis 787a. -, helminthology, research reviewed 786a. -, helminths in animals 345a. -, - - cattle 412a. - — domestic animals 67a. -, Heterodera marioni 263a, 312b. -, hydatid 348a. -, schistosome dermatitis 110b. Australorbis, Cercaria n.spp. in 454a. - & Tropicorbis, calcium content 783 1. — glabratus, Cercaria n.spp. in 454c. ——, effect of desiccation 783k. ----, growth rate & morphology 783d, 783f. -, morphology 783a. ---, saponins tested against 783j. --, Schistosoma mansoni cercariae in 454d. Austria, eelworms in potato 477a. -, Gordiacea 158a. , Onchocerca gutturosa 157a. phrys 841b. Avellariaea n.tribe 1002.

-, Australorbis glabratus 783k. -, Cyclocoelum microstomum 675f. -, Dictyocaulus 536h. -, - spp. 196c. -, - viviparus 196b, 855a. -, Ditylenchus sp. 257f. -, eelworms in plants 257r. —, — — soil 592a, 860a. -, free-living eelworms 867a. -, Haemonchus contortus 196b. -, helminths in sheep 155d. -, Heterodera rostochiensis 117b, 164b. -, - schachtii 870a. -, microfilariae 154d. -, nematodes in pheasant 843b. —, — sheep 97c. -, Plagitura salamandra ova 854a. Austrofilaria rhipidurae n.sp. in Rhipidura leuco--, Schistosoma mansoni 61b. -, Schistosomatium douthitti 55a. -, Simulium spp. 599a, 599b. Avioserpens bifidus n.sp. in Glaucionetta islandica —, — damnosum 289b. -, Stegodexamene anguillae 428c. - galliardi Chabaud & Campana, 1949 & -, strongyle larvae 536j. Petroviprocta vigissi Schachtachtinskaja, 1951 —, Telogaster opisthorchis 428c. —, Thermocyclops nigerianus 61a. identical 58 L Axonchium dudichi n.sp. 56a. -, trichostrongyle larvae 29a. Aythya marila nearctica, Hymenolepis n.sp. in -, trichostrongyles in sheep 539a. 150c. -, Trichuris ovis 196b. Azim, M. A., bibliography & obituary 430a. , Tropicorbis centimetralis 783k. Birds, Acanthocephala in 529d. —, Capillaria in 961bp. Balistes capistratus, Cucullanus n.sp. in 491a. -, Choanotaenia n.sp. in 961bt. Basilichthys, Diplostomulum n.sp. in 500a. -, Corrigia n.sp. in 961bt. -, Tylodelphys n.sp. in 500a. —, helminths in 166, 688a, 961q. -, Leucochloridium n.sp. in 961bt. -, microfilariae in 390a. -, nematodes in 669a, 961y. -, Porrocaecum in 961e. , Trichostrongylus n.sp. in 961bt. 362

Batrachoseps attenuatus, Thelandros n.sp. in 428b. Baylisia baylisi n.g., n.sp. in Lobodon carcinophagus Baylisiella n.g. for Bothriocephalus tectus 72a. Bdellonyssus bacoti, Litomosoides carinii in 983c.
—, technique for culturing 20v. Beet, Ditylenchus dipsaci in 684a. Beetles, Carabonema n.g., n.sp. in 555a. -, parasites in 983a.

Boehmeria utilis, Meloidogyne arenaria thamesi n.subsp. in 128a. Böhmiella wilsoni in Neotoma floridana 320d.

Bolbolaimus denticulatus n.sp, 610d.

Bolbosoma bobrovoi n.sp. in Callorhinus ursinus

Bothriocephalus, see also Diphyllobothrium. - abyssmus in Echinostoma tanneri 230br.

- apogonis n.sp. in Apogon lineatus 49a. - branchiostegi n.sp. in Branchiostegus 49a.

- brotulae n.sp. in Brotula multibarbata 49a. - claviceps in Gasterosteus in Britain 332a. - fluviatilis n.sp. in Hymenophysa curta 49a.

- lateolabracis n.sp. in Lateolabrax japonicus 49a.

- tectus to Baylisiella n.g. 72a.

Box salpa, Mesometra orbicularis in 610b. Brachygobius xanthozona, Brachyphallus n.sp. in

Brachylaemus, see also Brachylaima.

- brauni nom.nov. for Harmostomum caudale of Diesing 320e.

- erinacei in pig, general account 627a.

- virginiana intermediaries 323a.

- metacercariae in Ariolimax columbianus 20g. Brachylaima, see also Brachylaemus.

- condylura n.sp. in Condylura cristata 529c. - virginianum metacercaria, morphology 693a.

Brachylecithum attenuatum parinum n.subsp. in Parus palustris 1001.

 capilliformis n.sp. in Turdus dauma 1001.
 chivosca n.comb. for Olssoniella chivosca 1001. - colorosum n.comb. for Lyperosomum colorosum

- coturnixi n.sp. in Coturnix coturnix 1001.

- cuculi n.sp. in Cuculus canorus 1001. - donicum n.comb. for Lyperosomum donicum

1001. - eophonae n.comb. for Lyperosomum eophonae 1001.

- kirghisensis n.sp. in Montifringilla alpicola 1001.

- lobatum glandarii n.comb. for Lyperosomum lobatum glandarii 1001.

- strixi n.subsp. in Strix uralensis 1001.

- pici n.sp. in Picus canus 1001. - praetenuis n.sp. in Apus pacificus 1001.

- uigurica n.sp. in Eremophila penicillata 1001. Brachyphallus brachygobii n.sp. in Brachygobius xanthozona 859a.

Brama rayi, Gymnorhynchus gigas plerocercoids in 381a.

Branchellion lobata n.sp. on Raja sp. 749a. Branchiostegus japonicus, Bothriocephalus n.sp. in

Branta canadensis, Catatropis n.sp. in 428a.

- -, Paramonostomum n.sp. in 428a.

-, Renicola n.sp. in 230da. Brazil, cysticerciasis 626a, 807b.

-, filariasis 586v.

-, - bancrofti 800a, 800b, 800d. -, helminths in man 343a, 399c.

__, __ pigeon 807a. -, hookworm 322b.

-, Hymenolepis nana 680a.

-, Schistosoma mansoni 783h, 783i, 948a. -, - intermediaries 798b.

-, schistosomiasis mansoni 399d, 503b, 809b.

Brazil, Strongyloides 322b.

-, Syngamus laryngeus 783c. , Wuchereria bancrofti 34b, 783m.

Britain, Acuaria (Echinuria) uncinata 41a.

-, Bothriocephalus claviceps 332a. -, Dictyocaulus viviparus 287b. -, eelworms in plants 793a.

-, Erpobdellidae 488d. -, fascioliasis 165g.

-, free-living eelworms 100a.

-, helminths in fresh-water fish 332a.

-, - - sheep 142a.

-, Heterodera rostochiensis 44, 45.

-, - schachtii 15a, 46.

-, Hirudo medicinalis 462a, 519a.

-, "husk" 51a.

-, Moniezia expansa 165b. megalorchis -, Plagiorchis (Multiglandularis) nom.nov. 29c.

-, trichinelliasis 108b.

, Trichuris trichiura 217a. Brotula multibarbata, Bothriocephalus n.sp. in 49a. Brumpt, Emile, bibliography & obituary 58a.
Brumptaemilius sclerophorus n.g., n.sp. in Julus

sp. 58d. Bucephalidae, germ cell cycle 713b.

— reviewed 281a.

Bucephalopsis tenuis n.sp. in Platycephalus indicus

- tylosuris n.sp. in Tylosurus coromandelicus 432b.

Bucephalus retractilis n.sp. in Caranx sp. 568a. sphyraenae n.sp. in Sphyraena sp. 568a. Buffalo, Schistosoma spindale in 273a, 773c.

-, stephanofilariasis in 92c. , Thelazia n.sp. in 221b.

Bulinus abyssinicus, possible intermediary for Schistosoma haematobium in Somaliland 514e.

contortus in France 852b. - truncatus, cercariae in 38h. Burma, Hirudinea 414a.

Caesio kuning, Aphanurus n.sp. in 568a. Calicophoron ijimai in cattle, life-history 151a. Calidris minuta, Dicranotaenia n.sp. in 961x.

subminuta, Dicranotaenia n.sp. in 961x. Callorhinus ursinus, Bolbosoma n.sp. in 961z. Callorhynchicola branchialis n.g., n.sp. on Callorhynchus callorhynchus 443a.

Callorhynchus callorhynchus, Callorhynchicola n.g., n.sp. on 443a.

Calotes versicolor, Camallanides prashadi in 821a. , Conispiculum guindiensis in 120h.

Camallanata reviewed 961bb.

Camallanides prashadi in Rana cyanophlyctis & Calotes versicolor 821a.

Camallanus baylisi in Echis carinatus 821a. Camel, Fasciola hepatica in 659a.

Cameroons, Acanthocheilonema perstans 179a.

-, paragonimiasis 289e. Canada, see also North America.

—, Amidostomum anseris 77a, 204b. —, eelworms in plants 203a, 794a, 943a

-, helminths in fish 252a.

-, Heterodera 277a. -, hydatid 656a.

-, Trichinella 10d, 11a, 204a.

Canines, trematodes in 132c. Cantaloupe, Meloidogyne incognita var. acrita 30e. Cape Verde Islands, ancylostomiasis 586f. Capella gallinago, Praeorchitrema n.g., n.sp. in 1001. - solitaria, Aploparaksis n.sp. in 961x. Capillaria in birds in Russia 961bp. - mink, phenothiazine 917a. — — poultry 989. — — sable, phenothiazine 917a. — aerophila in man, treatment 145a. - cholidicola n.sp. in Sorex araneus 329a. - contorta trasmitted by Eisenia foetida 230w. - hepatica in Ateles geoffroyi in Mexico 588e. oesophagicola n.sp. in Sorex araneus 329a.
pirangae n.sp. in Piranga erythromelas 236a. - plica in dog 850c. - salvelini n.sp. in Salvelinus alpinus 962a. - urinicola n.sp. in Sorex araneus 329a. - vanelli (Rudolphi, 1819), synonymy 58k. - ventricola n.sp. in Sorex araneus 329a. Capillospirura to Ascarophis 962a. Capingentidae n.fam. 305. Caprimulgus indicus, Krusadia n.g., n.sp. in 912e. Capsala spp. redescribed 784a. - gouri n.sp. on Thynnus thunnina 784a. Carabonema hasei n.g., n.sp. in beetles 555a. Carabonematidae n.fam. 555a. Caranx sp., Aponurus n.sp. in 568a. ---- , Bucephalus n.sp. in 568a. -, Opisthomonorchis n.g., n.sp. in 568a. Caridinicola platei n.sp. in prawn in Ceylon 78b. Cardium edule, Cercaria n.sp. in 228a. Carnation, Heterodera marioni in 759a. Carnivores, ascarids in 1004. Carnoya dubia n.sp. in Rhinocricus sp. 58d. Carp, helminths in 675g.
Carpiodes carpio, Neoechinorhynchus n.sp. in 20f. Caryophyllidea n.ordo 305. Cat, Aelurostrongylus abstrusus in 888a, 974. -, Ancylostoma longespiculatum in 895a. -, Dirofilaria immitis in 251d. -, Gnathostoma spinigerum in 67c. -, helminths in 251b, 251e, 428n, 501a. —, Opisthorchis felineus in 963b. —, vermiplex 251c. Catalaimus setifer n.sp. 610d. Catatropis harwoodi n.sp. in Branta 428a. Catenotaeniidae n.fam. 305. Cattle, Ascarops strongylina in 132 1. —, Calicophoron ijimai in 151a. —, cysticerciasis in 626a, 807b, 894a. -, Cysticercus bovis in 165a, 197a, 234a, 285a, -, Dictyocaulus in 536a, 536f. -, - viviparus in 156a, 287b, 813a. -, Eurytrema coelomaticum in 802a. —, Fasciola in 618c. __, _ in lung 618b. __, _ hepatica in 428m, 772a, 896a. —, fascioliasis in 6442, 849a. —, — & Bacterium coli infection in 988. -, - tuberculin reactions in 536k. - - tuberculosis in 985. -, Gongylonema pulchrum in 546d. -, helminths in 65a, 93a, 148b, 154a, 154b,

189a, 412a, 535b, 538a, 698a, 777c.

Cattle, "husk" in 51a. -, hydatid in 348a, 618a. —, lungworms in 902a, 904a, 920b, 1008. —, nematodes in 723a, 813b. -, - & gastro-enteritis in 8242. —, Onchocerca gutturosa in 157a. —, Ornithobilharzia n.sp. in 29f. -, Ostertagia ostertagi in 187b. -, Paramphistomum cervi in 245e. -, parasites in 702c. -, phenothiazine 230ci, 230ci, 230cv, 428r. - keratitis in 165c. -, Protuvlasan 850b. -, Schistosoma bovis in 38i. -, schistosomiasis bovis in 59a. -, Stephanurus dentatus in 132 1. -, Strongyloides in 666a. -, Thelazia in 536d. -, - rhodesii in 849d. - skrjabini in 213b. Celebes, new trematodes in fish 568a. Celeriac, Ditylenchus dipsaci in 681a. Celery, Dolichodorus heterocephalus in 30q. —, Hemicycliophora n.sp. on 261a. -, Paratylenchus hamatus in 30g, 478c. Centrorhynchus batrachus n.sp. in Rana tigrina 784d. spinosus, intermediaries listed 676a. Cephalobus (Tricephalobus) similigaster n.sp. 56a. Gephaloscyllium umbratile, Monorygma n.sp. in Ceramonema pisanum n.sp. 610d. - pselionemoides n.sp. 610d. Cercaria, new, in Helix pomatia 610a. - from Pyrene mercatoria 230cd. - n.sp. from Musculium sp. 230ce. — — — Pisidium sp. 230cf.
— acaudata n.sp. in Australorbis glabratus 454c. - C in fish, pathogenicity 120g. - chandleri n.sp. in Helisoma corpulentum 281c. - columbellae redescribed 327a, 368a. - franci n.sp. in rediae of C. tregouboffi 327a. 368a. - fulbrighti n.sp. in Cardium edule 228a. - goodmani n.sp. in Limnaea palustris 20bb. - metacercariae in Limnephilus 230dh, 230di. - hartmanae n.sp. in Lanicides vayssierei in Antarctic 230i. - hemiura n.sp. in Australorbis 454a. -littorinalinae in Littorina pintado in Hawaii - macrogranulosa n.sp. in Australorbis glabratus 454c. - mathiasi n.sp. 819b. — meyersi n.sp. in Physopsis africana 328g. — minense n.sp. in Australorbis glabratus 454c. — myae n.sp. in Mya arenaria 20bc. — nipponensis, morphology 410b. — naticae n.sp. in Natica clausa 675i. - nuda n.sp. in Bithynia striatula 696k. okabei redescribed 410e. - santense n.sp. in Australorbis 454a. — sensa n.sp. in Oncomelania hupensis 696k. - sturniae, morphology 696a. - takahashii redescribed 410e.

- tregouboffi n.sp. in Columbella rustica 3272,

368a.

Chile, helminths in man & dog 190a.

Cercaria tregouboffi parasitized by C. franci n.sp. 327a, 368a. - variglandis causing dermatitis in man 101f. - yoshidae, morphology & distribution 696j. Cercariae in Bulinus truncatus in Sardinia 38h. - from Cerithium 230cb. -, development of excretory system 37b. from marine gastropods 230cc. -, morphology 610a. , new, causing dermatitis 646a. Cercaricidal efficacy of ozone 20t. Cercopithecus, Filaria sp. in 69b. Cercorchis in Pseudemys scripta, life-history 262b. - for teaching helminthology 262a. Cereals, Ditylenchus dipsaci in 508a. Cericithidea californica, intermediary for Cloaci-trema michiganensis 230n. Cestoda (text-book) 305. Cestode, new, in muskrat 159b. Cestodes, amitosis 213a. -, development of vitelline glands 675d. - in dingo, new records 348a. — — dog 957a. - domestic animals 927a. _____, diagnosis 865a. _____ fish in Japan 49a. _ _ _ Mediterranean 852c. -, life-history, historical account 327c. - in man 968a. ———, diagnosis 766a. ———, effect on liver function 571a. _ — —, pheniodol 612c. -, phosphatases present 864d, 864e. - of Pinnipedia, systematics 422b. - in seals in Antarctic 72a. - - sheep 917d. - shrews in U.S.A. 21a. -, technique for mounting 241a. Cetacea, helminths in 961v, 961w. Cetorhinus maximus, Dinobothrium spp. in 540a. --, - n.sp. in 49a. Ceylon, Caridinicola n.sp. 78b. -, Monodiscus n.sp. 78c. , paramphistomes 78e. Chabertiinae n.subf. 1002. Chaenobryttus coronarius, Urocleidus n.sp. in 55b. Chaetodon awuga, Diplolasiotocus n.g., n.sp. in 568a. Chaeturichthys hexanemus, Pterobothrium n.sp. in Charadrius sp., Skrjabinoclava n.sp. in 961ba. - vociferus, Anomotaenia n.sp. in 132d. Chaunax sp., Contracaecum n.sp. in 491a. Cheiropteronematinae n.subf. 1002. Chelonia mydas, Haemoxenicon n.g., n.spp. in

Chemistry, Acanthocephala 230bx.

-, piperazine analogues 417a, 417b.

Chile, ancylostomiasis 633a. —, Dipylidium caninum 190b, 803a.

—, helminths in man 352b.

, santonin 102a, 223c, 223d, 899c.

—, 2:3-dihydro-3-ketobenzo-1:4-thiazine 223a. —, Haemopis sanguisuga 635c. —, hydatid membrane 398a.

-, ascaridole 899b.

-, miracil-D 223b.

-, Hymenolepis spp. 944a. —, — diminuta 190g. —, Trichostrongylidae 190d. Chimaericoloidea n.superf. 532a. Chimpanzee, Oesophagostomum stephanostomum China, eelworms in plants 480b. -, Fasciola hepatica 659a. -, filariasis bancrofti 861b. , helminths in man 659a. -, hookworm disease 552b. , Spirocerca lupi 327d. Chironectes panamensis, Amphimerus n.sp. in 588c. Chironomus thummi, 2nd intermediary for Plagiorchis maculosus 859b. Choanotaenia emended 33a. — spp. to Dictymetra 132d.

— burti nom.nov. for Paricterotaenia tringae 33a.

— cayennensis to C. joyeuxibaeri n.sp. 33a.

— hypoleucia n.sp. in Tringa hypoleucis 688a. - joyeuxibaeri n.sp. for C. cayennensis 33a. - sylvarum n.sp. in bird in Russia 961bt. Chordodes spp. in Japan 864a. — japonensis n.sp. in Tenodera spp. in Japan 603b. — lichyi n.sp. in Microcentrum 565a. Chorinemus moadetta, Prosorhynchus n.sp. in 568a. Chromadorita brevisetosa n.sp. 610d. Chrysanthemum, Aphelenchoides sp. in 127b. - ritzema-bosi in 96a, 258a, 325a, 360a, 394a, 475b, 506a, 517a.
—, eelworms in 661a, 828a, 907a.
Cionella lubrica, biology & life-history 374a.
—, intermediary for Dicrocoelium dendriticum in U.S.A. 20n. Circlus pallasi, Lyperosomum mosquensis n.subsp. in 1001. Citellus leucurus, Hymenolepis diminuta in 428g. Citrus, eelworms in 779a. -, Tylenchulus semi-penetrans in 9a, 30a, 30n, 133a, 494a, 664a. Cleidodiscus stentor on Pomoxis nigromaculatus, first record 713g. Clementeia trispiculata n.sp. in Leptodesmus 58d. tubulifera n.sp. in Leptodesmus jucundus 58d. Cloacitrema michiganensis, life-history 230n. Clonorchis sinensis, glyceromonophosphatases demonstrated 915a. Clover, Ditylenchus dipsaci in 164h, 973. -, eelworms in 523a, 856a. , red, Ditylenchus dipsaci in 921a. Clupea clupeoides, Prosogonotrema n.sp. in 568a. Cobbia dentata n.sp. 610d. Coelorhynchus sp., Microbothriorhynchus n.g., n.sp. in 49a. Coenuriasis in sheep, control 536i. Coenurus, see also Multiceps. - cerebralis in goat 963a. - glomerctus, morphology 339b. Coffee, eelworms in 191a. Colaptes campestroides, Aprocta n.sp. in 572a. Colinus virginianus, helminths in 840a. Colombia, helminths in man 922a. -, microfilariae in vertebrates 50b. Columbella rustica, Cercaria n.sp. in 3272, 3682. Condylura cristata, Brachylaima n.sp. in 529c. Conepatus suffocans, Filaria n.sp. in 572a.

Control, Meloidogyne hapla 125c, 483b. Conispiculum guindiensis in Galotes versicolor, survival in formalin 120h. -, - incognita 776h. -, onchocerciasis 450c. Conspicuum icteridorum, life-history 20bd. Contracaecum chaunaxi n.sp. in Chaunax sp. 491a. -, Pratylenchus sp. 75b. -, schistosome dermatitis 38 L - cyclopteri n.sp. in Cyclopterus lumpus 851a. -, - intermediaries 59b, 463a. - histiophori, male described 491a. -, schistosomiasis 229b, 328i, 650a. - lari n.sp. in Larus marinus 851a. -, Simulium neavei 38a. - longicaecum n.sp. in Platystoma sp. 572a. melichthysi n.sp. in Melichthys buniva 491a.
ogcocephali n.sp. in Ogcocephalus radiatus 491a. -, strongyles 198b. -, Strongyloides 536g. -, taeniasis 788a. - (Simplexonema) n.subg. 851a. -, Thelazia 536d. - (Synthetonema) n.subg. 851a. —, — rhodesii 849d. —, Triaenophorus spp. 643a. Cooperiinae n.subf. 1002. Control, ancylostomiasis 320a, 632a, 633a. -, Anguina agrostis 126c. -, Aphelenchoides spp. 397a, 543a. Copper oleate controlling Schistosoma japonicum i -, - besseyi 601b. -, - oryzae 30c, 257e, 341a. cercariae 13d. -, - ritzema-bosi 96a, 360a, 475b. Coregonus lavaretus, Proteocephalus exiguus in i , ascariasis 906a. 380b. -, Ascaris 719b. - pollan, Proteocephalus n.sp. in 408a. Corrigia bonasia n.sp. in birds in Russia 961bt. --, -- ova 442a. —, Belonolaimus gracilis 257 1. —, cerebrospinal filariasis 702b. - separatiorchis n.sp. in Actitis hypoleucos 1001. - viktori n.sp. in Coturnix coturnix 961q. -, coenuriasis 536i. Corynorhinus rafinesquii, Longibucca lasiura in -, Culex fatigans 800c. -, Cysticercus cellulosae 453a. Cosmocephalus australiensis n.sp. in Hydromys: -, Dictyocaulus 536a. chrysogaster 841b. —, — filaria 849c. Cottocomephoronema hamulatum n.comb. for ' —, Dipetalonema evansi 536b. Haplonema hamulatum 9610. -, Ditylenchus dipsaci 164j, 236b. Cotton, Belonolaimus gracilis & wilt in 125d, 483c. —, eelworm diseases 387b. , eelworms in 885a. -, eelworms in citrus 779a. Cotton-rat, see Sigmodon hispidus. -, - - clover 523a, 856a. Cottus cognatus, Triaenophorus nodulosus pleto--, — — mushroom 30b. -, — — plants 117a, 127a, 174a, 256a, 511a, cercoids in 370a. Cotugnia dayali n.sp. in Psittacula eupatria 688a. 677a, 678b, 992. —, — — & soil 26a. Coturnix coturnix, Brachylecithum n.sp. in 1001. - -, Corrigia n.sp. in 961q. -, -- potato 301. Crabs, Simulium spp. larvae & pupae on 120t. Creeping eruption in man 367a. -, - rice 512a. -, - rye 30d. ---, gammexane 687a. __, _ _ rye-grass 30d. __, _ _ soil 776c, 776d, 776g, 860a. ---, stibanose 278a. ---, treatment 584a. —, — — strawberry 825a. —, — — tobacco 510a, 776i. -- & Strongyloides in man 19a. -, visceral, in man 544a. -, fascioliasis 644a, 849a. Crenosoma, see also Lungworms. -, filariasis bancrofti 800b. coloradoensis n.sp. in Martes caurina origenes -, helminthiasis in domestic animals 453c. 101c. -, - - pig 453b. Criconema (Criconemoides) annulifer forma hygro--, helminths in animals 726b, 992. philum n.f. 56a. -, - cattle 154a, 538a, 698a. Criconemoides curvatum n.sp. 261d. —, — — domestic animals 138c, 961bo, 1003. — cylindricum n.sp. 261d. -, - - horse 755a. - informe redescribed 261d. -, — — pig 14a. -, — — poultry 14a. - lobatum n.sp. 261d.
- mutabile redescribed 261d. — parvum n.sp. 261d. — teres n.sp. 261d. -, - - sheep 83b, 155e, 165f. -, -- - & goat 878. -, Heterodera avenae 476b. - xenoplax n.sp. 261d. -, - marioni 383a. Crops, Meloidogyne spp. in 30k. -, - - & H. rostochiensis 205b. Cruciferae, Heterodera cruciferae in 760a. —, — rostochiensis 5a, 44, 45, 60c, 117d, 125b, 164c, 164d, 257c, 257t, 373a.
—, — schachtii 15a, 46, 678d. Cryptogonimea n.tribe 1000. Cucullaninea n.tribe 961bb. Cucullanus hansoni n.sp. in Balistes capistratus -, hydatidosis 598a, 652a, 781a, 850d. 491a. -, lungworms in pig 239a. longipapillatus n.sp. in Melichthys buniva -, Meloidogyne 354a, 509a, 747a, 771a, 910b, Cuculus canorus, Brachylecithum n.sp. in 1001. 913a. -, - spp. 31b. Cucumber, Meloidogyne in 257b, 479a.

Culex fatigans, control 800c. - - transmitting Wuchereria bancrofti 800f. Culicoides austeni, intermediary for Acanthocheilonema perstans 334e. - nubeculosus transmitting Onchocerca spp. 50a. - pifanoi, Microfilaria in 816b. Cyathocephalus truncatus in Gammarus in Yugoslavia 535a. Cyathocotylidae reviewed 230f. Cybium guttatum, Pricea n.spp. on 784c. Cyclocoelum microstomum, larval migration 675f. - mutabile redescribed 269c. Cyclops n.spp. in Nigeria 488c. Cyclopterus lumpus, Contracaecum n.sp. in 851a. Cylindropharyngea n.tribe 1002. Cylindrophorus suppressed 651d. - typicus to Platybothrium baeri nom.nov. 651d. Cynoscion spp., trematode in 230db. Cyprus, hydatid in domestic animals 82a. Cysticerci containing ova 853a. Cysticerciasis in cattle 894a. ——— & pig in Brazil 626a, 807b. –, cerebral, in man 108a, 193a, 404a, 673a, 891a, 928a, 960a, 960c. - in domestic animals 138a. - -- man 830b, 933a. —, ocular, in man 923a. — in pig in Mexico 632b. - - sheep 352f. Cysticercoids, new, in Gammarus 120a. Cysticercus, see also Taenia. - bovis in cattle in Denmark 165a. - — — France 197a. - -- -- Germany **234a.** — — — Russia 285a. — — — Yugoslavia 850e. - --- man 340f. - cellulosae in dog 813d. - - killed by sodium chloride 453a. — in man 396e, 719a. — — — in Gold Coast 61c. --- -- India 219a. - fasciolaris, development of hooks 389a. - in rat, splenic lesions 635b. - madoquae, life-history 618e. - tenuicollis in sheep, diagnosis 453d. Cystocaulus & pneumonia in sheep 961bh. — nigrescens in sheep in Italy 618d. Cytology, Ascaris lumbricoides 410d, 774a. —, Bucephalidae 713b. -, cestodes 213a, 675d. -, Fasciola hepatica 658a. -, Mesocestoides variabilis 428p. -, Parascaris equorum 47a, 71a, 489a. -, Raillietina urogalli 675a. -, Rhabditis belari 337a. -, - elegans 12b, 62a, 368b. -, Spirorchis magnitestis 713a. , Strongyloides ratti 359a. Czechoslovakia, enterobiasis 43b. -, helminths in man 201a, 932a. -, Skrjabinema ovis 964a.

—, helminths in man 201a, 932a.

—, Skrjabinema ovis 964a.

—, trichinelliasis 934a.

Dactylella acrochaeta n.sp. attacking eelworms

Dagda asymmetrica n.sp. 610d. Daffodil, eelworms in 365a. Dahlia, Aphelenchoides ritzema-bosi in 57a. Dasybatus akajei, Acanthobothrium n.spp. in 49a. -, Anthobothrium n.sp. in 49a. - zugei, Acanthobothrium n.sp. in 49a. Dasymetra natricis n.comb. for D. nicolli 981. nicolli to D. natricis n.comb. 981. Deer, helminths in 276a. Delamurella hyperoodoni n.g., n.sp. in Hyperoodon ampollatus 1002. Demaniellidae n.fam. 961bi. Delphinapterus leucas, Leucasiella n.g., n.sp. in Denmark, Aphelenchoides fragariae 527a. -, - ritzema-bosi 527a. -, Cysticercus bovis 165a. -, Ditylenchus dipsaci 527a, 681a. -, nematodes in birds 669a. -, Taenia saginata 165a. Deropristiinae emended 29b. Desmoscolex falklandiae n.sp. 868a. - latus n.sp. 868a. Diacetabulum curvicolon n.g., n.sp. in Tringa incana 1000. Diagnosis, ascariasis 918b. -, cestodes in domestic animals 865a. —, — — man 766a. -, Eurytrema coelomaticum 802a. -, Fasciola hepatica 426a. —, filariasis 877. —, — bancrofti **399i.** —, helminthiasis 578a. —, — in cattle 148b. -, helminths in domestic animals 662a. -, - ruminants 866a. -, hydatid 352a, 451b, 748a. -, paragonimiasis 639a. -, Schistosoma japonicum 552c. -, schistosomiasis 58i. -, - haematobia 625a. -, - mansoni 355a, 585a, 799a, 799b, 799c, 799d. -, trichinelliasis 549a, 653a. Diaptomus franciscanus, intermediary for Proteocephalus primaverus 428t. Dichelininea n.tribe 961bb. Dichromadora abnormis n.sp. 610d. Dicranotaenia guschanskoi n.sp. in Calidris minuta – skrjabinissima n.sp. in Calidris subminuta 961x. Dicrocoelium, see also Liver-fluke. -- dendriticum in Cionella lubrica, development -, formation of slimeballs 81c. - —, glyceromonophosphatases demonstrated 915a. - -, infection of definitive hosts 81d. — —, life-history 230ch, 374c. — — in man, intracranial invasion 363a. — — sheep in Australia 546a. ———— & goat in Germany, 23a. ——— U.S.A. 777b. _____, life-history 20n. - lanceatum sciuri n.var. in squirrel 1001. - moschiferi n.sp. in Moschus moschiferus 1001. - petrovi n.sp. in Alectoris graeca 1001.

Diplogaster coprophagus of Goodey, 1929 too Paroigolaimella anchicoprophaga nom.nov. Dicrostonyx groenlandicus, Andrya n.sp. in 428e. -, Taemia crassiceps in 428e. Dictymetra for Choanotaenia 132d. 961bi. — numenii n.g., n.sp. in Numenius 132d. — paranumenii n.sp. in Numenius 132d. Dictyocaulus, see also Lungworms. de Man, 1876 to Paroigolaimella n.g. 961bi. - dubius to Anchidiplogaster n.g. 961bi. - lheritieri to Pristionchus n.comb. 961bi. - in cattle, control 536a. - paesleri n.sp. 192a. - similis to Paradiplogaster n.g. 961bi. - striatulus to Pareudiplogaster n.g. 961bi. - horse, bionomics 536h. - subdentatus n.sp. 192a. - sheep in Russia 917b. - spp., survival of free-living larvae 196c. Diplogasterata n.subordo 961bi. — filaria in sheep, experimental 319b.
————, inhibition of development 120b.
————, treatment & control 849c. Diplogasteritus n.g. for Diplogaster spp. 961bi. Diplogasterium n.g. for Diplogaster spp. 961bi. Diplogasteroides africanus to Goffartia africanad n.comb. 867a. - viviparus in cattle in Britain 287b. ————, epidemiology 156a. ————, Protuvlasan 850b. variabilis to Goffartia variabilis n.comb. 867a. Diplogasteroididae n.fam. 961bi. Diplogastrellus n.g. for Diplogaster spp. 961bi. - - -, treatment 813a. - -, survival of free-living larvae 196b. Diplolasiotocus chaetodontis n.g., n.sp. in Chaetodonn awuga 568a. -, - on pasture 855a. Dihemistephanus sturionis to Pristicola n.g. 29b. Diplostomulum mordax n.sp. in Basilichthys in Dilepis ardeolae n.sp. in Ardeola grayi 688a. Argentina 500a. "Diplostomum volvens" in stickleback, life-history Dingo, cestodes in 348a. 120g. -, Echinococcus granulosus in 348a. Dinobothrium spp. in Cetorhinus maximus 540a. Diplotriaena in Paroaria, morphology 796a. — modesta n.sp. in Asthenes modesta 572a. — muscisaxicolae n.sp. in Muscisaxicola maculi-- humile n.sp. to D. spinosum Baylis, 1950 651d. - planum Linton, 1922 to Gastrolecithus n.g. rostris 572a. Dipodomys spp., helminths in 230br. - spinosum Baylis, 1950 for D. humile n.sp. 651d. - spinulosum n.sp. in Cetorhinus maximus 49a. Dipsacus fullonum, Ditylenchus dipsaci in 236b. Dioctophyme renale in dog in France 492c. Dipsosarus dorsalis, Alaeuris n.sp. in 529b. Dipylidiinae reviewed 33a. Diomedea chrysostoma, Diomedenema n.g., n.sp. Dipylidium caninum in man 815b. in 841b. Diomedenema diomedeae n.g., n.sp. in Diomedea —— in Chile 190b, 803a. chrysostoma 841b Dirofilaria, see also Filaria, Microfilaria. Diorchis reviewed 961f. - reviewed 530a. - (Acanthodiorchis) n.subg. 961f. - conjunctivae in man 2b. - tilori n.sp. in Acridotheres tristis 688a. - desportesi n.sp. in Euarctos americanus 372a. Dipetalonema, see also Acanthocheilonema, Filaria, - immitis in cat in U.S.A. 251d. - - dog 286a. Microfilaria. ---, arsenamide ineffective 177b. reviewed 58f. - blanci n.sp. in Meriones libycus 58f. ----, hetrazan 155h. — —, periodicity 40b, 287a.
— —, transplantation of adults 230m. - in Meriones libycus, life-history 79c. - evansi in Camelus spp., life-history & control - intermediaries 713f. Diphyllobothrium, see also Bothriocephalus, Spar-- - in mosquitoes, infection rates 230dk. ganum. - repens in dog, caricide 41c. - & anaemia in man 168a. - ursi, male & microfilaria described 530a. - anaemia, monographed 567a. Disculicepitidae n.fam. 305. — in man, haematology 167a. — sp. from Salmo gairdnerii, in animals 230bq. Disculicepitidea n.ordo 305. Dispharynx, see also Acuaria.

— brevicordon n.sp. in Falco sp. 572a. - latum in dog, experimental 2f. — — man, general account 121a. - ketupae n.sp. in Ketupa zeylonensis 912d. —, morphology 586w. — plerocercoids in fish 553d. Distormon sp. Linton, 1901 to Tormopsolus lintoni n.sp. 162. Ditestolepis n.g. for Hymenolepis diaphana 329a. Ditylenchus, see also Anguillulina. - in Salmo irideus in Argentina 130a. — perfoliatum to Glandicephalus n.comb. 72a. Diplobulbus scari n.sp. in Scarus sp. 568a. - sp., bionomics 257f. - in lucerne 126b. Diplocotylidae n.fam. 305. Diplogaster spp. to Diplogasterium n.g. 961bi. - spp., cellulase & chitinase present 559. - Diplogasteritus n.g. 961bi. -, host range 60b. - Diplogastrellus n.g. 961bi. - destructor, effect of host on size 331c. - Eudiplogaster n.g. 961bi. -, host range 164k. — Fictor n.g. 961bi. — Oigolaimella n.g. 961bi. — Pristionchus 961bi. - — in potato, immunity 794a. — — Stachys palustris 60b. - dipsaci in beet 684a. - butschlii to Fuchsia butschlii n.comb. 961bl. - celeriac in Denmark 681a.

Earwig, Mermis nigrescens in \$42a. Ditylenchus dipsaci in cereals, popular account Echeneibothrium bifidum u.sp. in Trachurus trachurus 49a. - clover & lucerne, resistance 973. Echinococcus, see also Hydatid. _____, resistance 164h. — in animals in Arctic 613a.
— granulosus in dog in Italy 275a. - -, control 164j. - in Dipsacus fullonum & other plants 236b. - - dingo 348a. - - garlic 692a. — —, histology 765c. - -, host range 123a, 164i. -----, life-history, popular account 27a. - in mangold in Belgium 123a. - -, musculature of scolices 58b. - - oats in Sweden 533b. Echinocotyle glareolae n.sp. in Tringa glareola - - onion in Sweden 886a. - - red clover, resistance 921a. - hypoleuci n.sp. in Tringa hypoleucos 688a. -, revival after hot-water treatment 141b. - minutissima n.sp. in Querquedula circia 688a. - in rye, technique for testing susceptibility Echinoparyphium flexum metacercariae in frog, life-history 230dg. — — strawberry in Denmark 527a. - recurvatum in fowl 936a. Dog, arsenamide 177b. Echinostomatid metacercariae in fish 696e. -, Capillaria plica in 850c. Echiostoma tanneri, Bothriocephalus abyssmus in -, caricide 41c. -, cestodes in 957a. Echis carinatus, Camallanus baylisi in 821a. -, chenopodium oil 528b. Ecology, Limnaea truncatula 165g. -, Cysticercus cellulosae in 813d. , schistosome dermatitis 88a. -, Dioctophyme renale in 492c. Eelworm diseases, control regulations 387b. Eelworms attacked by Dactylella acrochaeta 460a. -, Diphyllobothrium latum in 2f. -, Dirofilaria immitis in 40b, 230m, 286a, 287a. - causing disease in tobacco 198a. -, Dracunculus insignis in 433a. - in chrysanthemum, Aralo 828a. -, Echinococcus granulosus in 275a. ———, parathion 661a. ———, popular account 907a. -, Eucoleus hepaticus in 135a. —, filariasis in 1872.
—, helminths in 28a, 28b, 63a, 95b, 95c, 118e, 135a, 190a, 251e, 428n.
—, hetrazan 155h, 407a, 711a. - citrus, control 779a. clover seed, control 523a, 856a.
coffee, "guamo" & plantain 191a.
cotton, popular account 885a. -, phenothiazine 230ck. - - crops in Sweden 533a. -, Spirocerca lupi in 154c, 395a. - — daffodil, Systox 365a. -, Taenia brauni in 819a. -, Tenuiproboscis n.sp. in 63b. - fig 935a. - Hibiscus sabdariffa, resistance 126f. -, Terit 528e. - iris bulbs, morphologically identical with —, 6-tertiary-butyl-m-cresol 261c.

Dolichodorus heterocephalus on celery 125a. Ditylenchus destructor 141a. - Lycopersicum & Physalis, resistance 483a. _ _ _ _, causing decline 30q. — — mushroom, control 30b. - in Eleocharis dulcia 30p. - narcissus, general account 364a. Domestic animals, see Animals, domestic. - oats, general account 712a. Dominican Republic, Schistosoma mansoni 176d, - pineapple in Mexico 387a. — — plants 966a. Dorylaimus beaumonti n.sp. 384a.

— carteri brevis n.subsp. 384a. — — —, bionomics 257r. — — in Canada 203a, 794a, 943a. - modestus n.sp. 384a. ———, causing root-knot 663a. ——— in China 480b. - nivalis n.sp. 384a. - paraconfusus n.sp. 384a. — — —, control 117a, 256a, 678b, 992. — — —, — regulations 677a. — — — by soil fumigants 127a, 174a. — — in France 57a. — pseudoagilis n.sp. 384a. - subacutus n.sp. 384a. Dracontiasis in man, notezine 647b. Dracunculus intermediaries in Nigeria 544b. — — —, general account 795a, 847a, 881a. — — in Holland, general account 446a. - insigmis in dog in U.S.A. 433a. — medinensis, calcified 224a. --- Iraq 216a. --, - in man 833a. _ — — — Ireland 313a, 313b, 942a. — —, popular account 392a, 767a.
— —, production of toxins 117c, 119b, 376a. _ _ in man 38e, 18oc. Duck, Acuaria (Echinuria) uncinata in 41a.

—, Hymenolepis n.sp. in 688a. ____, research in Britain 793a. _____, ___ progress 505a. _____, ___ reviewed 335a. -, Porrocaecum crassum in 213f, 961j. - Trichobilharzia n.sp. in 158b. __ ____, reviewed 961bi. - - & soil, control 26a. _____, soil fumigants 311a. ____ in South Africa 388c. Earthworm, intermediary for Porrocaecum crassum

----, control 511a.

213f.

Earthworms, anthelmintics tested on 827b.

Eelworms in plants (text-book) 558.	rinterobiasis in man, phenothiazine 340b, 573a.
— in plants in U.S.A. 126h.	— —, piperazine diphenylacetate 526a.
— on plants in U.S.A. 30f, 2570.	— —, technique for diagnosing 605a.
, reviewed 776b.	———, terramycin 467a.
— in potato 999.	, treatment 43b, 147a, 906c, 955a.
— — in Austria 477a.	& poliomyelitis in man 269a.
— — —, control 301.	-, technique for diagnosing 36b, 846a.
— — —, general account 169a.	Enterobius & appendicitis in man 112a, 609a,
& predatory fungi in soil 114a.	863a.
- & "rayadilla" in Musa spp. 306a.	— in man 672e.
— in rice, control 512a.	———, Atrimon 210a, 718a.
— — , resistant varieties 721a.	— — in Italy 514d, 822a.
—— rye 722a.	——————————————————————————————————————
— — & rye-grass, control 30d.	— —, technique for detecting 552a.
— — soil, bionomics 592a.	———, treatment 292a.
, control by soil fumigants 776c, 776d,	— — —, Vermalin 718a.
776g.	—, oviposition 456a.
———, effect of environment & control 860a.	- & peritonitis in man 683a.
———, reviewed 776a.	- vermicularis in man 293a, 578c, 708a, 765b.
- strawberry, control 825a.	———— in Italy 34c, 785e.
sweet-potato, D-D 30h.	, triphenylmethane dyes 547a.
-, technique for freeing from bacteria 790a.	in U.S.A. 20bf.
— in tobacco, control by crop rotation 776i.	Entobdella brattströmi n.sp. on Paralichthys
	adspersus 443a.
— — , —, popular account 510a.	Eopsetta jordani, helminths in 20m.
— —, ethylene dibromide 30i.	
— —, resistance 86a.	Epidemiology, ascarids 1004.
———, soil fumigants 202a.	—, Dictyocaulus viviparus 156a.
——— in Switzerland 508c.	Equines, Parascaris in 965a.
-, free-living, bionomics 867a.	Eremophila penicillata, Brachylecithum n.sp. in
-, -, in compost 192a.	1001,
-, -, contaminating blood films 84a.	Erpobdellidae in Britain, revised 488d.
—, —, from Italy, n.spp. 610d.	Euarctos americanus, Dirofilaria n.sp. in 372a.
-, -, - Kenya 342a.	Eubothrioides lamellatus n.g., n.sp. in Zenopsis
—, —, on marine algae 100a.	nebulosa 49a.
-, -, in marine mud in Sweden 869a.	Eucoleus hepaticus in dog in Spain 135a.
-, -, osmotic adaptation 550a.	Eudiplogaster n.g. for Diplogaster spp. 961bi.
-, -, phylogeny 848b.	Euparadistomum to Platynotrema 688b.
,, reviewed 961bi.	Euryhelminae n.subf. 1000.
—, —, from Rumania 56a.	Euryhelmis pacificus n.sp. in Mustela vison 428i.
-, -, - soil in Switzerland 384a.	Eurystomus orientalis, Pancreatrema n.g., n.sp. in
-, -, - Spain 939a.	1001.
,, new records 782a, 782b.	Eurytrema antigen, technique for preparing 802a.
Egg counts, technique for statistical examination	- spp., glyceromonophosphatases demonstrated
422a.	915a.
Egypt, Ascaridia galli 200.	— coelomaticum in cattle, diagnosis 802a.
—, helminths in poultry 20be.	— pancreaticum in ruminants in Mauritius 234b.
-, Schistosoma mansoni 452a.	Evolution, Acanthocephala 524a.
Eisenia foetida transmitting Capillaria contorta	
230w.	
Eleocharis dulcia, Dolichodorus heterocephalus in	Falco sp., Dispharynx n.sp. in 572a.
30p.	- subbuteo, Skrjabinus n.sp. in 1001.
Embryology, cercariae 37b.	Farancia, Stomatrema guberleti in 981.
-, Nematoda 848a.	Fasciola, see also Liver-fluke.
Enchodelus rhaeticus n.sp. 384a.	— in cattle 618c.
— vestibulifer n.sp. 384a.	lung of cattle, lymph node changes 618b.
Endocotyle incana n.g., n.sp. in Tringa incana	—, protein metabolism 230co.
1000.	- sp. in sheep, longevity 261g.
Enoploides brunettii n.sp. 610d.	- gigantica, life-history 912a.
Enoplolaimus enoploidiformis n.sp. 610d.	- hepatica, anthelmintics tested against 702j.
— subterraneus n.sp. 610d.	— in camel & pig in China 659a.
- villosus n.sp. 610d.	——— cattle in Hawaii 428m.
Enterobiasis în Czechoslovakia, incidence 43b.	Puerto Rico 896a.
— — man 24a.	
———, bilberries 470a.	, chemicals tested against 157c, 157d.
, Egressin 200a, 232a.	, cytology 658a.
— —, hexylresorcinol enemas 811c, 814a.	- effect of 2:3-dihydro-3-ketobenzo-1:4-
——— in Italy 275b.	thiazine in vitro 195b.
	2000.

Fasciola hepatica, effect of low temperatures on	Filariasis in dog 187a.
eggs 177c.	, treatment 711a.
, phenothiazine oxidation products	- & elephantiasis in man in Tanganyika 789a.
in vitro 195a.	— —, report of conference 830a.
, formation of egg-shell 26c.	— — eosinophilia in man 949a.
——, life-history 679a.	— in man 54b, 358c.
— in man 441a, 497c, 812a.	— —, Arsobal 624a.
——————————————————————————————————————	— — in Belgian Congo 3281. — — Brazil 586v.
————, emetine 271a.	———, diagnosis, distribution & treatment 877.
— —, metabolism 702h.	— — , diamino-diphenyl-sulphone 690a.
— — ova, chemicals tested against 20h.	——— in Fiji, corrigenda 531d.
— — —, technique for detecting 17a. — — in pig 245c.	————, general account 152a.
——— rabbit, antimony compounds 702a.	———— French West Africa 362b.
in Belgium 593a.	———, general account 180d.
Rattus rattus, new host record 20s.	, hetrazan 235a, 358b, 379a, 430j.
——— ruminants in Belgium 542a.	———, — & lymphadenitis 674a.
———— sheep in Australia 171a.	— — in Indo-China, survey 58g.
———, carbon tetrachloride 285b.	————— Lebanon 817a.
— —, transaminase activity 385a.	———— Maldive Islands 654a.
——, treatment 702g.	— —, nervous symptoms 328c.
, vitellaria atrophied 33c.	— — in Tanganyika 791a.
Fascioliasis & Bacterium coli infection in cattle	——————————————————————————————————————
988.	— ————————————————————————————————————
— in Britain, epidemiology 165g.	399i.
— — cattle, control 644a, 849a.	— — & lymphangiectasis in man 768c.
— — man, emetine 797a.	— in man 7a.
— & tubercular reactions in cattle 536k.	in Brazil 800a.
— — tuberculosis in cattle 985. — hepatica in man, carbon tetrachloride 586y.	, control 800b.
— —, popular account 109a.	, survey 800d.
Felis (Leptailurus) serval, Ancylostoma para-	China 861b.
duodenale in 64b.	, hetrazan 308a.
Fellodistomatoidea n.superf. 339a.	———— in New Caledonia 941a.
Fellodistomum fellis, life-history 553b.	, survey 728c.
- preovaricum n.sp. in Galeichthys seemanni	———, pathology 445a.
588b.	— — — in Venezuela 816a.
Fictor n.g. for Diplogaster spp. 961bi.	— — — Society Islands 176a.
Fig, eelworms in 935a.	— malayi in man, report of case 91a. Filarinematinae n.subf. 1002.
Fiji, filariasis 152a.	Filicapitea n.tribe 1002.
-, helminths in domestic animals 172a.	Fish, Cercaria C in 120g.
Filaria, see also Acanthocheilonema, Dipetalonema, Dirofilaria, Litomosoides, Loa, Microfilaria,	—, cestodes in 49a, 852c.
	— controlling schistosome intermediaries 463a.
Wuchereria. — sp. in Cercopithecus, periodicity 69b.	
- conepati n.sp. in Conepatus suffocans 572a.	-, echinostomatid metacercariae on 690e.
Filariae, effect of diethylcarbamazine on 80a.	-, Glossobothrium n.g., n.sp. in 49a.
— in man, phenosafranine compounds ineffective	—, helminths in 252a, 843a.
357a.	-, metacercariae in 696d.
— — mosquitoes 248a.	—, Marsipobdella n.g., n.sp. on 749a.
— phylogenetic classification 581.	—, Monogenea on 532a, 541a, 713g.
Filariasis & adenolymphocele in man in Belgian	—, nematodes in 564a.
Congo 925b.	-, new trematodes in 568a, Opisthorchis felineus metacercariae in 553d.
-, cerebrospinal, see also Lumbar paralysis,	-, Paracryptogonimus n.sp. in 568a.
Neurofilariasis.	— parasites in 972.
,, in domestic animals 221c, 251a, 283a,	, Pseudamphistomum truncatum metacercariae
569a.	in 553d.
	-, Rhodatrema quadrilobata in 213g.
—, — goat, antimony compounds 702e.	—, trematodes in 588b.
-, -, goat, antimony compounds 702f.	—, Triaenophorus spp. in 643a.
	fresh-water, helminths in 332a.
-, -, in man & domestic animals 358a.	Floriceps uncinatus n.comb. for Rhynchobothrium
-, -, treatment & control 702b.	uncinatum Linton, 1924 49a.
— & chyluria in man 448a.	Formica fusca, 2nd intermediary for Dicro-
solustibosan 2190.	coelium dendriticum 374c.
-, dichlorphenarsine hydrochloride 702i.	Formosa, Schistosoma japonicum 2e.

Fowl, Ascaridia galli in 200, 87d, 150a, 281d.

-, carbon tetrachloride 534a, 536e.

-, Echinoparyphium recurvatum in 936a.

-, penicillin 129a.

—, phenothiazine 534a, 938a. France, Ancylostoma 607b. —, Aphelenchoides ritzema-bosi 325a.

-, Bulinus contortus 852b. -, Cysticercus bovis 197a. -, Dioctophyme renale 492c. -, celworms in plants 57a. -, helminths in man 400a, 607b.

-, Heterodera schachtii 325a.

Francolinus francolinus, Lyperosomum n.sp. in 1001.

- orientalis, Lemdana n.sp. in 961t. French Sudan, schistosomiasis 728b.

- West Africa, filariasis 7b, 362b.

———, hydatid 362d.

————, Onchocerca volvulus 362a. Frog, Echinoparyphium flexum in 230dg. Frostia pellucida n.g., n.sp. 342a.

Fuchsia butschlii n.comb. for Diplogaster butschlii 961bi.

Fulica americana, Amidostomum raillieti in 76a.

Gadus morrhua, Ascarophis n.sp. in 962a. - virens, Hemiurus communis in 97g.

Gaigeria pachyscelis in Aepyceros melampus in

South Africa 434a. Galeichthys seemanni, Fellodistomum n.sp. in 588b. Galeocerdo tigrinus, Staphylorchis n.sp. in 484a. Gallus lafayetti, Leucochloridium n.sp. in 78d. Gambia, helminths in man 152 l.

Gammanema conicauda n.sp. 610d.

Gammarus, Cyathocephalus truncatus in 535a.

pulex, cysticercoids in 120a. Garlic, Ditylenchus dipsaci in 692a.

Gasterosteus aculeatus, Bothriocephalus claviceps in

Gastrolecithus n.g. for Dinobothrium planum Linton, 1922 49a.

Gavia stellata, Syngamus n.sp. in 961 L.

Geotrupinae, helminths in 514b.

Gerbillus pyramidum, Schistosoma mansom in 261i.

Germany, Cysticercus bovis 234a.

-, Dicrocoelium dendriticum 23a. -, helminths in man 737a.

-, - & meat inspection 234a.

—, Heterodera rostochiensis 117d, 393a. -, parasites in animals 1007, 1010.

Gelanostrongylus n.g. for Macropostrongylus macro-stoma 1002.

Germo alalunga, helminths in 418a.

Gerres punctatus, Marsupioacetabulum n.g., n.sp.

Gigantobilharzia huronensis in Spinus tristis & Richmondena cardinalis, life-history 991. Ginglymostoma cirratum, Terranova n.sp. in 491a.

Glandicephalus perfoliatus n.comb. for Diphyllobothrium perfoliatum 72a.
Glaucionetta islandica, Avioserpens n.sp. in 150d.

Glomeris spp., Thelastoma n.spp. in 58d. Glossobothrium nipponicum n.g., n.sp. in fish 49a. Gnathostoma spp. larvae diff_rentiated 402c.

- in man in Japan 338a.

- brasiliense n.sp. in Lutreolina crassicaudata 454b.

doloresi larvae in Hynobius spp. 402a.

- kyushuense, synonym of G. spinigerum 402b.

- spinigerum in cat 67c. -, life-history 402b.

Goat, antimony compounds 702e. -, arsenic compounds 702f.

-, Coenurus cerebralis in 963a. -, Dicrocoelium dendriticum in 23a. -, emetine hydrochloride 230cu.

-, helminths in 878.

—, lumbar paralysis in 155a. -, Marshallagia marshalli in 29e.

-, Oesophagostomum venulosum in 20e.

—, parasites in 993. —, phenothiazine 314a. -, - derivatives 207a.

-, Setaria digitata & lumbar paralysis in 41d.

Skrjabinema ovis in 964a.

Gobius sp., Opegaster n.spp. in 568a. Goffartia heteroceri n.g., n.sp. 867a.

- africana n.comb. for Diplogasteroides africanus

- variabilis n.comb, for Diplogasteroides variabilis 867a.

Gold Coast, see also West Africa.

-, Cysticercus cellulosae 61c. - -, Simulium damnosum 289b.

Gongylonema neoplasticum in rat, pathology 427a.

- pulchrum in cattle in Australia 546d.

— — — man in Morocco 645a. — — — — U.S.A. 261j.

Goose, Amidostomum anseris in 77a, 204b.

-, carbon tetrachloride 528d. , helminths in 961br.

Gordiacea in Austria 158a. - Italy 610c.

Gordionus austriacus n.sp. 158a.

- conglomeratus n.sp. 158a.

— fragmentarius n.sp. 158a. — scripturus n.sp. 158a.

- subalpinus n.sp. 158a. Gordius doriae Camerano, 1897 to Acutogordius n.g. 158a.

- feae Camerano, 1897 to Acutogordius n.g. 158a.

- incertus Villot to Acutogordius n.g. 158a.

- parvaquaticus n.sp. 158a. plicatissimus n.sp. 158a.

Gorgodera (Antodera) n.subg. 1001.

- (Extremodera) n.subg. 1001. (Gorgodera) n.subg. 1001. (Mediodera) n.subg. 1001.

- pawlowskyi n.sp. in Rana sp. 1001.

— (Postodera) n.subg. 1001.

- vivata n.sp. in Rana spp. 982. Gorilla, Oesophagostomum stephanostomum 200b, 200c.

Grape, Meloidogyne in 75a. Grass, Tylenchorhynchus n.sp. on 97d. Grasses, Anguina sp. in 26b.

Greece, hydatidosis 652a.

"Guamo", eelworms in 191a. Guatemala, Onchocerca volvulus 230d.

-, onchocerciasis 176g, 243c.

Guinea-fowl, helminths in 961n. Guinea-pig, Ascaris lumbricoides extract & eosinophilia in 231a. Trichinella in 970. Gyalocephalea n.tribe 1002. Gymnophallinae, systematic position 230dd. Gymnophalloides tokiensis in Ostrea gigas, pathology 714a. Gymnorhynchus gigas plerocercoids in Brama тауі 381а. Gymnostreptus sp., Rhigonema n.sp. in 58d. -, Thelastoma n.sp. in 58d. Gynaecotyle adunca redescribed 230c. -, technique for culturing 20q. Gynaecotylinae n.subf. 961r. Gyrocoelia reviewed 498a. - albaredai n.sp. in Himantopus himantopus 498a. Gyrocotylidea n.ordo 305. Haemadipsa zeylanica on man, repellent 26d. Haematology, ancylostomiasis 340d, 586i. -, Ascaris in pig 953a. - lumbricoides extract 231a. -, Diphyllobothrium 167a. -, Haemonchus contortus 649a. —, helminths in man 260b, 430c, 444a. —, schistosomiasis mansoni 680f, 680g, 680h. —, Strongylus edentatus 377a. —, Trichinella spiralis 230v, 590a. Haematopota larva, intermediary for Porrocaecum crassum 961j. Haemonchiasis in sheep in Australia 787a. Haemonchinae n.subf. 1002. Haemonchus, see also Stomach worms, Trichostrongyle. - & Nematodirus in sheep, effect on resistance - contortus, anaemia-producing substance 649a. - - in cattle, phenothiazine 230ci. - -, survival of free-living larvae 196b. Haemopis sanguisuga, chemistry of skin 635c.
—, structure of teeth 675h. Haemoxenicon chelonenecon n.sp. in Chelonia 20x. - stunkardi n.g., n.sp. in Chelonia mydas 20x. Halalaimus cirrhatus n.sp. 610d. Halocercinae n.subf. 1002. Hamatospiculum flagellispiculosum n.sp. in Myio-dynastes solitarius 572a. Hamster, Schistosoma mansoni in 13b. , Trichinella spiralis in 230v, 230bh, 590a. Haplocleidus reviewed 132h. gomtius n.sp. on Wallagonia attu 912b. Haplonema hamulatum to Cottocomephoronema hamulatum n.comb. 9610. Haplonematidae n.fam. 961o. Haploporoidea revised 339a. Haplorchis, key to spp. 78a. pearsoni redescribed 78a. Haplorchoidinae n.subf. 1000. Haplosplanchnus, life-history 230de. Harmostomum caudale of Diesing to Brachylaemus brauni nom.nov. 320e. Hawaii, Fasciola hepatica 428m., helminths in animals 471b. schistosome dermatitis 35a.

Hedruris mucronifer, morphology 572b.

-- n.sp. in Telmatobius schreiteri 551b, 572a.

Heligmodentostoma n.g. in Microtus sp. 1002. Heligmonellea n.tribe 1002. Helisoma corpulentum, Cercaria n.sp. in 281c. Helleborus niger, Pratylenchus sp. in 479c. Helminth anaemia in man 435a. - enzyme systems 415a. - immunity 295a. - --, general account 326a. - larvae, technique for mounting 74a. - ova in faeces, technique for concentrating 272a, 616a, 733a. ---, -- detecting 655a, 672d. ——— sewage 961bq. --- & swimming pool in Yugoslavia 668a. ----, technique for collecting from faeces 230by. ____, ___ _ soil 843b. ____, ___ concentrating 230s. ——, —— counting 28b.
——, —— fixing 703a.
— physiology (text-book) 304.
Helminthiases in sheep (text-book) 302. Helminthiasis in animals 546c. — — cattle, diagnosis 148b. —, diagnosis 578a. - in domestic animals, control 453c. -, latent 961bd. - in man 1005. ----, chemotherapy 110a. — — in Japan, survey **696f.** — — —, treatment 899a, 906f. — of man, world distribution 16a. - in pig, control 453b. Helminthological research in British Colonies Helminthology in Australia, research reviewed 786s. - & crime detection 116a. -, historical 735b. -, medical (text-book) 557, 873. — in Russia 553c, 726a, 961a, 961b. -, summary of lectures 118f. -, technique for teaching 115a. — (text-book) 871. — in U.S.A., historical review 20w. veterinary (text-book) 876. Helminthoxys effilatus n.sp. in Lagidium sp. 572a. Helminths in Aepyceros melampus 434a. — — animals 813c. — — in Australia 345a. — — —, control 726b, 992. — — —, general account 245a. — — in Hawaii 471b. - of animals, identification 163. — in animals in Ireland 942a. ----, reviewed 516b. - - in Russia 961p. - - - Switzerland 516a. - of animals transmissible to man 600a. - & appendicitis in man 497a, 815a. - in birds, general account (text-book) 166. — — in India 688a. — — —, popular account 577b. - - in Russia 961q. — — carp, experimental 675g. — — cat 501a. _ - - & dog 251e. ____ in U.S.A. 428n.

·	
Helminths in cat in U.S.A. 251b.	Helminths in man in Japan, literature reviewed
cattle in Australia 412a.	696h.
, control 154a, 538a, 698a.	Madeira 586b.
, general account 93a, 154b.	Mozambique 586c, 586d, 586j.
———, resistance 65a.	Poland 525a.
	, popular account 388a.
— — in U.S.A. 189a, 777c.	& primates 622a.
——————————————————————————————————————	, pulmonary lesions 340a.
— — Cetacea 961v, 961w.	———— in Sicily 785d.
— — Colinus virginianus 840a.	
—— deer in Switzerland 276a.	——————————————————————————————————————
— — Dipodomys spp. 230br.	Spain 451a.
— & disease in sheep 209b.	———— Thailand 230bn, 230bo.
— in dog 95b.	———, treatment 811e, 946c, 946d.
in Algeria 63a.	— — —, — reviewed 838a.
, general account 118e.	— — in U.S.A. 936b.
— — in Spain 135a.	——————————————————————————————————————
, 6-tertiary-butyl-m-cresol 261c.	———, Vermella 735c.
———, treatment 28a.	— — in Yugoslavia 909a.
in U.S.A. 28b, 95c.	- & meat inspection in Germany 234a.
— domestic animals 40c, 245b, 245d, 629a.	———— (text-book) 303.
	in U.S.A. 2222.
———— in Australia 67a.	- in medicine, historical account 769a.
————, control 138c, 961bo.	
	— — mink 678c.
————, diagnosis 662a.	— mosquitoes 34a.
———— in Fiji 172a.	Ondatra in Russia 553a.
——————————————————————————————————————	zibethica 554a.
——————————————————————————————————————	in U.S.A. 20 L
— — Eopsetta jordani 20m.	— Ovis canadensis in U.S.A. 132g.
— — fish in Canada 252a.	- & paradental disease in man 929a.
———— U.S.A. 843a.	- in pig 284a, 678a.
- fresh-water fish in Britain 332a.	———, control 14a.
— — Geotrupinae 514b.	popular account 96b.
— — Germo alalunga 418a.	— — —, symptomatology 507a.
-, glycogen distribution 864g.	— — in U.S.A., general account 486a.
— in goat in Fiji, phenothiazine 314a.	— pigeon in Brazil 807a.
— goose in Kussia 961br.	— Pinnipedia 961v.
— guinea-fowl in Russia 961n.	— — poultry 903a.
— horse, carbon tetrachloride with turpentine	, control 142.
oil 492b.	— —, effect of built-up litter 322, 259a, 858a.
———, control 755a.	——— in Egypt 20be.
———, effect on growth 537a.	———— India 92a.
— — Leptodactylus pentadactylus 583a.	———, popular account 178a, 468a.
— Leucichthys artedi, new records 428 1.	———— (text-book) 556.
— — mammals 927c.	— — rabbit 324b.
— — in Russia 961i.	— — Rattus norvegicus in Aleutian Islands 99a.
man 2a, 358d, 358e, 399f, 399h, 616b,	——————————————————————————————————————
616c, 672a, 750a.	Yugoslavia 808a.
— —, allergic reactions 366a.	rodents in Alaska 428e.
, symptoms 190e, 399e.	— ruminants, diagnosis 866a.
— — —, atebrin-924a.	— Russia, oecology 961bm.
——— in Brazil 343a, 399c.	—— sheep 374b.
——————————————————————————————————————	
——————————————————————————————————————	———, effect of supplementary diet 356a.
	in Britain, general account 142a.
———— Colombia 922a.	———, control 83b, 155e, 165f.
——————————————————————————————————————	———, effect on weight gains 214a.
— — in Czechoslovakia 201a, 932a.	——— & goat, symptoms & control 878.
——— & dog in Chile 190a.	— — —, phenothiazine 10a.
— — —, electrical treatment 145b.	———, — photosensitization 473a.
— —, cosinophilia 260b, 430c, 444a.	, seasonal variations 155d.
——— in France 607b.	——— in U.S.A. 10e.
—————, incidence 400a.	— — Sus scrofa 961bg.
———— Gambia 152 L	—, technique for detecting 230r.
———, general account 697a.	- disgressing 161 200h
in Germany, post-war increase 737a.	—, — diagnosing 161, 399b.
, hetrazan 353a.	-, - staining & mounting 146a.
	— in Tarsius carbonarius 488a.
——— in Iran 294d.	-, vitamin B ₁₂ content 567a.
——————————————————————————————————————	- in zoo animals 488e, 488f.

Hemicycliophora emended 201a.	Heterodera rostochiensis in potato, control 60c.	
- parvana n.sp. on celery 261a.	————, D-D 257n.	
Hemigymnostreptus sp., Ichthyocephalus n.sp. in	, _ & ethylene dibromide 331a.	
58d.	, general account 194a, 667a, 717a, 984.	
Hemiurus communis in Gadus virens 97g.	———— & other plants in Peru 351a.	
Herpestes cafra, Metathelazia n.sp. in 58j.	———— in Peru 479b.	
Heterakis gallinae & "blackhead" in turkeys,	———, resistant varieties 447a.	
2-amino-5-nitrothiazole 10b.	——— soil, effect of crop rotation 257c.	
, new drugs 10c.	—— in Solanum, resistant crosses 518a.	
, new drugs 10c.	spp., resistant lines 463c.	
against 10c.	———— tuberosum, resistant subspecies 463d.	
— in pheasant 118b.	, survival affected by temperature 300.	
——, respiratory metabolism 416a.	, technique for field sampling 164a.	
—— in turkeys, phenothiazine 312a.	— in tobacco, resistance 760b, 760c.	
Heterodera cysts, technique for collecting 97b.	tomato, gall formation 257i.	
, detecting 479d.	- schachtii, see also H. glycines.	
——, —— dissecting 97a.	— in Beta cicla 943a.	
——, —— examining 97a.	——, bionomics 870a. ——, control regulations in Britain 15a, 46.	
— in plants in Canada 277a.	hoes made 31a	
-, technique for inoculating plants with 149a.	— —, host range 31a.	
- sp. in legumes 794a.	— —, new host records 794a. — in sugar-beet, control 678d.	
——— tobacco 257).	———— in France 325a.	
& tomato 126a.	, general account 984.	
- spp., differentiation 164f, 205a.	———, popular account 475a.	
——, — of cysts 164g.	in Sweden 886a.	
——, hatching factor 205c.	Heterodontus zebra, Phyllobothrium n.sp. in 49a.	
— avenae, see also H. major.	Heterodorus magnificus n.g., n.sp. 384a.	
—— in oats, control 476b.	Heterophus to Mediorhynchus 784b.	
- cruciferae in Cruciferae in U.S.A. 760a.	Heterophyoidea reviewed 961bn.	
— glycines, see also H. schachtii.	Heterotyphlum eurycheilum n.sp. in Promicrops	
— in soya bean, disease index 265a.	itaiara 491a.	
— n.sp. in soya bean in Japan 254a.	Heth artigasi n.sp. in Rhinocricus sp. 58d.	
— major, see also H. avenae.	perarmatum n.sp. in Khinocricus 38a.	
—— in oats in Sweden 533b.	- schubarti n.sp. in Rhinocricus cachoeirensis 58a.	
— marioni, see also Meloidogyne. — in carnation, resistant varieties 759a.	— travassosi n.sp. in Leptodesmus jucundus 58d.	
, effect of D-D on population 3a.	- transfilhai n.sp. in Rhinocricus padvergi 380.	
- & H. rostochiensis, control by soil fumigation	- tuzetae n.sp. in Pseudonannolene tricolor 58a.	
205b.	Hexabothriinae Price, 1942, emended 532a.	
— in lespedeza, resistant varieties 144a.	Hexatylinae n.subf. 961bi.	
—— ova in faeces of man 111a.	Hexatylus vigissi n.sp. in plants in Russia 961bj.	
—— in potato in Australia 312b.	Hibiscus snn., eelworms in 1261.	
soil, control by electric current 383a.	Himantopus himantopus, Gyrocoelia n.sp. in 4968.	
sweet-potato, varietal resistance 246a.	Hippeastrum, Rotylenchus n.sp. on 31c.	
— — tobacco in Australia 263a.	Hirudinea in Belgian Congo, monographed 3938.	
Valeriana officinalis 119a.	— — Burma 414a.	
— munctata in Holland 117a.	-, catalase activity 635d.	
- rostochiensis attacked by amoeboid organism	—, copper content 635e.	
120c, 249a.	in Italy 610c.	
——, bionomics 164b.	-, nomenciature 588d.	
——, control 5a, 257t, 373a.	-, physiology 208c, 208d.	
, _ by nematicides 164d.	-, popular account 908a.	
, - regulations in Britain 44, 45.	— in Switzerland 651c.	
Germany 117d.	, taxonomy 472a.	
——————————————————————————————————————	-, trematodes in 651c.	
	Hirudo spp., bacterial fauna 979.	
- cysts on Convallaria, hot-water treatment	——, physiology 978.	
1176	- lefeurei n.sp. 595a.	
effect of potato root leachings 257g.	- medicinalis in Britain 462a, 519a.	
effect of crop rotation on population 10-10-	, symbionts 43a.	
, factors controlling population 1170.	Hirundo rustica, Hymenolepis n.sp. in 688a.	
— — general account 508b.	Histology, Echinococcus granulosus 765c.	
- in Germany, popular account 3832.	-, Raillietina urogalli 213e.	
——————————————————————————————————————	History, Schistosoma 103b.	
in Lycopersicum, resistance 25 / Q.	Holland, eclworms in plants 446a.	
means of spread 104c.	-, Heterodera punctata 117a.	
—— in plants, general account 476a.	-, - rostochiensis 386a.	

```
Hydatid in man & animals in Canada 656a.
Holostephanus ictaluri n.sp. in Ictalurus punctatus
                                                                                                        — —, diagnosis 352a, 451b.
— — in French West Africa 362d.
Hookworm, see also Ancylostoma, Creeping
                                                                                                         — —, immunology & diagnosis 748a.
— — in Lebanon 229c.
        eruption.
- anaemia in man 22a, 576a, 586h, 826a, 945a,
                                                                                                        ———, surgery 186a, 186b, 425a, 574b, 694c, 701b, 727d, 765e, 768a, 768d, 829b, 882a.
———, technique for removing 682a.
____, iron therapy 531b.
___ in Malaya 449a.
                                                                                                         - -, treatment 574d, 959a.
_____, treatment 762a.
                                                                                                      — — in Turkey 694b.
- disease, see also Ancylostomiasis.
                                                                                                      --- U.S.A. 602a.
-- in China 552b.
                                                                                                      - in man in Brazil 322b.
- membrane, chemistry 398a.
                                                                                                      — in pig 245c.
— — —, Diospyros mollis berries 230bp.
— — —, hexylresorcinol & tetrachlorethylene
                                                                                                          - - in Uruguay 631a.
                                                                                                         - — rabbit, experimental 597b.
      270c.
                                                                                                         - - sheep 113b.
   - - in Lebanon 705b.
                                                                                                      Hydatidosis, control 598a, 781a.
Horse, carbon tetrachloride 850f.
—, — — with turpentine oil 492b.
—, cerebrospinal filariasis & kumri in 221a.
                                                                                                      - in domestic animals, fertility of cysts 153a.
                                                                                                      — — Greece, control 652a.
— — Yugoslavia, control 850d.
-, Dictyocaulus in 536h.
-, helminths in 537a, 755a.
-, Onchocerca cervicalis in 243d, 243e.
                                                                                                      Hydromys chrysogaster, Cosmocephalus n.sp. in
                                                                                                              8415.
-, - reticulata & periodic ophthalmia in 832a.
                                                                                                      Hyla arborea, Polystoma n.sp. in 269b.
-, Parascaris equorum, in 702d.
                                                                                                      Hymenolepididae reviewed 961g.
                                                                                                      Hymenolepis, protein metabolism 230co.
-, phenothiazine 40a.
-, Setaria digitata & lumbar paralysis in 41d.
                                                                                                      - spp., morphological variations 39a.
                                                                                                      — in rat in Chile 944a.
— ababili n.sp. in Hirundo rustica 688a.
-, strongyle larvae in 118d.
—, strongyles in 198b, 998.
—, Strongyloides westeri in 880c.
                                                                                                      - barrowensis n.sp. in Anser albifrons 20d.
-, strongylosis in 961bl.
                                                                                                      - californicus cysticercus in Artemia salina 436a.
-, Strongylus edentatus in 83a, 377a.
                                                                                                      - - valid 230 L
  -, - equinus in 118c.
                                                                                                      - cordobensis n.sp. in pigeon 33b.
                                                                                                      - crecca n.sp. in Nettion crecca 688a.
-, - vulgaris in 41b.
                                                                                                      — creplini in Anser albifrons 20d.
— diaphana to Ditestolepis n.g. 329a.
-, technique for drenching 67b.
-, toluene 177a.
  -, Trichonema (Cylicocyclus) radiatum in 490a.
                                                                                                      - diminuta, abnormal eggs 208b.
   -, Triodontophorus in 630a.
                                                                                                      - - in Citellus leucurus 428g.
Hungary, Trichinella spiralis 309a. "Husk" in cattle in Britain 51a.
                                                                                                       - -, cytochrome oxidase 385b.
                                                                                                      — — in man 344a.
Hydatid, see also Echinococcus.
                                                                                                         --- in Chile 190g.
-, alveolar, in man 613a.
                                                                                                      - - Muridae in India 344a.
— in animals, effect on milk 823a.
                                                                                                         - —, physiology 230bu.
- antigen, technique for preparing 349a.
                                                                                                        - in rat, effect of gonadectomy & gonadal
— in brain, technique for removing 612a.
                                                                                                          hormones 13a.
                                                                                                         ———, physiology 230bs.
———, pumpkin seed extract ineffective
— — cattle in Australia 348a.
   - -, protein changes 618a.
   - — domestic animals 577c.
                                                                                                              887a.
 - doineste annias 5//c.
- - in Cyprus 82a.
- - - Italy 275a.
- - Uruguay 628a.
- - Man 59d, 106a, 137a, 137b, 180b, 188a, 190f, 233a, 233b, 233c, 233d, 233f, 233g, 233i, 267a, 268a, 268b, 274a, 352c, 403a, 404b, 423c, 428b, 420a, 455c, 406b, 574a, 420a, 455c, 406b, 420a, 455c, 406b
                                                                                                     — fraterna, see also H. nana.
                                                                                                     — — & Trichinella spiralis in mouse 230bi.
                                                                                                     - gaugi n.sp. in Turdoides somervillei 688a.
                                                                                                     - hamanni for H. neomidis 651b.

    horrida, morphological variation 428u.
    integra for H. polyacantha 651b.

       233i, 267a, 268a, 268b, 274a, 352c, 403a, 404b, 423c, 438b, 439a, 451c, 496b, 574a, 574c, 574c, 574c, 574c, 574s, 574h, 574i, 576c, 576d, 576c, 579a, 587a, 587b, 591a, 596c, 607a, 611a, 612b, 612d, 619a, 620a, 634a, 634b, 634c, 640a, 641a, 660a, 665a, 671a, 671b, 672b, 672c, 701a, 706b, 715a, 715b, 716a, 724a, 727a, 727b, 727c, 730a, 732a, 732b, 732c, 734a, 736a, 738a, 741a, 741c, 745a, 763a, 768b, 780a, 797b, 804a, 804b, 806a, 818a, 818b, 820b, 820c, 820c, 829c, 836a, 837a, 862a, 863b, 918a, 933b, 947a, 951a, 951b, 952a, 952b, 953d, 954a, 959b.
                                                                                                     - intricatus n.sp. in Sorex vagrans vagrans 21a.
                                                                                                     - johnsoni n.sp. in Microtus pennsylvanicus
                                                                                                             drummondii 21b.
                                                                                                         - kenaiensis n.sp. in Aythya marila nearctica
                                                                                                     — kenki n.sp. in Sorex vagrans vagrans 21a.

macyi n.sp. in Sorex vagrans vagrans 21a.
magna n.sp. in Acridotheres tristis 688a.

                                                                                                     — makundi n.sp. in duck 688a.
— nana, see also H. fraterna.
                                                                                                        - in man, atebrin 105b.
       951a, 951b, 952a, 952b, 953d, 954a, 959b, 959c, 959d, 960b.
                                                                                                        - - ova in man in Brazil 680a.
                                                                                                        - neomidis to H. hamanni 651b.
```

Hymenolepis olsonis n.sp. in Scapanus townsendi 428d.

- omissa, life-history 651b.

oregonensis n.sp. in Ondatra zibethica 428d.
 pitymi n.sp. in Pitymys pinetorum 104c.

- polyacantha to H. integra 651b.

- sphenomorphus n.sp. in Sorex vagrans vagrans

Hymenophysa curta, Bothriocephalus n.sp. in 49a. Hynobius spp., Gnathostoma doloresi larvae in 402a.

Hyostrongylea n.tribe 1002.

Hyperoodon ampollatus, Delamurella n.g., n.sp. in

Hypomesus olidus, Philometra opsalichthydis in 695a.

Iceland, acanthocephalans in animals 1009. Ichthyocephalus almeidai n.sp. in Hemigymnostreptus sp. 58d.

ichthyocephaloides n.sp. in Rhinocricus sp. 58d. Ichthyotrema vogelsangi n.g., n.sp. in Xesurus punctatus 588a.

Ichthyotreminae n.subf. 588a.

Ichtyobronema conoura to I. gnedini nom.nov.

gnedini nom.nov. for I. conoura 9610.

Ictalurus punctatus, Holostephanus n.sp. in 230f. Iheringtremea n.tribe 1000.

Ilisha elongata, Pterobothrium n.sp. in 49a. Immunity, Ascaridia galli 87d, 150a.

-, Dictyocaulus filaria 120b. -, Ditylenchus destructor 794a. -, — dipsaci 164h, 921a, 973. -, eelworms 126f, 483a.

- in tobacco 86a.

-, Haemonchus & Nematodirus 230p.

-, helminths 295a, 326a. - in cattle 65a.

-, Heterodera marioni 144a, 246a, 759a. — rostochiensis 257d, 447a, 463c, 463d, 518a, 760b, 760c.
— schachtii 31a.

-, Hymenolepis fraterna & Trichinella spiralis 230bi.

-, Litomosoides carinii 230ba.

, Meloidogyne 75a, 257m, 483e, 636a, 636b, 770a, 771b.

-, — spp. 30k, 126e, 128a. -, — incogmita 242a, 401a.

-, Schistosomatium douthitti 230cz, 424a.

-, schistosomes 230ct.

, Trichinella spiralis 225a, 225b, 230t, 230bd, 230be, 230bf, 230bg, 419a, 419b, 775a, 961c, 961d.

—, Trichostrongylus axei 19c. —, — retortaeformis 120d, 120e, 155c.

Immunology, ascariasis 845a, 918b.

—, Ascaris 403b.

-, Cysticercus tenuicollis 453d.

-, helminths 98b. -, hydatid 748a.

-, Schistosoma mansoni 19b. -, Trichinella spiralis 371b, 995.

-, trichinelliasis 549a.

Incidence, enterobiasis 43b.

-, helminthiasis in man 16a. , helminths in man 400a.

India, ancylostomiasis 220a, 423a.

-, Cysticercus cellulosae 219a. -, helminths in birds 688a.

—, — — poultry 92a.

-, Hymenolepis diminuta 344a. —, lumbar paralysis 92b. —, Marshallagia marshalli 29e. —, Ornithobilharzia n.sp. 29f.

-, schistosomiasis haematobia 689a.

-, Thelazia n.sp. 221b.

-, Wuchereria bancrofti 586t. Indo-China, filariasis 58g. -, Schistosoma spindale 273a.

Indonesia, Spirocerca lupi 395a. Inermicapsifer cubensis in rat 270a, 811b. Insectivores in Hungary not carrying Trichinella

309a. Insects, 2nd intermediaries for Gorgodera vivata 982.

mermithids in 852a.

Iran, helminths in man 294d.

, schistosomiasis haematobia 229a.

Iraq, eelworms in plants 216a.

-, schistosomiasis haematobia 227a, 229a.

-, trichostrongylosis 420a.

Ireland, Anguillulina tritici 700a.

-, eelworms in plants 313a, 313b, 942a.

, helminths in animals 942a. Iris, bulbous, eelworms in 141a.

Isopods, 2nd intermediaries for Conspicuum icteridorum 20bd.

Italy, ancylostomiasis 820a.

-, Cystocaulus nigrescens 618d. —, Echinococcus granulosus 275a. —, enterobiasis 275b.

-, Enterobius 514d, 822a.

-, - vermicularis 34c, 785e. -, free-living eelworms 610d.

-, Gordiacea 610c.

-, helminths in man 48a, 514a.

-, Hirudinea 610c.

-, hydatid in domestic animals 275a.

-, parasitic enteritis 597a.

-, trichinelliasis 785a, 785b.

Japan, Ascaris lumbricoides 396a, 396b, 396c, 396d.

-, cestodes in fish 49a. -, Chordodes spp. 864a.

—, — n.sp. 603b.

-, Gnathostoma spp. 338a.

-, helminthiasis in man 696f. -, helminths in domestic animals 926a.

- man 696h.

-, Heterodera n.sp. 254a. —, paragonimiasis 696g, 831a.

-, Stephanurus dentatus 905a. , Wuchereria malayi 409a.

Johanssonia abditovesiculata n.sp. on Tetraodon hispidus 749a.

Julus sp., Brumptaemilius n.g., n.sp. in 58d.
—, Thelastoma n.sp. in 58d.

- terrestris, Severianoia n.sp. in 58d.

Kenya, free-living eelworms 342a. Ketupa zeylonensis, Dispharynx n.sp. in 912d. Korea, spargana in man 230bt. Kowalewskiella emended 33a. Krusadia indica n.g., n.sp. in Caprimulgus indicus Lagidium sp., Helminthoxys n.sp. in 572a. peruanum, Paranoplocephala n.sp. in 713c. Lampsilis siliquoidea, intermediary for Rhipi-docotyle septpapillata 529a. Langeronia provitellaria n.sp. in Rana 281b. Lanicides vayssierei, Cercaria n.sp. in 230i. Lapwingia reticulosa n.g., n.sp. in Lobipluvia malabarica 688a. Larus marinus, Contracaecum n.sp. in 851a. Larva migrans, see Creeping eruption. Lateolabrax japonicus, Bothriocephalus n.sp. in Lates calcarifer, Opecoelus n.sp. in 568a. -, Pseudometadena n.g., n.sp. in 568a. Lebanon, filariasis 817a. -, hookworm 705b. -, hydatid in man 229c. -, trichinelliasis 482b, 705a. Lecanicephala n.ordo 305. Lechriorchis primus, development 101a. Lecithochirium lobatum n.sp. in Sphyraena sp. Legumes, Meloidogyne spp. in 126b. Lemdana skrjabini n.sp. in Francolinus 961t. Lemmus spp., Paranoplocephala n.sp. in 428e. Lemniscomys striatus, intermediary for Taenia brauni 819a. Leopard, ancylostomiasis in 642a. Leparis leparis, and intermediary for Prosorhynchus squamatus 213d. Leptodactylus ocellatus, Rhabdias n.sp. in 572a.

— pentadactylus, helminths in 583a. Leptodesmus jucundus, Clementeia n.spp. in 58d. - -, Heth n.sp. in 58d. - paulistus, Rhigonema n.sp. in 58d. Leptophyllinae to Travtrematinae nom.nov. 981. Lespedeza, Heterodera marioni in 144a. —, Meloidogyne in 483e. Lethrinus sp., Plagioporus n.spp. in 568a. — —, Podocotyle n.sp. in 568a. -, Pseudolepidapedon n.sp. in 568a. Leucasiella mironovi n.g., n.sp. in Delphinapterus leucas 961z. Leucichthys spp., Triaenophorus crassus plerocercoids in 421a. - artedi, helminths in 428 1. Leucochloridiinae revised 320e. Leucochloridium, key to spp. 320e. - beauforti n.sp. in Ammospiza maritima 101d. - ceylonicum n.sp. in Gallus lafayetti 78d. - heckerti n.sp. for Distomum macrostomum of Heckert 320e. — insigne to L. heckerti 320e. - paradoxum, life-history 25a. — soricis n.sp. in Sorex araneus 329a. - tetrastae n.sp. in bird in Russia 961bt.

Libya, schistosomiasis haematobia 327b.

Life-history, Anguina agrostis 126c.

-, Ascaridia galli 230a, 281d.

-, ascarids 98d, 961k.

-, Calicophoron ijimai 151a. -, Capillaria contorta 230w. -, Centrorhynchus spinosus 676a. -, Cercorchis 262b. —, cestodes 327c. —, Cionella lubrica 374a. -, Cloacitrema michiganensis 230n. -, Conspicuum icteridorum 20bd. -, Cysticercus madoquae 618e. -, Dicrocoelium dendriticum 20n, 81c, 81d, 209a, 230ch, 374c. -, Dipetalonema blanci 79c. - evansi 536b. -, — evansi 536b. -, "Diplostomum volvens" 120g. -, Dirofilaria immitis 230dk. -, Echinococcus granulosus 27a. -, Echinoparyphium flexum 230dg. -, Fasciola gigantica 912a. -, - hepatica 679a. -, Fellodistomum fellis 553b. -, Gigantobilharzia huronensis 991. -, Gnathostoma spinigerum 402b. -, Gorgodera vivata 982. -, Haplosplanchnus 230de. -, Hymenolepis omissa 651b. -, Lechriorchis primus 101a. -, Leucochloridium paradoxum 25a. -, Litomosoides carinii 38b, 38c, 38f. -, Megapera 230de. —, Meloidogyne spp. 30 l. —, Microbilharzia chapini 101f. -, Moniezia expansa 165b. -, Monoecocestus spp. 20y. -, - sigmodontis 230h. -, Multiceps endothoracicus 675c. —, Neoleucochloridium problematicum 37a. -, Onchocerca spp. 50a. -, - gutturosa 157a. -, Ochetosomatidae 713e. -, Opisthorchis felineus 553d. -, Paragonimus westermanii 696b, 696 l. -, Parascaris equorum 880b. -, Patagifer bilobus 675b. —, Physaloptera hispida 428h. -, Physocephalus sexalatus 961m. -, Plagiorchis maculosus 859b. -, — (Multiglandularis) megalorchis 29c. -, — muris 696c. -, Porrocaecum 961e. -, - crassum 213f, 961j. -, Prohemistomum chandleri 230f. --, Prosorhynchus squamatus 213d. -, Proteocephalus exiguus 380b. -, - primaverus n.sp. 428t. -, Pseudamphistomum truncatum 553d. -, Ratzia joyeuxi 12a. -, Rhipidocotyle septpapillata **529a.** -, Rhodatrema quadrilobata **213g.** -, Schistosomatium douthitti 55a, 320b. -, Stegodexamene anguillae 428c. -, Stomatrema guberleti 981. -, Strongyloides ratti 359a. -, Strongylus edentatus 83a. -, Syphacia obvelata 52a. -, Taenia brauni 819a. -, - jakhalsi 514c, 618e.

Life-history, Brachylaemus virginiana 20g, 323a.

- Life-history, Telogaster opisthorchis 428c. -, Thelazia skrjabini 213b, 536c. -, Triaenophorus spp. 643a.
 - -, Trichobilharzia n.sp. 158b. -, Wuchereria bancrofti 674b.

Liga emended 33a.

Lima bean, Meloidogyne incognita in 257a, 261e. Limnaea acuminata, intermediary for Fasciola gigantica 912a.

luteola, intermediary for Ornithobilharzia dattai 29f.

- palustris, Cercaria n.sp. in 20bb.

-, embryology 372b.

stagnalis, intermediary for Trichobilharzia szidati 158b.

- truncatula, ecology 165g. Limnaeid snails, molluscicides tested against 20h. Limnatis ealensis n.sp. 595a.

Limnephilus, Cercaria goodmani metacercariae in 230dh, 230di.

Litomosoides, see also Filaria, Microfilaria.

- carinii in Bdellonyssus bacoti, infection rate 983c.

-, embryology 38f. - -, life-history 38b, 38c.

- in Sigmodon, effect of superinfection 152c.

----, MSb 152d.

——— hispidus, immunity 230ba. ————, MSb as prophylactic 334d, 334g. Littorina pintado, Cercaria littorinalinae in 35a. Liver-fluke, see also Dicrocoelium, Fasciola.

- in domestic animals 904b, 920a. - - sheep, carbon tetrachloride 546b.

Loa, see also Filaria, Microfilariae. - loa in man, hetrazan 379c, 465b.

---, microfilariae in cerebrospinal fluid 328d.

Loaiasis in man, general account 289c.

---, hetrazan 544c.

Lobipluvia malabarica, Lapwingia n.g., n.sp. in 688a.

-, Progynotaenia n.sp. in 688a.

Lobodon carcinophagus, Baylisia n.g., n.sp. in 72a. Longibucca lasiura in Corynorhinus rafinesquii & Myotis lucifugus 261k.

Longicyatholaimus lineatus n.sp. 610d.

Longistriata adunca in Sigmodon hispidus 20r. - neotoma n.sp. in Neotoma floridana 320d.

Longistriatea n.tribe 1002. Longistriatinae n.subf. 1002.

Lucerne, Ditylenchus sp. in 126b. -, - dipsaci in 973.

Lumbar paralysis, see also Filariasis, cerebrospinal, Neurofilariasis.

- in goat, treatment 155a. - - ruminants in India 92b.

- — & Setaria digitata 702.

Lungworms, see also Aelurostrongylus, Crenosoma, Dictyocaulus, Metastrongylus.

- in cattle 902a, 904a, 920b. ———, treatment 1008.

- — domestic animals, popular account 238a.

- - pig 471a. ——, control 239a.

- - sheep 751a. - - & goat, emetine hydrochloride 230cu.

----, treatment 990.

Lungworms in sheep in Turkey 288a.

_ _ _ U.S.A. 772b.

Lutianus analis, Raphidascaris n.sp. in 491a. Lutreolina crassicaudata, Gnathostoma n.sp. in 454b.

Lutztrema sp. of Boyd, 1951 to Lutztrema sturni n.sp. 1001.

· sturni n.sp. 1001.

Lycopersicum, eelworms in 483a. —, Heterodera rostochiensis in 257d.

—, Meloidogyne spp. in 478b. —, — incognita in 242a.

Lymnaea, see Limnaea.

Lynx lynx, Petrowospirura n.g., n.sp. in 961be. - rufus, parasites in 854b.

Lyperosomum colorosum to Brachylecithum colorosum n.comb. 1001.

- donicum to Brachylecithum donicum n.comb. 1001.

- eophonae to Brachylecithum eophonae n.comb. 1001.

- lobatum glandarii to Brachylecithum lobatum glandarii n.comb. 1001.

- mosquensis cineli n.subsp. in Circlus pallasi 1001.

- petrovi n.sp. in Francolinus francolinus 1901. — schikhobalovi n.sp. in Alectoris graeca 1001. Lytocestidae n.fam. 305.

Macracanthorhynchus, protein metabolism 230co. hirudinaceus, glycogen consumption 428k.

Macrobdella decora in Mexico 588d.

Macroderinae nom.nov. 981.

Macropostrongylus macrostoma to Gelanostrongylus n.g. 1002.

Madagascar, schistosomiasis 739a.

-, - mansoni 367e.

, Wuchereria bancrofti 367d. Madeira, helminths in man 586b. Maize, Pratylenchus zeae in 760d. Malaya, hookworm anaemia 449a.

-, parasitology 131a.

Maldive Islands, filariasis 654a. Mammals, Acanthocephala in 230bv.

—, helminths in 927c, 961i.

, nematodes in 586x. Mammanidula asperocutis n.g., n.sp. in Sorex sp.

Mammomonogamus auris n.comb. for Syngamus auris 961 1.

Man, Acanthocheilonema streptocerca in 38k.

- acridine 742a.

_, _ derivatives 765a.

--, ACTH 467c.

-, Ancylostoma in 607b.

—, — duodenale in 58e, 586e, 694a.
—, ancylostomiasis in 220a, 260a, 330a, 340c, 340d, 340e, 350b, 423a, 465a, 482a, 580a, 586f, 586g, 586i, 589a, 611b, 633a, 657a, 657b, 725a, 810a, 820a, 820d.

-, anthiomaline 624b.

-, antrypol 68a, 152m, 586p. -, arsenamide 54a.

-, Arsobal 624a.

-, ascariasis in 190c, 406a, 438a, 594a, 834a, 845a, 906a, 911a, 918b, 946a, 953c, 956a, 969b.

```
Man, gentian violet 514f, 946e.
Man, ascaridole 827c.
                                                                                                 -, glucantime 566a.
-, ascarids in 294b.
-, Ascaris in 106b, 266a, 310a, 405a, 423b,
                                                                                                    -, Gnathostoma spp. in 338a.
       544d, 576b, 608a, 670a, 670b, 682b, 719b,
                                                                                                  -, Gongylonema pulchrum in 261j, 645a.
                                                                                                 —, helminth anaemia in 435a.
       743a, 752a, 757a, 845b, 861a, 884a.
                                                                                                 —, helminthiasis in 110a, 696f, 899a, 906f, 1005.
  -, - & appendicitis in 958a.

    helminths in 2a, 20p, 36a, 48a, 143a, 145b,
    152 l, 190a, 190e, 201a, 230bn, 230bo,
    294d, 343a, 352b, 358d, 358e, 366a, 388a,

   -, - intestinal occlusion in 685a, 710a, 741b,
       901a.
—, — larvae in 515a.
                                                                                                         399c, 399f, 399h, 400a, 451a, 514a, 521a, 525a, 586b, 586c, 586d, 586j, 600a, 607b,
 —, — & linitis plastica in 735d.
—, — lumbricoides in 107a, 190h, 270b, 396a,
                                                                                                        616b, 616c, 622a, 659a, 672a, 696h, 697a, 737a, 750a, 785d, 811e, 838a, 909a, 922a, 932a, 936b, 946c, 946d.
        396b, 396c, 396d.
-, atebrin 94a, 105b, 210b, 352e, 637a, 924a,
                                                                                                   -, - & appendicitis in 497a, 815a.
 -, Atrimon 210a, 718a.
                                                                                                    -, — eosinophilia in 430c.
-, bilberries 470a.
—, Capillaria aerophila in 145a.
                                                                                                    -, - - paradental disease in 929a.
                                                                                                          - - pulmonary lesions in 340a.
-, carbon tetrachloride 586y, 761a.
                                                                                                 -, Heterodera marioni ova in faeces 111a.
-, cashew nut-shell oil 690c.
                                                                                                   -, hetrazan 152h, 176e, 235a, 308a, 321a, , 353a, 358b, 379a, 379c, 399g, 405b, 430j, , 450d, 465b, 531a, 531c, 544c, 674a.
-, cerebrospinal filariasis in 358a.
—, cestodes in 571a, 766a, 968a.
-, chloroform 598b.
-, chloromycetin 499a.
                                                                                                 -, hexamethylenetetramine 720a.
                                                                                                 —, hexylresorcinol 136a, 185a, 270e, 270f, 294c, , 811a, 811c, 814a, 946b.

—, — & tetrachlorethylene 270c.
-, Contaverm 735a.
 -, cortisone 53a.
-, creeping eruption in 367a, 544a, 584a.
-, cysticerciasis in 108a, 193a, 404a, 496a,
                                                                                                 -, hookworm in 322b, 705b.
                                                                                                          - anaemia in 22a, 449a, 531b, 576a, 586h,
        673a, 830b, 891a, 923a, 928a, 933a, 960a,
                                                                                                         762a, 826a, 945a, 953b.
                                                                                                 -, hydatid in 59d, 106a, 137a, 137b, 180b,
 —, Cysticercus bovis in 340f.
                                                                                                         186a, 186b, 188a, 190f, 229c, 233a, 233b, 233c, 233d, 233e, 233f, 233g, 233h, 233i, 267a, 268a, 268b, 274a, 352a, 352c, 362d, 403a, 404b, 423c, 425a, 438b, 439a, 451b,
-, - cellulosae in 61c, 219a, 396e, 719a.
-, diamino-diphenyl-sulphone 690a.
 -, Dicrocoelium dendriticum in 363a.
 -, Diospyros mollis berries 230bp.
                                                                                                         451c, 496b, 574a, 574b, 574c, 574d, 574e, 574f, 574e, 574h, 574i, 576c, 576d, 576e, 579a, 587a, 587b, 591a, 596c, 602a, 607a, 611a, 612a, 612b, 612d, 613a, 619a, 620a,
 -, Diphyllobothrium in 167a.
 -, - & anaemia in 168a.
        — latum in 121a.
-, Dipylidium caninum in 190b, 803a, 815b.
 -, Dirofilaria conjunctivae in 2b.
                                                                                                         634a, 634b, 634c, 640a, 641a, 656a, 660a, 665a, 671a, 671b, 672b, 672c, 694b, 694c,
 —, Dracunculus medinensis in 38e, 180c, 224a,
        690b, 833a.
                                                                                                         701a, 701b, 706b, 715a, 715b, 716a, 724a, 727a, 727b, 727c, 727d, 730a, 732a, 732b, 732c, 734a, 736a, 738a, 741a, 741c, 745a, 748a, 763a, 765e, 768a, 768b, 768d, 780a, 797b, 804a, 804b, 806a, 818a, 818b, 820b, 820a, 820a, 820a, 826a, 
-, Egressin 200a, 232a.
 -, emetine 271a, 797a.
 —, — hydrochloride & sulpha drugs 696i.
 -, enterobiasis in 24a, 43b, 147a, 275b, 906c,
                                                                                                          820c, 820e, 829b, 829c, 836a, 837a, 862a,
         – & poliomyelitis in 269a.
                                                                                                          863b, 882a, 918a, 933b, 947a, 951a, 951b,
   -, Enterobius in 292a, 514d, 672e, 785c, 822a.
                                                                                                          952a, 952b, 953d, 954a, 959a, 959b, 959c,
 -, - & appendicitis in 112a, 609a, 863a.
                                                                                                          959d, 960b.
   -, - - peritonitis in 683a.
                                                                                                     -, Hymenolepis diminuta in 190g, 344a.
    -, - vermicularis in 20bf, 34c, 293a, 578c,
                                                                                                     -, — nana ova in 680a.
                                                                                                     -, Karlsbad salt 857a.
         708a, 765b, 785e.
       Fasciola hepatica in 426a, 441a, 497c,
                                                                                                     -, Loa loa in 328d.
                                                                                                    -, loaiasis in 289c.
 -, filariasis in 7b, 54b, 58g, 152a, 180a, 180d,
                                                                                                  -, magnesium sulphate 573c.
   328c, 328f, 358c, 362b, 586v, 654a, 791a, 817a, 877, 983b.

-, — & adenolymphocele in 925b.
                                                                                                  -, Meloidogyne ova in 809a.
                                                                                                   -, - sp. in 399b.
                                                                                                    -, microfilariae in 38m.
                                                                                                  -, miracil-D 69d, 235b, 289d, 328b, 801b.
   -, — — chyluria in 448a.
 -, - elephantiasis in 789a.
                                                                                                   -, nematode larva in 183a.
 -, - - eosinophilia in 949a.

-, - bancrofti in 7a, 176a, 445a, 728c, 800a,

800b, 800d, 816a, 861b, 941a.

-, - - & lymphangiectasis in 768c.
                                                                                                    -, nematodes in 578b.
                                                                                                    -, - & dermatology in 898a.
                                                                                                            - - sarcoidosis in 184a, 184b.
                                                                                                  -, Nematolyt 764a.
          - malayi in 91a.
                                                                                                  -, nilodin 586n, 586o, 586r, 624c, 625a.
   -, fouadin 170a.
                                                                                                  -, notezine 647b.
   -, gammexane 687a.
                                                                                                     -, Onchocerca volvulus in 362a, 573b, 586u.
```

Man, onchocerciasis in 68a, 243a, 367f, 892a, 893a. -, - & trachoma in 334a. —, Opisthorchis felineus in 906e. —, oxyurids in 104a. -, paragonimiasis in 289e, 425b, 639a, 696g, 706a, 831a. -, Paragonimus westermanii in 168b, 704a. -, parasites in 844a, 996. -, pheniodol 612c. -, phenosafranine compounds 357a. -, phenothiazine 340b, 573a. -, piperazine diphenylacetate 526a. -, potassium antimony tartrate 38j. -, Schistosoma haematobium in 182a, 367c. -, — japonicum in 13d, 552c, 707a. -, — & "kabure" in 291b. -, — mansoni in 176d, 391a, 586k, 783i, 798a, 799f, 801c, 948a. -, - - & splenomegaly in 452a. , schistosome appendicitis in 503a. - dermatitis in 381, 101f, 110b, 230df, 338b, 375a, 646a, 746a. -, schistosomiasis in 6a, 20b, 58i, 90a, 175a, 212a, 255a, 328e, 330a, 334c, 429a, 430b, 430d, 430e, 430f, 430g, 586a, 614a, 615a, 650a, 728b, 739a, 768e, 801a, 801d, 801e, 805b, 826b, 872. -, — & appendicitis in 648a. -, — cappendichs in 647a, 879a.
-, — naematobia in 69c, 105a, 105c, 227a, 229a, 26oc, 362c, 586j, 586m, 606a, 689a.
-, — japonica & cancer in 758a.
-, — mansoni in 250a, 334f, 355a, 367e, 522a, 585a, 586 l, 680f, 680g, 680h, 799e, 801g, 809b, 874, 977. 8095, 874, 977. -, Setaria digitata in 210c. -, sodium antimony tartrate 38j, 85a, 382a, 586q, 801f. -, -- tri-gluconate 282a. -, solustibosan 219b. -, Sparganum in 230bt. _, — mansoni in 20c. —, stibanose 278a. —, Strongyloides in 322b, 729a. -, - & creeping eruption in 19a. -, - stercoralis in 680c, 906d. -, strongyloidiasis in 350a, 691a. -, Supatonin 409a. -, Syngamus laryngeus in 783c. -, Taenia in 950a. -, - saginata in 152k, 165a, 237a, 811d, 829d. —, taeniasis in 497b, 788a. —, Ternidens deminutus in 334b. —, terramycin 467a.
—, tetrachlorethylene 423d. -, Tetrapetalonema berghei in 152i.
-, toxicity of chenopodium oil to 279a.
-, - Systox to 479a. -, Toxocara canis & visceral larva migrans in -, Trichinella spiralis in 230k. -, trichinelliasis in 108b, 211a, 378a, 459a, 466a, 467b, 467d, 481a, 482b, 485a, 504a, 549a, 581a, 582a, 653a, 705a, 785a, 785b, 890a, 914a, 934a. —, Trichostrongylidae in 190d.

Man, trichostrongylosis in 420a. -. Trichostrongylus in 940a. -, trichuriasis in 906b. -, Trichuris trichiura in 217a, 270d, 289a, 426b. -, triphenylmethane dyes 547a. —, Vermalin 718a. —, Vermella 735c. -, Wuchereria bancrofti in 2b, 34b, 318a, 367b, 367d, 586t, 783m, 800g. —, — — & elephantiasis in 728a. -, - malayi in 409a. Mangold, Ditylenchus dipsaci in 123a. Mardonius piceus, Thelastoma pteroton in **361a.** Maritrema sp. of Afanasev, 1941 to M. afanassjewi n.sp. 1000. - afanassjewi n.sp. 1000. - magnicirrus n.sp. in Upupa epops 1000. Marshallagia marshalli in goat in India 29e. - for M. orientalis 29e. - orientalis to M. marshalli 29e. Marsipobdella sacculata n.g., n.sp. on fish 749a. Marsupioacetabulum marinum n.g., n.sp. in Gerres punctatus 568a. Marsupiobothrium alopias n.g., n.sp. in Alopias vulpinus 49a. Martes spp., Ascaris n.sp. in 132e. — caurina origenes, Crenosoma n.sp. in 101c.
— flavigula, Soboliphyme baturini in 961h.
— zibellina, Sobolevingylus n.g., n.sp. in 961bf. — -, Taenia n.sp. in 961bf. Mastophorus for Protospirura 230bk Mauritius, Eurytrema pancreaticum 234b. Mediorhynchus for Heterophus 784b. passerus n.sp. in Passer domesticus 784b. Mediterranean, cestodes in fish 852c. Megapera, life-history 230de. Meggittiella to Dilepidinae 961g. Melichthys buniva, Contracaecum n.sp. in 491a. ——, Cucullanus n.sp. in 491a.
Meloidogyne, see also Heterodera marioni. - in cucumber & tomato, Systox 257b, 479a. -, diagnosis emended 132i. - in grape, resistant varieties 75a. __ lespedeza, resistant strain 483e. - ova in man 809a. - in peach, resistant varieties 75a. -- plants 910a. — —, control 747a, 913a. _ — —, resistance 257m. - - in Southern Rhodesia, control 509a. - --, susceptibility tested 771b. — — in Turkey 623a. — — — Venezuela 480a. --- soil, D-D 354a. - soya bean, resistance 636b. _ - - -, resistant varieties 636a. - sugar-beet, control by crop rotation 771a. -, technique for culturing 230bb. - in tomato, control 910b. ———, resistant lines 770a. - trees & vines, D-D 75a. - sp. in man 399b. - spp. in Antirrhinum & Lycopersicum, compared 478b. — — — majus, experimental 478a. — — — —, life-history 30 l. — —, control 31b.

Meloidogyne spp. in crops, resistance 30k. — — legumes 126b. esculentum, life-history - Lycopersicum 30 1. - peach, resistance 128a. - - South Africa 138b. -, technique for identifying 126d. - in tobacco, resistance 126e. - arenaria thamesi n.subsp. in Boehmeria utilis - hapla in peanut, control 125c, 483b. - U.S.A. 479e. - incognita, effect of potassium on reproduction - - in Hibiscus cannabinus 126f. - lima bean, effect of potassium 257a, 261e. - - Lycopersicum, resistant crosses 242a. - plants, control by crop rotation & flooding 776h. - - vine, resistance 401a. - war. acrita in Asparagus officinalis, resistance - cantaloupe 30e. Mephitis sp., Oochoristica n.sp. in 320c. Meriones libycus, Dipatelonema n.sp. in 58f. -, - blanci in 79c. Mermithid larvae in Odontria zealandica 744a. - in silkworm 916a. Mermithids in insects, oecology 852a. - mosquito larvae 20z. - - Russia 961bk. Mermis nigrescens in earwig 842a. Mesacanthion hirsutum n.sp. 610d. Mesocricetus auratus, Trichinella spiralis in 883a. Mesodon thyroidus, 1st & 2nd intermediary for Brachylaemus virginiana 323a. Mesogyna hepatica n.g., n.sp. in Vulpes macrotis 529e. Mesometra orbicularis in Box salpa, morphology 610b. Mesomphis cupreus, 2nd intermediary for Brachylaemus virginiana 323a. Metacercariae, technique for inducing excystment - in fish, distribution in body 696d. -- Xenopus laevis, destroying neuromast organs 488b. Metadesmolaimus aversivulva n.sp. 610d. Metastrongylus, see also Lungworms. — in pig 889b.

Metathelazia reviewed 132k. - for Osleroides of Skryabin 132k. - exilis n.sp. in Herpestes cafra 58j. Mexico, see also North America. —, Capillaria hepatica 588e. -, cysticerciasis 632b. -, eelworms in pineapple 387a. -, Macrobdella decora 588d. —, Onchocerca cervicalis 243d. —, onchocerciasis 243c, 450a, 450c. Microbilharzia chapini, life-history 101f. Microbothriorhynchus coelorhynchi n.g., n.sp. in Coelorhynchus sp. 49a. Microbothrium tolloi n.sp. on Mustelus 443a. Microcentrum, Chordodes n.sp. in 565a.

Microphallus bariurus n.comb. for Monocoecuma bariurum 1000. Micropterus dolomieu, Rhipidocotyle septpapillatad in 529a. Micropus affinis, Neoangularia n.g., n.sp. in 688a... — —, Neoliga n.g., n.sp. in 688a. -, Notopentorchis n.sp. in 688a. ---, Vitta n.sp. in 688a. Microstrongylea n.tribe 1002. Microtus sp., Heligmodentostoma n.g. in 1002. - spp., Rictularia n.sp. in 461a. - pennsylvanicus drummondii, Hymenolepis n.sp. . in 21b. Milvus migrans, Paricterotaenia n.sp. in 688a. Mink, helminths in 678c. -, phenothiazine 917a. Molluscicides, chemicals tested 176c, 428q. -, copper sulphate 680e. —, dinitro-o-cyclohexylphenol 229b. —, "resin soap" 800e. —, saponins 783j. —, sodium pentachlorophenate 176f, 680e, 783e.
—, various 20h, 875. Molluscs, cercariae in 230cc. , classification & identification 74b. Monhystera somereni n.sp. 342a. Moniezia expansa, life-history in Britain 165b. —, nitrogen metabolism 961bs. Monkey, schistosomiasis in 833b. Monocoecum bariurum to Microphallus bariurus n.comb. 1000. Monodhelmis arii n.sp. in Arius sp. 568a. Monodiscus macbridei n.sp. in prawn in Ceylon Monoecocestus spp., development in oribatid mites 20y. -, effect of temperature on development 87b. - sigmodontis, life-history 230h. Monogenea from fish in Norway, monographed 532a. on fish in U.S.A., new host records 541a. - fresh-water fish in U.S.A. 713g. Monorygma megacotyla n.sp. in Cephaloscyllium umbratile 49a. Montifringilla alpicola, Brachylecithum n.sp. in 1001. Morocco, Gongylonema pulchrum 645a. Morphology, Acanthocephala 87e. -, Ancylostoma braziliense & A. ceylanicum 642. -, - longespiculatum 895a. -, ascarids 9672.

Microfilaria, see also Acanthocheilonema, Dipetalo

Onchocerca, Wuchereria.

—, cause of periodicity 69b. — in man in Tanganyika 38m.

- - skunk, invasion of brain 95a.

uptake by Culicoides spp. 152e.
in vertebrates in Colombia 50b.

Microlaimus crassiceps n.sp. 610d.

— microseta n.sp. 610d.

— monstrosus n.sp. 610d.

—, periodicity 152g, 154d. — in rodents in U.S.A. 20u.

- undulatus n.sp. 610d.

— in Culicoides pifanoi 816b. Microfilariae in birds in U.S.A. 390a.

nema, Dirofilaria, Filaria, Litomosoides, Load

Mouse, ascarids in 98d. Morphology, Ascaris lumbricoides 152f.

—, — of man & pig 463b. -, fouadin 230cp. -, Rictularia n.sp. in 961i. -, Ascarophis morrhuae 962a. -, Schistosoma mansoni in 2d, 4a, 13b. - Australorbis glabratus 783a, 783d, 783f. _, - rodhaini in 596a. , bivesiculid cercariae 230cb. -, schistosomes in 230ct. -, Brachylaima virginianum metacercaria 693a. -, Syphacia obvelata in 52a. -, Capsala spp. 784a. -, Trichinella spiralis in 225a, 225b, 230t, 230bc, 230bd, 230be, 230bh, 419a, 775a, 961d. - Cercaria columbellae 327a, 368a. , - nipponensis 410b. - - & Ancylostoma caninum in 230bf, - okabei 410e. 230bg, 419b. Mozambique, helminths in man 586c, 586d, - sturniae 696a. — takahashii 410e. 586j. - yoshidae 696j. -, schistosomiasis 586n, 586o. , cercariae 610a. - haematobia 586j. -, Coenurus glomeratus 339b. Mule, Setaria equina in 963d. -, Contracaecum histiophori 491a. Multicalyx cristatus for Stichocotyle (Multicalyx) -, Cysticercus fasciolaris 389a. cristata 1000. -, cystophorous cercaria 230cd. Multiceps, see also Coenurus, Taenia. -, Diphyllobothrium latum 586w. endothoracicus in Vulpes vulpes, life-history -, Diplotriaena 796a. -, Dirofilaria ursi 530a. Muridae, Hymenolepis diminuta in 344a. -, Echinococcus granulosus 58b. Murielea n.tribe 1002. -, Fasciola hepatica 33c. Murshidiea n.tribe 1002. Musa spp., eelworms & "rayadilla" in 306a. -, Gynaecotyla adunca 230c. Musca amica, intermediary for Thelazia skrjabini -, Haemopis sanguisuga 675h. -, Haplorchis pearsoni 78a. 213b, 536c. -, Hedruris mucronifer 572b. Muscisaxicola maculirostris, Diplotriaena n.sp. in -, Heterodera cysts 164g. 572a. -, - spp. 164f. Musculium sp., Cercaria n.sp. in 230ce. -, heterophyid cercariae 230cc. Mushroom, eelworms in 30b. -, Hymenolepis spp. 39a. Muskrat, cestode in 159b. Mustela frenata effera, Skrjabingylus nasicola in -, - diminuta 208b. -, - horrida 428u. 20bg. -, Meloidogyne 132i. vison, Euryhelmis n.sp. in 428i. -, Mesometra orbicularis 610b. Mustelus edulus, Microbothrium n.sp. on 443a. -, Nordosttrema messjatzevi 675e. - manazo, Nybelinia n.sp. in 49a. -, Onchocerca spp. microfilariae 2c. -, Platybothrium n.sp. in 49a. Mya arenaria, Cercaria n.sp. in 20bc. -, Paramphistomum spp. 432a. Myiodynastes solitarius, Hamatospiculum n.sp. in - hiberniae 97h. -, Phyllodistomum lysteri 101e. -, Raillietina spp. cysticercoids 864f. Myolaimus stammeri n.sp. 867a. -, Schistorophus longicornis 931a. Myotis lucifugus, Longibucca lasiura in 261k. -, Physocephalus sexalatus larvae in 4280. - skrjabini 961s. -, Schistosoma intercalatum cercaria 328g. Myriapods, nematodes in 58d. Mytilus edulis, 1st intermediary for Prosorhynchus squamatus 213d. -, - japonicum miracidium 603a. -, - mansoni 740c. - cercariae 38d. -, Strongylus vulgaris 551a. -, Taenia saginata 765d. Narcissus, Ditylenchus dipsaci in 141b. - taeniaeformis 29d. -, eelworms in 364a. -, Tetrameres australis 841b. Narke japonica, Acanthobothrium n.sp. in 49a. -, Thelandros alatus 20i. Natica clausa, Cercaria n.sp. in 675i. Natrix sipedon, Prohemistomum n.sp. in 230f. -, Thysanosoma actinioides 132f. -, Trichobilharzia szidati 159a. Nematicides (plant eelworm) 897a. -, Triodontophorus 630a. - -, acridine dyes 464a. —, Vigisolepis spinulosa 329a. —, Wuchereria bancrofti 152b. ---, Aralo 828a. -, Alalo ozoa.
-, chlorobromopropene 75b, 126g.
-, chloropicrin 776f, 776g, 919b.
-, D-D 3a, 9a, 30h, 42a, 75a, 75b, 126g, 202a, 257n, 331a, 354a, 431a, 776c, 776f, 776g, 779a. , Zygocotyle lunatum 230cg. Moschus moschiferus, Dicrocoelium n.sp. in 1001. Mosquito larvae, mermithids in 20z. Mosquitoes, filariae in 248a. - tainting tobacco 463e. -, helminths in 34a. Mouse, antibiotics 98a, 98c. --, E. 605 forte 397a. --, ethylene dibromide 30i, 202a, 257 l, 331a, 776c, 776f, 776g.
--, methyl bromide 336a, 341a, 431a. -, arsenical compound 134a. -, N-(p-arsenoso-benzyl)-glycine amide hydrochloride 385d.

-, ascarid larvae in 230ca.

Nematicides (plant eelworm), parathion 31b, 96a, 127b, 506a, 517a, 661a. — —, Pestox III 127b. - -, phosphate insecticides 341a. - - , research reviewed 792a. - -, reviewed 257s. - -- , sodium selenate 96a. — —, soil fumigants 311a, 776e, 919a. — —, Systox 127b, 257b, 365a, 479a. - -- , technique for assaying 257p. ---, -- injecting 253a, 469a. ---, -- screening 563a. - -, tung-nut meal 31b. --, various 26a, 30c, 125b, 125c, 164d, 164j, 257e, 257h, 475b, 483b, 776d. Nematoda, free-living & plant-parasitic, see Eelworms. -, "left inversion" in development 848a. Nematode larva in eye of man 183a. Nematodes in birds 961y. -- in Denmark, new host records & synonymy 669a. - - cattle, phenothiazine 230cj, 230cv. — — —, popular account 723a. — — in Venezuela 813b. - & dermatology in man 898a. — in dog, chenopodium oil 528b.
— — —, phenothiazine 230ck.
— — —, Terit 528e. - domestic animals, phenothiazine 963c. - & gastro-enteritis in cattle 824a. -, histotropic phase 165d. — in insects, phylogeny 848b.
— mammals in Africa 586x. — man, Nematolyt 764a.
— —, technique for detecting 578b. — — myriapods 58d. -, new, in animals in Argentina 572a. -, -, - marine fish 564a. -, nitrogen catabolism 66a. - from Periplaneta, bacteria on cuticle 230bz. - in pheasant, bionomics 843b. - pig, hexylresorcinol 649b. ———, sodium fluoride 545a. ——— in U.S.A. 206a. ruminants, popular account 173a. - & sarcoidosis in man 184a, 184b. - in sheep, effect of glucoside 264a. ----, egg-count variations 97c. — — & goat, control 577a. — in Norway 165e. Nematodirus, see also Trichostrongyles. - in sheep, phenothiazine 230q. Neoangularia ababili n.g., n.sp. in Micropus affinis 688a. Neobythites macrops, Nybelima n.sp. in 49a. Neocaridina denticulata, 2nd intermediary for Plagiorchis muris 696c. Neochasmea n.tribe 1000. Neochordodes europaeus n.g., n.sp. 158a.

Neodiplogasterinae n.subf. 961bi.

carpio 20f.

in 20ba.

Neoechinorhynchus prolixus n.sp. in Carpiodes

Neofiber alleni nigrescens, Paranoplocephala n.sp.

---- Quinqueserialis n.sp. in 20ba.

Neoleucochloridium, key to spp. 320e. - problematicum, early development 37a. Neoliga diplacantha n.g., n.sp. in Micropus affinis 688a. Neophron percnopterus, Aporina n.sp. in 688a. Neotoma floridana, Böhmiella wilsoni in 320d. -, Longistriata n.sp. in 320d. --- , Taenia taeniaeformis in 320d. - -, Trichuris muris in 320d. Nettion crecca, Hymenolepis n.sp. in 688a. Neurofilaria cornellensis n.g., n.sp. in sheep in Ú.S.A. 81b. Neurofilariasis, see also Filariasis, cerebrospinal, Lumbar paralysis. in sheep 81a. New Caledonia, filariasis bancrofti 728c, 941a. Nigeria, see also West Africa. -, Cyclops n.spp. 488c. , Dracunculus intermediaries 544b. Nippostrongylus muris, water balance regulation Nochtiinae n.subf. 1002. Nomenclature 8a. -, Acanthocephala 132a. -, Hirudinea 588d. -, Stephanurus dentatus 369a, 369b, 369c, 369d. -, zoological 140a. Nordosttrema messjatzevi, morphology 675e. North America, see also America, Canada, Mexico, U.S.A. -, Acanthocephala 230bv. - -, schistosome dermatitis 88a. - -, Taenia crassiceps 428e. Norway, Monogenea 532a. -, nematodes in sheep 165e. Notemigonus crysoleucas, Octomacrum n.sp. in 55b. Notopentorchis micropus n.sp. in Micropus affinis 688a. Numeniotrema musculosa n.g., n.sp. in Numenius phaeopus 1000. Numenius americanus, Dictymetra n.g., n.spp. in 132d. — arquata, Schistorophus longicornis in 931a. — phaeopus, Numeniotrema n.g., n.sp. in 1000. Nutria, Trichinella spiralis in 211a. Nybelinia anguillicola n.sp. in Anguilla japonica - manazo n.sp. in Mustelus manazo 49a. - nipponica n.sp. in Neobythites macrops 49a. - (Syngenes) sphyrnae n.sp. in Sphyrna zygaena Nygolaimus plan-posae n.sp. 384a.

Oats, Ditylenchus dipsaci in 533b. -, eelworms in 712a. -, Heterodera avenae on 476b. , — major in 533b. Ochetosoma lampropelti n.sp. 981.
— rauschi n.sp. 981. - salmonicola Dollfus, 1951 to Pseudochetosoma n.g. 339a. Ochetosomatidae, hatchability of ova 713e. - reviewed 981. Octomacrum microconfibula n.sp. in Notemigonus

crysoleucas 55b.

Odagmia ornata, intermediary for Onchocerca gutturosa 157a.

Odontria zealandica, mermithid larvae in 744a.

Oecology, helminths 961bm.

, mermithids in insects 852a.

Oedemia fusca, Spelotrema n.sp. in 1000. Oesophagostomum & Hyostrongylus in pig, vitamin B₁₂ & aureomycin 230cl.

dentatum in pig, phenothiazine 113a.

- radiatum in cattle, phenothiazine 428r.

- stephanostomum in gorilla 200b. ___ _ & chimpanzee 200c.

- venulosum in sheep & goat, experimental 20e. Ogcocephalus radiatus, Contracaecum n.sp. in 491a. Oigolaimella n.g. for Diplogaster spp. 961bi.

Oligorchis, key to spp. 230g.
— nonarmatus n.sp. in Tamiasciuris douglasii

Olive, Tylenchulus semi-penetrans in 30n, 664a.

Ollulanidae n.fam. 1002.

Olssoniella chivosca to Brachylecithum chivosca n.comb. 1001.

Onchocerca, see also Filaria, Microfilaria.

, effect of hetrazan on adults & microfilariae 69a, 731a.

spp., differentiation of microfilariae by caudal nuclei 2c.

-, life-history 50a.

- cervicalis in horse in Mexico 243d. - microfilariae in eye of horse 243e.

- & O. reticulata identical 208a.

gutturosa in cattle in Austria, life-history 157a.
reticulata & O. cervicalis morphologically identical 50a.

- periodic ophthalmia in horse 832a. - volvulus in Guatemala, transmission 230d.

- - man 573b.

- - in Belgian Congo 586u. ___ _ French West Africa 362a.

- microfilariae, physiology 450b.

Onchocerciasis in Guatemala, new intermediaries 176g.

- man 892a.

_ _ in America 243a.

-----, antrypol 152m. - - in French West Africa 7b.

— — —, hetrazan 450d.

_ __, ocular lesions 367f.

---, pathology & treatment 893a.

- Mexico, control 450c. ---, distribution 450a.

___ & Guatemala, distribution 243c.

, ocular, in man, treatment 68a. & trachoma in man 334a.

Oncomelania formosana resistant to Schistosoma japonicum from Japan 428j.

- hupensis, Cercaria n.sp. in 696k.

- nosophora, chemicals tested against 176c. -, molluscicides tested against 428q.

Ondatra, helminths in 553a.

- zibethica, helminths in 201, 554a. -- Hymenolepis n.sp. in 428d.

. -, Taenia taeniaeformis in 854c.

Onion, Ditylenchus dipsaci in 886a. Oochoristica pedunculata n.sp. in Mephitis sp.

wallacei n.sp. in Spilogale interruptus 320c.

Opecoelus piriformis n.sp. in Lates calcarifer 5682. Opegaster gobii n.sp. in Gobius sp. 568a.

longivesicula n.sp. in Gobius sp. 568a. Opisthomonorchiinae n.subf. 568a.

Opisthomonorchis carangis n.g., n.sp. in Caranx sp.

Opisthorchis felineus in cat in Turkey 963b.

— — man 906e.

- metacercariae in fish 553d.

- guayaquilensis to Amphimerus n.comb. 588c. Orange, see Citrus.

Oribatid mites, intermediaries for Monoecocestus spp. 20y.

Ornithobilharzia dattai n.sp. in cattle 29f.

- -, life-history 29f.

- turkestanicum in sheep in Pakistan 773b.

Oryzomys palustris, Schistosoma mansoni in 20i. Osleroides of Orlov, Davtian & Liubimov to Vogeloides Skryabin 132k.

- Skryabin to Metathelazia 132k.

Ostertagia, see also Stomach worms, Trichostrongyles.

- ostertagi in cattle 187b.

(Skrjabinagia) spp. to Skrjabinagia 1002. Ostrea gigas, Gymnophalloides tokiensis in 714a. Oswaldocruzia filiformis of Travassos, 1917 to O. goezei nom.nov. 1002.

goezei nom.nov. for O. filiformis of Travassos,

1917 1002.

Oswaldocruziinae n.subf. 1002. Ovis aries, Skrjabinagia n.sp. in 1002.

- canadensis, helminths in 132g.

Oxyurids in man & domestic animals 104a. Oxyuris equi, technique for detecting 880a.

Pakistan, Ornithobilharzia turkestanicum 773b.

-, Schistosoma spp. 773c.

Panagrolaimus thienemanni n.sp. 867a. Panama, trematodes in fish 588b.

Pancreatrema disacetabulum n.g., n.sp. in Eurystomus orientalis 1001.

Panopistus europaeus n.sp. in Sorex araneus 329a. Paracanthocheilus rajae n.g., n.sp. in Raja 851a. Paracryptogonimea n.tribe 1000.

Paracryptogonimus ovatus n.sp. in fish 568a. Paradiplogaster n.g. for Diplogaster similis 961bi. Paragonimiasis, cerebral, in man 425b, 706a.

- in man in Cameroons 289e.

- ---, diagnosis 639a.

, emetine hydrochloride & sulpha drugs 696i.

-- in Japan 696g, 831a.

Paragonimus westermanii, incidence in 2nd intermediaries 696 l.

— in man 704a.

____, invasion of brain 168b.

Paragordius varius in Venezuela, first record 565a. Paraheterophyes pirum to Spelotrema pirum n.comb. 1000.

Paralichthys adspersus, Entobdella n.sp. on 443a. Paramonostomum brantae n.sp. in Branta 428a. Paramphistomes in cattle, egg-count variations

- ruminants in Ceylon, survey 78e. Paramphistomiasis in sheep, hexachlorethane 773a.

Paramphistomum spp., lymph system 432a. cervi in cattle 245e. - hibernias miracidium, development & morphology 97h. Paranoplocephala lemmi n.sp. in Lemmus spp. 428e. neofibrinus n.sp. in Neofiber alleni nigrescens 20ba. – threlkeldi n.sp. in Lagidium peruanum **713c.** Parascaris in equines, treatment 965a. - equorum, bactericidal action of extracts 604a. - in horse 702d. ————, carbon tetrachloride 850f. ———, toluene 177a. - - ova, cytoplasmic movements 489a. ---, factors affecting development 880b. - , ovum cytoplasm 47a. - -, physiology 864c. -, tetrad formation 71a. Parasites in animals in Germany, 1007, 1010. - — —, transmission 976. - beetles 983a. - — cattle 702c. - & disease in animals 986, 987. _ - - - sheep 994. - in domestic animals 971. - — fish 972. - goat, environmental factors 993. - Lynx rufus 854b. - - man 844a. - - & domestic animals (text-book) 996. - pig 756a. — poultry 927b. Parasitic enteritis in sheep in Italy 597a. Parasitism (text-book) 298. Parasitology, bibliography 560, 561, 562. -, clinical (text-book) 160. -, evolution 215a. — in Malaya 131a. -, teaching of 155g. — (text-book) 997. Parastrigea caballeroi n.sp. for P. cincta of Caballero & Vogelsang, 1949 651a. cincta of Caballero & Vogelsang, 1949 to P. caballeroi n.sp. 651a. Paratylenchus hamatus in celery 30g. -, causing stunting 478c. Pareudiplogaster n.g. for Diplogaster striatulus 961bi. Paricterotaenia milvi n.sp. in Milvus migrans 688a. - tringae to Choanotaenia burti nom.nov. 33a. Paroaria dominicana, Diplotriaena in **796a.** Paroigolaimella n.g. for Diplogaster coprophagus de Man, 1876 961bi. - anchicoprophaga nom.nov. for Diplogaster coprophagus of Goodey, 1929 **961bi.** is palustris, Brachylecithum atten attenuatum n.subsp. in 1001. Passalurus in rabbit, crystal violet 617c. Passer domesticus, Mediorhynchus n.sp. in 784b. Patagifer bilobatus, life-history 675b. Pathology, ancylostomiasis 260a. —, Aphelenchoides ritzema-bosi 258a. -, Ascaris 440a. -, cestodes 571a. -, filariasis bancrofti 445a.

-, Gongylonema neoplasticum 427a.

Pathology, Gymnophalloides tokiensis 714a. -, onchocerciasis 893a. -, Schistosoma spp. & Schistosomatium douthitti 1a. -, - bovis 38i. -, - mansoni 2d, 4a, 783g. -, - rodhaini 586s. -, schistosomiasis 430b, 430d, 430e. —, — mansoni 874, 977. -, Strongyloides 666a. -, Strongylus equinus 118c. Peach, Meloidogyne in 75a. _, _ spp. in 128a. Peanut, Belonolaimus gracilis on 257 1. -, Meloidogyne hapla in 125c, 483b. Pericrocotus roseus, Unilaterilecithum n.g., n.sp. in 1001. Peru, Heterodera rostochiensis 351a, 479b. Petalodistomum (Petalodistomum) n.subg. 1001. - (Staphylorchis) n.subg. 1001. Petasiger chandleri n.sp. in Podilymbus podiceps 230di. Petroviprocta vigissi Schachtachtinskaja, 1951 & Avioserpens galliardi Chabaud & Campana, 1949, identical 58 L Petrowospirura lynxi n.g., n.sp. in Lynx lynx 961be. Phaneropsolus philanderi n.sp. in Philander laniger 436b. Pharyngodon mearnsi n.sp. in Streptosaurus mearnsi 281e. Pharyngostrongylea n.tribe 1002. Pheasant, Heterakis gallinae in 118b.

—, nematodes in 843b. Philander laniger, Phaneropsolus n.sp. in 436b. Philippines, schistosomiasis 429a. Philometra opsalichthydis in Hypomesus olidus 695a. Phoreiobothrium validated 651d. Phyllobothrium filiforme n.sp. in Alopias oulpinus - loculatum n.sp. in Heterodontus zebra 49a. - serratum n.sp. in Triacis scyllium 49a. - squali n.sp. in Squalus sucklii 49a. - triacis n.sp. in Triacis scyllium 49a. Phyllodistomum folium in stickleback 120g. lysteri redescribed 101e. Phylogeny, free-living eelworms 848b. , nematodes in insects 848b. Physa spp., 1st intermediaries for Stomatrema guberleti 981. gyrina, intermediary for Gigantobilharzia huronensis 991. Physalis, eelworms in 483a. Physaloptera hispida larvae in Blattella germanica in Sigmodon hispidus, life-history 428h. Physiology, Ancylostoma caninum 291a, 385c. —, — duodenale extract 635a. -, — — larvae 753d. -, Ascaridia galli 13e, 150b. -, Ascaris lumbricoides 79a, 79b, 79d, 230cw. 6218, 864b, 864c, 961bs, 975. -, — ova 87c, 213c. -, — var. suum 18a, 18b. -, cestodes 864d, 864e. -, Clonorchis sinensis 915a.

Physiology, Conispiculum guindiensis 120h. -, Dicrocoelium dendriticum 915a. -, Ditylenchus spp. 559. -, - dipsaci 141b. -, Eurytrema spp. 915a. -, Fasciola hepatica 385a, 702h. -, free-living eelworms 550a. -, helminth 230co, 567a, 864g. -, - enzymes 415a. -, Heterakis gallinae 416a. -, Heterodera rostochiensis 300, 257k. -, Hirudinea 208c, 208d, 635d, 635e. -, Hymenolepis diminuta 13a, 230bs, 230bu, 385b. -, Hirudo spp. 978. -, Macracanthorhynchus hirudinaceus 428k. -, Meloidogyne incognita 257a, 261e. -, Moniezia expansa 961bs. -, Monoecocestus spp. 87b. -, nematodes 66a. -, Nippostrongylus muris 385c. Onchocerca volvulus microfilariae 450b. -, Parascaris equorum 864c. -, phenothiazine 346a. — & derivatives 371a, 371c. Protoclepsis tesselata 964b. Rhabditis belari 79e. -, Schistocephalus solidus 226a. - - plerocercoids 13g. -, Schistosoma mansoni 638b —, — — cercariae 230cq, 424b. —, Trichinella spiralis larvae 87a. , Turbatrix aceti 455a. Physocephalus sexalatus larvae in Myotis lucifugus 428o. -, life-history 961m. Physopsis africana, Cercaria n.sp. in 328g. Picus canus, Brachylecithum n.sp. in 1001. Pig, ascariasis in 240a. -, Ascaris in 900a, 953a. -, Brachylaemus erinacei in 627a. -, cysticerciasis in 632b, 626a, 807b. -, Fasciola hepatica in 245c, 659a. -, helminthiasis in 453b. -, helminths in 14a, 96b, 284a, 486a, 507a, 678a. -, hexylresorcinol 649b. -, hydatid in 245c, 631a. -, lungworms in 239a, 471a. -, Metastrongylus in 889b. -, nematodes in 206a -, Oesophagostomum & Hyostrongylus in 230cl. -, parasites in 756a. -, phenothiazine 113a, 528c. -, sodium fluoride 157b, 545a, 575a, 850a, 889a. -, Stephanurus dentatus in 649c, 905a. -, Strongyloides in 536g. -, Suscaridin 118a. -, Trichinella in 10d, 11a. -, trichinelliasis in 777a. -, Trichuris trichiura in 849b. Pigeon, helminths in 807a. -, Hymenolepis n.sp. in 33b. Pinnipedia, cestodes of 422b.

—, helminths in 961v. Pineapple, eelworms in 387a. Piranga erythromelas, Capillaria n.sp. in 236a. Pisidium sp., Cercaria n.sp. in 230cf. Pithophorus vulpeculae n.sp. in Vulpecula marina 49 Plagioporus longivesicula n.sp. in Lethrinus sp. 568u. macassarensis n.sp. in Lethrinus sp. 568a. - synagris n.sp. in Synagris sp. 568a. Plagiorchis maculosus, life-history 859b. - (Multiglandularis) megalorchis, life-history 29c. — — nom.nov. in turkeys in Britain 29c. — muris in rat, life-history 696c. Plagitura salamandra ova, effect of environment Planorbidae, control by "resin soap" 800e.

—, intermediaries for Schistosoma mansoni in Brazil 805a. Planorbis kahuika, intermediary for Calicophoron ijimai 151a. tanganyikanus transmitting Schistosoma rodhaini 327e. Plantain, eelworms in 191a. Plants, Aphelenchoides spp. in 543a. -, - fragariae in 347a. - attacked by soil nematodes 125a. — attacked by soft hematodes 125a;
—, eclworms in 26a, 30f, 57a, 117a, 126h, 127a, 174a, 203a, 216a, 256a, 257o, 313a, 313b, 376a, 388c, 392a, 446a, 480b, 505a, 511a, 533a, 558, 663a, 678b, 776b, 793a, 794a, 847a, 881a, 942a, 943a, 966a, 992. -, Heterodera in 277a. -, - rostochiensis in 351a, 476a. -, Hexatylus n.sp. in 961bj. —, Meloidogyne in 75a, 257m, 480a, 509a, 623a, 747a, 771b, 910a, 913a.

—, — hapla in 479e.
—, incognita in 776h. Platybothrium auriculatum n.sp. in Prionace glauca 49a. Platybothrium baeri nom.nov. for Cylindrophorus typicus 651d. musteli n.sp. in Mustelus manazo 49a. Platycephalus indicus, Bucephalopsis n.sp. in 568a. Platynotrema for Euparadistomum 688b. upapai n.sp. in Upupa epops 688b. Platystoma sp., Contracaecum n.sp. in 572a. Plectonchus extrematus n.sp. 192a. Plectus kenyanus n.sp. 342a. longicaudatus var. opisthocirculus n.var. 56a. submersus n.sp. 867a. Pleorchiinae n.subf. 162. Pleuronectes limanda, Agamascaris n.sp. in 851a. Pneumatophilus tracheophilus n.sp. 981. Poa sp., Anguina n.sp. in 296b. Podilymbus podiceps, Petasiger n.sp. in 230dj. -, Polymorphus trochus in 713d. Podocotyle for Podocotyloides 568a. — gracilis n.sp. in Lethrinus sp. 568a.
— serrani n.sp. in Serranus sp. 568a.
Podocotyloides to Podocotyle 568a.
Poland, helminths in man 525a. Polar bear, Trichinella spiralis in 199a. Polymorphus trochus in Podilymbus podiceps Polystoma palancai n.sp. in Hyla arborea 269b. Pomatiopsis lapidaria, potential intermediary for Schistosoma japonicum in U.S.A. 230e. Pomoxis nigromaculatus, Cleidodiscus stentor on

Porrocaecum in birds, life-history 961e.

- crassum in duck, life-history 213f, 961j. - trichiuri & P. secundum to Terranova n.comb.

Posthodiplostomum prosostomum for P. microsicya var. prosostomum 651a.

Potato, Ditylenchus destructor in 794a. -, eelworms in 169a, 301, 477a, 999.

-, Heterodera marioni in 312b.

-, - rostochiensis in 60c, 194a, 257n, 331a, 351a, 447a, 479b, 667a, 717a, 984.

Potato, sweet, see Sweet-potato. Poultry, Butynorate 129b.

—, Capillaria in 989.

-, helminths in 14a, 20be, 32a, 92a, 178a, 259a, 556, 858a, 903a, 927b. -, Syngamus trachea in 778a.

Praeorchitrema praeorchis n.g., n.sp. in Capella gallinago 1001.

Pratylenchus, see also Anguillulina.

— sp. in begonia, general account 126g. — — Helleborus miger 479c.

- - strawberry, control 75b. - spp., general account 164m.

- pratensis in sweet-potato, chloropicrin 919b. - zeae in maize in U.S.A. 760d.

Prawn, Caridinicola n.sp. in 78b.

—, Monodiscus n.sp. in 78c.

Pricea armatum n.sp. on Cybium guttatum 784c.
— melane n.sp. on Cybium guttatum 784c.

- minutum n.sp. on Cybium guttatum 784c. - robustum n.sp. on Cybium guttatum 784c.

- tetracanthum n.sp. on Cybium guttatum 784c. - tricanthum n.sp. on Cybium guttatum 784c.

Primates, helminths in 622a.

Prionace glauca, Platybothrium n.sp. in 49a. Prismatolaimus verrucosus n.sp. 867a. Pristicola n.g. for Dihemistephanus sturionis 29b.

Pristionchus for Diplogaster spp. 961bi. Pristotrema manteri n.g., n.sp. in Scaphirhynchus

platorhynchus 29b. Probolitrema (Probolitrema) n.subg. 1001.

(Reduxotrema) n.subg. 1001.

Procamacolaimus dolichostylum n.sp. 610d.

Procamallanus spp. to Spirocamallanus n.g. 491a. Prochromadorella subterranea n.sp. 610d.

Procrassiphiala halcyonae Gogate, 1940 to Subuvulifer n.g. 651a.

Procyon cancrivorus, Schistosoma mansoni in 740b. Prodesmodora terricola n.sp. 384a.

Progynotaenia longicirrata n.sp. in Lobipluvia malabarica 688a.

Prohemistomum chandleri n.sp. in Natrix sipedon, life-history 230f.

Promicrops itaiara, Heterotyphlum n.sp. in 491a. Prosogonarium arii n.g., n.sp. in Arius sp. 568a. Prosogonotrema clupeae n.sp. in Clupea clupeoides

Prosorhynchus chorinemi n.sp. in Chorinemus moadetta 568a.

- squamatus, life-history 213d.

Proteocephala n.ordo 305.

Proteocephalus pollanicola n.sp. in Coregonus pollan 408a.

- primaverus n.sp. in Salmo, life-history 428t.

Proteocephalus exiguus in Coregonus lavaretus, lifehistory 380b.

Protoclepsis tesselata, colouration 964b.

Protoschelobates seghettii, intermediary for Monoecocestus sigmodontis 230h.

Protospirura syn. of Mastophorus 230bk. Psammomermis busuluk n.sp. 961bk.

Pselionema longissimum n.sp. 610d. Pseudamphistomum truncatum metacercariae in fish 553d.

Pseudemys scripta, Cercorchis in 262b.

Pseudoalaeuris waltoni n.sp. in Xantusia riversiana

Pseudochetosoma n.g. for Ochetosoma (Pseudochetosoma) salmonicola Dollfus, 1951 339a. Pseudohapladena scatophagi n.g., n.sp. in Scatophagus arsus 568a.

Pseudoheligmosomatea n.tribe 1002.

Pseudohexabothrium rajae n.g., n.sp. on Raja fyllae 532a.

Pseudolepidapedon lethrini n.sp. in Lethrinus sp. 568a.

Pseudomaritrema postlecithale n.g., n.sp. in Tringa incana 1000.

Pseudometadena celebesensis n.g., n.sp. in Lates calcarifer 568a.

Pseudonannolene tricolor, Heth n.sp. in 58d.

-, Rondonema n.sp. in 58d. Pseudothelandros n.g. for Thelandros spp. 230bj.

Psilenchus duplexus n.sp. 132j.
— terextremus n.sp. 132j.

Psittacula eupatria, Cotugnia n.sp. in 688a. Pterobothrium chaeturichthydis n.sp. in Chaeturich-

thys hexanemus 49a.

- hira n.sp. in Ilisha elongata 49a. Pteroplatea japonica, Anthobothrium n.sp. in 49a.

Ptiloscelis resplendens, Aprocta n.sp. in 572a. Pudicea n.tribe 1002.

Puerto Rico, ancylostomiasis 350b. -, Fasciola hepatica 896a.

Pyrene mercatoria, cercaria in 230cd. Pyrrhura cruentata, Aprocta n.sp. in 495a.

Querquedula circia, Echinocotyle n.sp. in 688a. Quinquarius japonicus, Amphicotyle n.sp. in 49a. Quinqueserialis floridensis n.sp. in Neofiber alleni nigrescens 20ba.

Rabbit, antimony compounds 702a.

—, crystal violet 617c.

—, Fasciola hepatica in 593a.

-, helminths in 324b. -, hydatid in 597b.

Trichostrongylus retortaeformis in 120e. Radix auricularia, 1st intermediary for Plagiorchis

maculosus 859b. - ovata, intermediary for Trichobilharzia szidati

158b. Radopholus, see also Anguillulina.

— oryzae in rice, general account 474a.
Raillietina cesticillus, diphenyl methane com-

pounds 230cn.

- in poultry, Butynorate 129b.
- echinobothrida cysticercoid, morphology 864f. - tetragona cysticercoid, morphology 864f.

Raillietina urogalli, development 213e. -, spermatogenesis 675a. Raja sp., Branchellion n.sp. on 749a.
— batis, Paracanthocheilus n.g., n.sp. in 851a. - fyllae, Pseudohexabothrium n.g., n.sp. on 532a. - kenojei, Anthobothrium n.sp. in 49a. Rana sp., Gorgodera (Mediodera) n.sp. in 1001. - spp., Gorgodera n.sp. in 982. - cyanophlyctis, Camallanides prashadi in 821a. - pipiens, Langeronia n.sp. in 281b. - tigrina, Centrorhynchus n.sp. in 784d. Ransomnema christiei n.sp. in Rhinocricus cachoeirensis 58d. Raphidascaris lutiani n.sp. in Lutianus analis 491a. Rat, Cysticercus fasciolaris in 635b. -, Gongylonema neoplasticum in 427a. -, hetrazan 2g. -, Hymenolepis spp. in 944a. -, — diminuta in 230bs. -, Inermicapsifer cubensis in 270a, 811b. -, Plagiorchis muris in 696c. -, pumpkin seed extract 887a. -, Trichinella in 204a. Trichosomoides crassicauda in 317a. Rattus norvegicus, helminths in 99a, 139b, 808a. - rattus, Fasciola hepatica in 20s. Ratzia joyeuxi, life-history 12a, 58c.
— in Vipera lebetina 12a, 58c. Renicola brantae n.sp. in Branta canadensis 230da. Resistance, see Immunity.

Rhabdias mucronata n.sp. in Leptodactylus 572a. - truncata n.sp. in Telmatobius schreiteri 572a. Rhabditis revised, n.subgg. & n.spp. 297a.

— belari, nucleic acids during gametogenesis 337a. -, sex determination 79e. - (Caenorhabditis) n.subg. 297a. - (Choriorhabditis) n.subg. 297a. - cristata n.sp. 867a. - elaphri n.sp. 867a. - elegans chromosomes, desoxyribonucleic acid during meiosis 12b. -, cytology 62a. ——, gametogenesis demonstrated 368b. — friderici n.sp. 867a. - (Telorhabditis) hermaphrodita n.sp. 297a. — inermiformis n.sp. 297a.
— (Caenorhabditis) körneri n.sp. 297a.
— limicola n.sp. 867a.
— (Mesorhabditis) n.subg. 297a. - (Protorhabditis) n.subg. 297a. _ (Rhabditis) n.subg. 297a. - (Telorhabditis) n.subg. 297a. - (Teratorhabditis) n.subg. 297a. tristis n.sp. 867a. Rhigonema glabrum n.sp. in Rhinocricus cachoeiren-sis 58d. -longicaudatum n.sp. in Leptodesmus paulistus - subtruncatum n.sp. in Gymnostreptus sp. 58d. Rhinocricus sp., Carnoya n.sp. in 58d. -, Ichthyocephalus n.sp. in 58d. — spp., Heth n.spp. in 58d. — cachoeirensis, Ransomnema n.sp. in 58d.

-, Rhigonema n.sp. in 58d.

Rhipicephalus sp. & Ornithodorus sp., inter-

mediaries for Dipetalonema blanci 79c.

Rhipidocotyle septpapillata in Micropterus dolomieu, life-history 529a. Rhipidura leucophrys, Austrofilaria n.sp. in 841b. Rhodatrema quadrilobata in fish, life-history 213g. Rhynchobothrium uncinatum Linton, 1924 to Floriceps uncinatus n.comb. 49a. Rhynconema brevituba n.sp. 610d. longituba n.sp. 610d. Rice, Aphelenchoides besseyi in 601b. -, - oryzae in 30c, 257e, 341a, 483d, 601a. —, eelworms in 512a, 721a. , Radopholus oryzae in 474a. Richmondena cardinalis, Gigantobilharzia huro-nensis in 991. Rictularia baicalensis n.sp. in mouse 961i. - halli redescribed 261h. - microti n.sp. in Microtus spp. 461a. Rodents, ascarids in 1004. -, helminths in 428e. — in Hungary not carrying Trichinella 309a. —, microfilariae in 20u. -, Schistosoma rodhaini in 58h, 328h, 586s. 596b. Rondonema pseudonannolenou n.sp. in Pseudonannolene tricolor 58d. Rotylenchus, see also Anguillulina. coheni n.sp. on Hippeastrum 97e. Rumania, free-living eelworms 56a. Ruminants, Eurytrema pancreaticum in 234b. -, Fasciola hepatica in 542a. -, helminths in 866a. —, lumbar paralysis in 92b. —, nematodes in 173a. -, paramphistomes in 78e. , Schistosoma sp. in 773c. Russia, Capillaria 961bp -, Choanotaenia n.sp. 961bt. —, Corrigia n.sp. 961bt. —, Cysticercus bovis 285a. -, Dictyocaulus 917b. —, helminthology **553c**, 726a, 961a, 961b. —, helminths 961bm. __, __ in animals 961p. -, - - birds 961q. -, -- goose 961br -, - guinea-fowl 961n. -, - - mammals 961i. -, - - Ondatra 553a. -, Hexatylus vigissi 961bj. -, Leucochloridium n.sp. 961bt. -, mermithids 961bk. -, Soboliphyme baturini 961h. - soricis 961h. -, Trichostrongylus n.sp. 961bt. Rye, Ditylenchus dipsaci in 280a. , eelworms in 30d, 722a. Rye-grass, eelworms in 30d. Sable, phenothiazine 917a.

Sable, phenothiazine 917a.

—, Taenia sibirica in 917c.

Salmo clarkii, Proteocephalus n.sp. in 428t.

— irideus, Diphyllobchrium latum in 130a.

Salvelinus alpinus, Capillaria n.sp. in 962a.

Samoa, Wuchereria bancrofti 176b.

Santo Domingo, see Dominican Republic.

Sardinia, cercariae in Bulinus truncatus 38h.

	ott to an in Decil Foot
Sauricolinae n.subf. 1002.	Schistosoma mansoni in man in Brazil 7831, 948a.
Scabious, Aphelenchoides n.sp. in 60a.	Dominican Republic 176d, 391a.
Scapanus townsendi, Hymenolepis n.sp. in 428d. Scaphirhynchus platorhynchus, Pristotrema n.g.,	————, miracil-D 328b.
n.sp. in 29b.	— — mouse, fouadin 230cp.
Scaptrella brevicaudata n.sp. 610d.	— — —, histopathology 2d.
Scarus sp., Diplobulbus n.sp. in 568a.	————, pathology 4a .
Scatophagus arsus, Pseudohapladena n.g., n.sp. in	— — Oryzomys palustris, experimental 20j.
568a.	— — Planorbidae, susceptibility 132b.
Schistocephalus solidus, effect of temperature on	— — Procyon cancrivorus 740b.
maturation in vitro 226a.	———— spinal cord of man 798a.
— plerocercoids, physiology 13g.	— — & splenomegaly in man in Egypt 452a.
Schistorophus longicornis in Numenius arquata,	, susceptibility of Tropicorbis spp. to 13f.
redescribed 931a.	— in Tropicorbis havanensis, survival 230cr.
— skrjabini, male described 961s.	— rodhaini in Belgian Congo, new focus 327e. — — mouse 596a.
Schistosoma, history of discovery 103b. — sp. in ruminants 773c.	——— rodents in Belgian Congo 58h, 328h,
- spp. & Schistosomatium douthitti in mouse,	596b.
pathology 1a.	———, pathology 586s.
- bovis in cattle, pathology 38i.	- spindale in buffalo in Indo-China 273a.
- haematobium in Bulinus truncatus, suscepti-	Pakistan, first record 773c.
bility 230cx.	Schistosomatium douthitti in mouse, acquired
——— man 367c.	immunity 424a.
———— in Algeria 182a.	————, immunity following treatment
————, hexamethylenetetramine 720a.	230cz.
———, miracil-D 235b.	, parthenogenetic development 320b.
- intercalatum cercaria, morphology 328g.	—— in Peromyscus maniculatus, unisexual &
— japonicum in brain of man 707a.	bisexual infections 55a.
— — cercariae causing "kabure" 291b. — — —, protective ointments tested 457a.	Schistosome antigen, technique for preparing 680b.
— in Formosa, survey 2e.	- appendicitis in man 503a.
from Japan, Oncomelania formosana	- cercariae, technique for exposing mice to 20a.
resistant to 428j.	- dermatitis caused by Gercaria littorinalinae
— in man, diagnosis 552c.	35a.
————, repellent action of copper oleate 13d.	— — in man in Australia 110b.
— — miracidium, epidermal plates 603a.	————, protective ointments 38 1.
—— in Oncomelania spp., susceptibility 230cy.	———— in Switzerland 375a.
——— Pomatiopsis lapidaria, susceptibility	————— U.S.A. 230df, 338b, 646a.
230cy.	, new records 746a.
- mansoni, acetylcholinesterase demonstrated	— — North America, oecology 88a.
— in Australorbis, effect on respiration 783b.	— — U.S.A. 122a.
glabratus, bionomics 61b.	— intermediaries in Belgian Congo 328a. — —, biological control 59b.
—————, effect of oxygen 230cq.	— —, classification 925a.
, survival 230cr.	, control by mollusc-eating fish 463a.
, tissue reactions 230j.	Schistosomes in mouse, age resistance 230ct.
— — cercariae in Australorbis glabratus 454d.	, miracil-D & organic antimonials tested
———, effect of dyes 680d.	against 424c.
———, —— sera 19b.	-, technique for recovering 740a.
———, emergence from snails 97i.	Schistosomiasis in Africa 290a.
killed by ozone 20t.	— & appendicitis in man 648a.
———, mode of skin penetration 152j.	— in Belgian Congo, control 650a.
— —, penetration mechanism 38d. — —, streptococcal decapsulation 424b.	— & cancer in man 647a, 879a.
, sucprocectal decapsulation 4240.	-, control of intermediaries & larvae by ultra-
230cs.	sonic & short waves 328i.
— — in Gerbillus pyramidum 261i.	—, — by molluscicides 229b. — in man 6a, 90a, 614a, 615a, 805b, 826b.
——— hamster & mouse, experimental 13b.	, aureomycin ineffective 176h.
— —, hermaphroditism 740c.	——— in Belgian Congo 328e.
, incidence in intermediaries in Brazil	, cardiac involvement 801a.
783h.	, causing aneurysm 212a.
, intermediaries in Africa 596a.	———, control, popular account 255a.
, Belgian Congo 59c.	———, cor pulmonale 175a.
——, —— Brazil 798b, 805a.	——— in French Sudan 728b.
— in laboratory animals, histopathology 783g.	— — , general account 801e.
——— man 799f, 801c.	———, glucantime 566a.
———— in Angola 586k.	——— in Madagascar, survey 739a.

Setaria digitata & lumbar paralysis 70a, 283a. Schistosomiasis in man, miracil-D 289d. --- in sheep, goat & horse 41d. - - in Mozambique, nilodin 586n. — —, pathology 430b, 430d, 430e. — —, reviewed 586a. __ in man 210c. - equina in mule in Turkey 963d. Severianoia julicola n.sp. in Julus terrestris 58d. ---, sodium antimony tartrate 382a. Sheep, carbon tetrachloride 285b, 546b. ---, -- tri-gluconate 282a. - cestodes in 917d. __ _ in Somaliland 330a. -, coenuriasis in 536i. _____, treatment 38j, 430g, 801d. _____ in Yemen, survey 20b. -, Cystocaulus nigrescens in 618d. -, cysticerciasis in 352f. — monkey & sheep, experimental 833b. — Philippines, survey 429a. -, Cysticercus tenuicollis in 453d. -, Cystocaulus & pneumonia in 961bh. -, pulmonary, in man 768e, 872. -, Dicrocoelium dendriticum in 23a, 546a. _, __, ___, diagnosis 58i. —, Dictyocaulus in 917b. —, — filaria in 319b, 849c. - & splenomegaly, treatment 430f. — streptomycin sensitivity unrelated 103a. — in Sudan 334c. -, emetine hydrochloride 230cu. -, Fasciola sp. in 261g. -, technique for detecting 522b. -, - hepatica in 171a. - bovis in cattle in Belgian Congo 59a. -, haemonchiasis in 787a. — haematobia in Iraq, Iran & Syria 229a. — — Libya, survey 327b. —, Haemonchus & Nematodirus in 230p.

—, helminths in 10e, 83b, 142a, 155d, 155e, 165f, 209b, 356a, 374b, 878.

—, hexachlorethane 773a. __ _ man 69c, 362c, 606a. __ _ _ in Angola 586m. _ _ _ _, anthiomaline 624b. -, hydatid in 113b. —, lungworms in 288a, 751a, 772b, 990. —, Magenwurmtabletten (Marienfelde) 458a. __ _ _ __, antrypol 586p. ____, diagnosis & treatment 625a. _____, general account 260c. _____ in India 689a. -, nematodes in 165e. -, Neurofilaria n.g., n.sp. in 81b. _ _ _ _ _ Iraq 227a. , neurofilariasis in 81a. ____, miracil-D **69d.** ____ in Mozambique **586j.** -, Oesophagostomum venulosum in 20e. -, Ornithobilharzia turkestanicum in 773b. ____, nilodin 586r, 624c. , parasites & disease in 994. parasitic enteritis in **597a**. ____, sodium antimony tartrate 85a, 586q. ____, vitamin A ineffective 105c. -, phenothiazine 10a, 230q, 458a, 528a. - - Sierra Leone, epidemiology 105a. -, - photosensitization 473a. - japonica & cancer in man 758a. -, schistosomiasis in 833b. -, control of intermediaries 176f. -, Setaria digitata & lumbar paralysis in 41d. - mansoni in Brazil 503b. -, technique for collecting faeces from 67d. ____, first record in São Paulo 399d. -, Thysanosoma actinioides in 230bw. -, diagnosis by rectal biopsy 799a, 799b, -, trichostrongyles in 539a. Trichostrongylin 458a.
Trichostrongylus axei in 19c. 799c, 799d. - in man 586 1, 801g --- in Brazil 809b. Shrews, cestodes in 21a. ______, diagnosis 355a, 585a. ______, endarteritis 874. Siam, see Thailand. Sicily, Enterobius 785c. ____, haematology 680f, 680g. —, helminths in man 785d. Sierra Leone, see also West Africa. ---, -- after treatment 680h. ____, lesion in spinal cord 250a. ____ in Madagascar 367e. - -, schistosomiasis haematobia 105a. Sigmodon, Litomosoides carinii in 152c. _____, miracil-D 801b. _____ in Mozambique, nilodin 5860. -, MSb 152d. - hispidus, Litomosoides carinii in 230ba. — —, Longistriata adunca in 20r. — —, MSb 334d, 334g. __ _ _ ___, pathology 977. ____, sodium antimony tartrate 801f. ____, South Africa 522a. ____, Sudan 334f. — -, Physaloptera hispida in 428h. -, Strongyloides sigmodontis in 20r. — — & rectal polyps in man 799e. Schrankia brasili Travassos, 1927 to Schrankiana Silkworm, mermithid in 916a. Simulium spp., biting habits 243b.
——, flight range & longevity 599a, 599b. n.comb. 583a. — larvata Vaz, 1933 to Schrankiana n.comb. 583a. Schrankiana brasili n.comb. for Schrankia brasili - larvae & pupae on crabs 120f. - damnosum in Gold Coast, bionomics 289b. Travassos, 1927 583a. larvata n.comb. for Schrankia larvata Vaz, - neavei, control 38a. Siphoderea n.tribe 1000. Siren, 2nd intermediary for Stomatrema guberleti 1933 583a. Seals, cestodes in 72a. Sebastodes caurinus, Ascarophis n.sp. in 491a. Skrjabinagia for Ostertagia (Skrjabinagia) spp. 1002. Semigordionus circumannulatus n.g., n.sp. 158a. Senegal, see French West Africa. - dagestanica n.sp. in Ovis aries 1002. Serranus sp., Podocotyle n.sp. in 568a.

Skrjabinema ovis in goat in Czechoslovakia 964a. Skrjabinocercina petrowi n.g., n.sp. in Alactaga saltator 961be. Skrjabingylus nasicola in Mustela frenata 20bg. Skrjabinoclava longifuniculata n.sp. in Charadrius sp. 961ba. Skrjabinomermis sukatschevi n.sp. 961bk. · tolski n.g., n.sp. 961bk. Skrjabinus lanciformis n.sp. in Palco subbuteo - popovi n.sp. in Tetraogallus caucasicus 961u. Skunk, microfilariae in 95a. Snails, aquatic, technique for feeding 38g. Sobolevingylinae n.subf. 961bf. Sobolevingylus petrovi n.g., n.sp. in Martes zibellina 961bf. Sobolephya oshmarini in Larus sp. 1000. Soboliphyme baturini in Martes flavigula 961h. - jamesoni n.sp. in Sorex vagrans 101b. soricis in Sorex araneus in Russia 961h. Soboliphymidae emended 961h. Society Islands, filariasis bancrofti 176a. Soil, eelworms in 26a, 592a, 776a, 776c, 776d, Solanum, Heterodera rostochiensis in 463c, 518a. tuberosum subsp., Heterodera rostochiensis in 463d. Somaliland, ancylostomiasis 330a. -, Bulinus abyssinicus 514e. -, schistosomiasis 330a. Sorex sp., Mammanidula n.g., n.sp. in 1002.
— araneus, Capillaria n.spp. in 329a. - -, Leucochloridium n.sp. in 329a. — —, Panopistus n.sp. in 329a. --- , Soboliphyme soricis in 961h. --- , Synhimantus n.sp. in 329a. vagrans, Soboliphyme n.sp. in 101b.
vagrans, Hymenolepis n.spp. in 21a. South Africa, see also Africa. -, eelworms in plants 388c, 511a. – —, Gaigeria pachyscelis **434a.** ----, helminths in domestic animals 388b. — —, — — man 36a, 143a, 521a. — —, Meloidogyne spp. 138b. — —, schistosomiasis mansoni 522a. Southern Rhodesia, Meloidogyne 509a. Soya bean, Heterodera n.sp. in 254a. - -, - glycines in 265a. ---, Meloidogyne in 636a, 636b. Spain, Eucoleus hepaticus 135a. —, free-living eelworms 782a, 782b, 939a. -, helminths in dog 135a. -, - man 451a. -, trichinelliasis 504a. Sparganum in man in Korea 230bt. - mansoni in man in U.S.A. 20c. Spathebothridea n.ordo 305. Spathebothriidae n.fam. 305. Spelophallus primas to Spelotrema prima n.comb. 1000. Spelotrema oedemia n.sp. in Oedemia fusca 1000. - pirum n.comb. for Paraheterophyes pirum 1000. prima n.comb. for Spelophallus primas 1000.

Sphaerium simile, 1st intermediary for Gorgodera

vivata 982.

- duplex n.sp. 555c.

Sphaerolaimus asetosus n.sp. 555c.

Sphaeronema californicum n.g., n.sp. in Umbellularia & Arctostaphylos 261b. Sphaeronematinae n.subf. 261b. Sphyraena sp., Bucephalus n.sp. in 568a. -, Lecithochirium n.sp. in 568a. Sphyrna zygaena, Nybelinia n.sp. in 49a. Spiculopteragia alcis n.sp. in Alces alces 1002. Spilogale interruptus, Oochoristica n.sp. in 320c. Spinus tristis, Gigantobilharzia huronensis in 991. Spirocamallanus n.g. for Procamallanus spp. 491a. monotaxis n.sp. in Monotaxis 491a. Spirocerca lupi in dog 154c. - -, hetrazan 407a. – — in Indonesia 395a. -, intermediary in China 327d-Spirorchis magnitestis chromosomes 713a. Squalus sucklii, Phyllobothrium n.sp. in 49a. Squamastrongylea n.tribe 1002. Squirrel, Dicrocoelium lanceatum n.var. in 1001. Stachys palustris, Ditylenchus destructor in 60b. Staphylorchis gigas n.sp. in Galeocerdo 484a. Stegodexamene anguillae in Anguilla spp., lifehistory & bionomics 428c. Stephanochasmus caducus Looss, 1901, of Lebour, 1908 to Stephanostomum lebourae n.sp. 162: carangis Yamaguti, 1951 to Stephanostomum carangium n.comb. 162. - ceylonicus Lühe, 1906 to Stephanostomun ceylonicum n.comb. 162. - robustus MacCallum, 1917 to Stephanostomum robustum n.comb. 162. Stephanofilariasis in buffalo, treatment 92c. Stephanolaimus paraflevensis n.sp. 610d. Stephanostomum carangium n.comb. for Stephanochasmus carangis Yamaguti, 1951 162. - ceylonicum n.comb. for Stephanochasmus ceylonicus Lühe, 1906 162.
- lebourae n.sp. for Stephanochasmus caducus
Looss, 1901, of Lebour, 1908 162. robustum n.comb. for Stephanochasmus robustus MacCallum, 1917 162. Stephanurus dentatus in cattle 132 l. ----, nomenclature 369a, 369b, 369c, 369d. — in pig, general account 649c. ———— in Japan 905a. Stichocotyle (Multicalyx) cristata to Multicalyx cristatus 1000. Stickleback, trematodes in 120g. Stomach worms, see also Haemonchus, Ostertagia, Trichostrongyles, Trichostrongylus. — in sheep, treatment 458a.

Stomatrema guberleti in Farancia, life-history 981. Strawberry, Aphelenchoides & Corynebacterium in 331d. -, — fragariae in 527a. -, — ritzema-bosi in 527a. -, Ditylenchus dipsaci in 527a. -, eelworms in 825a. -, Pratylenchus sp. in 75b. Streptosaurus mearnsi, Pharyngodon n.sp. in 281e.
—, Strongyluris n.sp. in 281e.
Strigea vaginata for Apatemon gracilis of Caballero & Vogelsang, 1949 651a. Strigeida redescribed 651a. Strix rufipes, Thelazia n.sp. in 572a. uralensis, Brachylecithum lobatum n.subsp. in 1001.

Strongylata catalogued 1002. Strongyle larvae in horse 118d. - of horse, chemicals tested against 638a. - -- , n-heterocyclic compounds 319a. — —, survival in soils 536j. ova & larvae, effect of detergents 230cm, 839a. Strongyles, see also Trichonema. - in horse 998. — — —, phenothiazine, continuous therapy 40a.
— — —, treatment & control 198b. Strongyloides in cattle, pathology 666a. - & creeping eruption in man 19a. - in laboratory animals, pathogenicity & treatment 247a. - larvae, technique for collecting from faeces 399a. - in man 729a. -- - in Brazil 322b. - ---, technique for diagnosing 322a. - - pig, control 536g. - spp., prenatal infections 294a. - ratti, gametogenesis & sex determination 359a. — sigmodontis in Sigmodon hispidus 20r. - stercoralis, giant strain 327f.
- in man 680c, 906d. — — — —, gentian violet 514f, 946e. — westeri in horse 880c. Strongyloidiasis in man 350a. — —, general account 691a.
— —, hetrazan ineffective 399g. Strongylosis in horse 961bl. Strongyluris riversidensis n.sp. in Streptosaurus mearnsi 281e. Strongylus edentatus in horse, haematology 377a. — — —, prepatent period 83a. — equinus in horse, pathology 118c. — vulgaris in horse 41b. — —, morphology 551a.

— — & verminous aneurysms, reviewed 155b. Stunkardionematea n.tribe 1002.
Subulura, taxonomy 20be.
Subuvulifer n.g. for Procrassiphiala halcyonae
Gogate, 1940 651a. Sudan, see also French Sudan. -, schistosomiasis 334c. —, — mansoni 334f. Sugar-beet, Heterodera schachtii in 325a, 475a, 678d, 886a, 984. , Meloidogyne in 771a. Sus scrofa, helminths in 961bg. Swaziland, taeniasis 788a. Sweden, Ditylenchus dipsaci 533b, 886a. -, eelworms in crops 533a. -, free-living eelworms 869a. —, Heterodera major 533b. —, — schachtii 886a. Sweet-potato, eelworms in 30h. -, Heterodera marioni in 246a. -, Pratylenchus pratensis in 919b. Switzerland, eelworms in tobacco 508c. -, free-living eelworms 384a. —, helminths in animals 516a. —, — — deer 276a. -, Hirudinea 651c. -, schistosome dermatitis 375a. -, Trichinella spiralis 211a.

-, trichinelliasis 211a.

Symbothriorhynchus uranoscopi n.g., n.sp. in Uranoscopus oligolepis 49a. Symptomatology, helminths in pig 507a. -, trichinelliasis 653a. Synagris sp., Plagioporus n.sp. in 568a. Syngamus in turkeys, iodine 492a.
— arcticus n.sp. in Gavia stellata 961 1. - auris to Mammomonogamus auris n.comb. - laryngeus in man in Brazil 783c. — trachea in poultry, general account 778a. Synhimantus, see also Acuaria. - rhopalocephalus n.sp. in Sorex araneus 329a. Synodontium monhystera n.sp. 610d. Synthetocaulus rufescens in goat, phenothiazine derivatives 207a. Syphacia obvelata in mouse, anthelmintics tested against 52b, 230z, 385e. ----, arsenical compound 134a. Syria, schistosomiasis haematobia 229a. Taenia, see also Cysticercus, Multiceps. -, abnormal specimen 230b. - in man 950a. - brauni in dog, life-history 819a. - crassiceps in Dicrostonyx groenlandicus in North America 428e. crassicollis, see T. taeniaeformis.
jakhalsi, adult of Cysticercus madoquae 514c. - -, life-history 618e. - saginata in man, acridine 742a. ————, — derivatives **765a.** ———— in Africa **152k.** — — — , allergic symptoms 829d.
— — — , atebrin 94a, 210b, 637a, 933c.
— — in Denmark 165a.
— — , hexylresorcinol & tetrachlorethylene 270c. _____, location by X-ray 237a. _____, treatment 811d. ----, sphincter muscle demonstrated 765d. — —, triradiate specimen 139a. — sibirica in sable 917c. — skrjabini n.sp. in Martes zibellina 961bf. — taeniaeformis in Neotoma floridana 320d. — — Ondatra zibethica 854c. ---, spiral torsion 29d. Taeniasis in man, general account 497b.

— — in Swaziland, control 788a. Tahiti, Wuchereria bancrofti 318a. Tamiasciuris douglasii, Oligorchis n.sp. in 230g. Tanganyika, filariasis 791a. -, — & elephantiasis 789a. Tarsius carbonarius, helminths in 488a. Tarvaia angusta n.sp. 610d. Taxonomy, Hirudinea 472a.

—, Leucochloridinae 320e. Subulura 20be. Technique for administering atebrin 352e. -- D-D 42a. — — phenothiazine by drenching gun 148c. — applying soil fumigants 776e. - assaying antimony in blood 430i. __ _ compounds 430h.

```
Telmatobius schreiteri, Hedruris n.sp. in 551b,
Technique for assaying enzymatic factor of
     Schistosoma mansoni cercariae 230cs.
                                                             -, Rhabdias n.sp. in 572a.
     - nematicides 257p.
                                                         Telogaster opisthorchis in Anguilla spp., life-
    - - potato root diffusate 331b.
                                                              history & bionomics 428c.
 - — collecting faeces from sheep 67d.
                                                         Tenodera, Chordodes n.sp. in 603b.
 - - helminth ova from faeces 230by.
                                                         Tenuiproboscis sergenti n.sp. in dog 63b.
 ____ soil 843b.
 - - Heterodera cysts 97b.
                                                         Ternidens deminutus in man 334b.
                                                         Terranova ginglymostomae n.sp. in Ginglymostoma
 - - Strongyloides larvae from faeces 399a.
  - - trematode ova 97j.
                                                              cirratum 491a.
                                                           - secundum n.comb. for Porrocaecum secundum
 - concentrating helminth ova 230s.
 491a.
                                                           trichiuri n.comb. for Porrocaecum trichiuri
  - - counting & culturing Turbatrix aceti 97f.
                                                              491a.
 -- - eosinophils 430c.
                                                         Tetrameres australis, female described 841b.
                                                         Tetraodon hispidus, Johanssonia n.sp. on 749a.
 - — helminth ova 28b.
 - - culturing Ascaris ova 410a.
                                                         Tetraogallus caucasicus, Skrjabinus n.sp. in 961u.
                                                            himalayensis, Ascaridia n.sp. in 961bc.
     - - Bdellonyssus bacoti 20v.
 - — — Gynaecotyla adunca 20q.
                                                         Tetrapetalonema berghei in man in Belgian Congo
  - - Meloidogyne 230bb
                                                              152i
     - Turbatrix aceti 230bl.
                                                         Thailand, helminths in man 230bn, 230bo.
 - - detecting Enterobius in man 552a.
                                                         Thaparocleidus wallagonius n.g., n.sp. on Walla-
                                                         gonia attu 912c.
Theilerianea n.tribe 1002.
 - — Fasciola hepatica ova 17a.
 - — helminth ova in faeces 655a, 672d.
                                                         Thelandros spp. to Pseudothelandros n.g. 230bj.
— alatus, male redescribed 20i.
— for T. avis 20i.
 - - helminths 230r.
  - - Heterodera cysts 479d.
  - nematodes in man 578b.
                                                           – – – micruris 20i.
    - - Oxyuris equi 880a.
                                                             — — sahariensis 20i.

— schistosomiasis 522b.

    - determining ascaridole in chenopodium oil
                                                         — avis to T. alatus 20i.
                                                         - bicaudatus n.sp. in Xantusia riversiana 428b.
       - concentration of molluscicides 783e.
                                                         — californiensis n.sp. in Xantusia vigilis 428b.
  - diagnosing enterobiasis 36b, 605a, 846a.
                                                          - micruris to T. alatus 201.
  - - helminths 161, 399b.
                                                         — minutus n.sp. in Batrachoseps 428b.
— sahariensis to T. alatus 20i.
  - - Strongyloides in man 322a.
                                                           - waltoni n.sp. for T. sp. of Walton 428b.
 - — dissecting Heterodera cysts 97a.
 - drenching horse 67b.
                                                         Thelastoma bulhoesi n.sp. in Gymnostreptus sp.
 - - examining egg counts statistically 422a.
 - — Heterodera cysts 97a.
                                                         - galliardi n.sp. in Glomeris sp. 58d.

    – exposing mice to schistosome cercariae 20a.

                                                         - glomericola n.sp. in Glomeris annulata 58d.
 - - feeding aquatic snails 38g.
                                                         — pteroton n.sp. in Julus (sensu lato) sp. 58d.
— in Mardonius piceus 361a.
 - - fixing helminth ova 703a.
 - identifying Meloidogyne spp. 126d.
                                                         Thelazia in cattle, control 536d.

    inducing excystment of metacercariae 20q.

                                                           - bubalis n.sp. in buffalo in India 221b.
 - - infecting fowl with Ascaridia galli 230x.

    longicaudata n.sp. in Strix rufipes 572a.

— injecting soil fumigants 253a, 469a.
— inoculating plants with Heterodera 149a.
— mounting cestodes & trematodes 241a.

                                                         - rhodesii in cattle, treatment & control 849d.
                                                          - skrjabini in cattle, life-history 213b.
                                                           - -, life-history 536c.
 - - helminth larvae 74a.
                                                         Theristus curvispiculum n.sp. 610d.
                                                         - heterospiculoides n.sp. 610d.
    - trematodes 55b.
 - — preparing Ascaris lumbricoides extracts 845a. — — bacteria-free eelworms 790a.
                                                         - inermis n.sp. 610d.
                                                          - longicaudatus-viviparus n.sp. 555b.
 - - Eurytrema antigen 802a.
                                                           microspiculum n.sp. 610d.
  --- hydatid antigen 349a.
                                                         Thermocyclops nigerianus, bionomics 61a.
 - - santonin from Artemisia 570a, 570b.
                                                         Thiara granifera, intermediary for trematodes,
 - - schistosome antigen 680b.
                                                              biology 487a.
 - — rearing Pomatiopsis lapidaria 230e.
                                                         Thynnus thunnina, Capsala n.sp. on 784a.
 - — recovering schistosomes 740a.
                                                         Thysanosoma
                                                                       actinioides, abnormal specimen
 - removing hydatid 682a.
                                                              132f.
           - from brain 612a.
                                                             - in sheep, seasonal incidence 230bw.
 - — sampling for Heterodera rostochiensis 164a.
- — screening nematicides 563a.
                                                         Tobacco, eelworms in 30i, 86a, 198a, 202a, 508c,
                                                              510a, 776i.
 - -- staining & mounting helminths 146a.
                                                         -, Heterodera sp. in 126a, 257j.
 - - teaching helminthology 115a.
                                                          -, — marioni in 263a.
-, — rostochiensis in 760b, 760c.
 - testing anthelmintics 155f, 617a, 887b.
                                                          Meloidogyne spp. in 126e.

tainted by D-D 463e.
     - susceptibility of rye to Ditylenchus
    dipsaci 280a.
```

Triaenophorus lucii in Anguilla anguilla 332a. Tomato, Heterodera sp. in 126a. - nodulosus plerocercoids in Cottus cognatus -, - rostochiensis in 257i. -, Meloidogyne in 257b, 479a, 770a, 910b. Trichinella in guinea-pig 970. Tormopsolus lintoni n.sp. for Distomum sp. - - pig in Canada 10d, 11a. Linton, 1901 162. - rat in Canada 204a. Toxicity of carbon tetrachloride to goose 528d. - & Sarcocystis in meat inspection 23b. - chenopodium oil to man 279a. - spiralis & Ancylostoma caninum in mouse, cross immunity 230bf, 230bg, 419b. - — filix mas 244a. - magnesium sulphate to man 573c. - in animals in Aleutian Islands 99a. - miracil-D to man 69d. _____, complement fixation 995. _____ in U.S.A. 230u. - plant-parasitic eelworms 117c, 119b, 376a. - Systox to man 479a. - - antigens from larva & adult compared 371b. Toxocara canis & visceral larva migrans in man - in hamster, vascular changes 230v, 590a. 124a. — — Hungary 309a.
— — laboratory animals 961c. Trachurus trachurus, Echeneibothrium n.sp. in 49a. Travtrematinae nom.nov. for Leptophyllinae - - larvae, metabolism 87a. - in man, prenatal infection 230k. Treatment, see also Anthelmintics, Nematicides _ — Mesocricetus auratus 883a. (plant eelworm). - - mouse, acquired resistance 225a, 225b. -, ancylostomiasis 589a. — —, N-(p-arsenoso-benzyl) glycine amide hydrochloride 385d. -, Aphelenchoides ritzema-bosi 394a. , ascariasis 176e. ---, effect of adrenalectomy 230bd, -, Capillaria aerophila 145a. -, cerebrospinal filariasis 702b. 230be. ----, experimental 230bc. -, creeping eruption 584a. ______ & hamster 230bh. ______, immunity 775a, 961d. ______, resistance 230t, 419a. -, Dictyocaulus 536f. -, — filaria 849c. -, — viviparus 813a. - - nutria in Switzerland 211a. -, enterobiasis 43b, 147a, 232a, 906c, 955a. - polar bear, longevity of infection 199a. —, Fasciola hepática 702g. —, filariasis 180a, 379a, 877. - rat, hetrazan ineffective against 2g. Trichinelliasis, general account 520a. -, helminthiasis in man 899a, 906f. — in man 459a, 467b, 467d, 485a, 582a, 914a. — — —, ACTH 467c. —, helminths in dog 28a, 261c. —, — man 145b, 811e, 838a, 946c, 946d. _ _ _ & animals 890a. -, Heterodera rostochiensis 117e. ____ in Italy 785a. -, hookworm anaemia 531b. ___ in Britain 108b. -, hydatid 574d, 959a. -, lumbar paralysis **155a.** - -- , chloromycetin 499a. ----, cortisone 53a. -, lungworms 990, 1008. -, onchocerciasis 68a, 152m, 893a. _ — in Czechoslovakia 934a. - -, diagnosis by complement fixation 549a. -, paragonimiasis 696i. ----, effect of post-insulin hypoglycaemia -, Parascaris 965a. 481a. -, schistosomiasis 430g, 801d. ---, fouadin 170a. -, — & splenomegaly 430f. -, — haematobia 85a, 625a. __ _ in Italy 785b. ____ Lebanon 482b, 705a. -, stephanofilariasis 92c. ----, nervous symptoms 378a, 466a. -, strongyles 198b. - -- , ocular symptoms 581a. -, Taenia saginata 210b, 811d. — — in Spain 504a. — — — Switzerland 211a. -, Thelazia rhodesii 849d. -, trichinelliasis 53a. __ pig in U.S.A. 777a. -, trichuriasis 906b. —, symptomatology & diagnosis 653a.

Trichobilharzia szidati cercariae, variability 159a. Trematoda, monographed 1000, 1001. Trematode intermediaries, control 875. - n.sp. in duck, life-history 158b. -, new, in Cynoscion spp. 230db. Trichodorus sp. on crops in U.S.A. 125a. ova, technique for collecting 97j Trichohelicea n.tribe 1002. Trematodes in canines in Alaska 132c. Trichonema, see also Strongyles. -, effect of fixation on dimensions 754a. (Cylicocyclus) radiatum in horse in Argentina - in fish in Panama 588b. __ _ Hirudinea 651c. Trichonematea n.tribe 1002. -, new, in fish in Celebes 568a. Trichosomoides crassicauda in rat 317a. - in snails, cercarial shedding rate 230dc, 428s. Trichostrongyle larvae on pastures, oecology 29a. -, technique for mounting 55b, 241a. Trichostrongyles, see also Haemonchus, Nemato-Triacis scyllium, Acanthobothrium n.sp. in 49a. dirus, Trichostrongylus. -, Phyllobothrium n.spp. in 49a. in sheep, egg count variations 539a. Triaenophorus spp. in fish, life-history & control 643a. Trichostrongylidae in man in Chile 190d. - crassus plerocercoids in Leucichthys spp. 421a.

Tylenchulus semi-penetrans in citrus, reviewed | Trichostrongylosis in man in Iraq 420a. Trichostrongylus, see also Stomach worms, Trichostrongyles. — — olive & citrus 30n, 664a. - U.S.A. 133a. — in man 940a. — axei in sheep, resistance 19c. Tylenchus, see also Anguillulina. - aberrans n.sp. 384a. - medius n.sp. in bird in Russia 961bt. - retortaeformis in rabbit, inhibition of developsachsi n.sp. 867a. Tylodelphys destructor n.sp. in Basilichthys in ment 120e. Argentina 500a. Trichromadora ariminiensis n.sp. 610d. Tylopharingidae n.fam. 961bi. Trichuriasis in man, hexylresorcinol enemas 136a, Tylosurus coromandelicus, Bucephalopsis n.sp. in 185a, 270e, 270f, 294c, 811a. -, treatment 906b. Trichuris in man, hexylresorcinol enemas 946b. - muris in Neotoma floridana 320d.
- ovis, survival of free-living larvae 196b. Udonella caligorum, systematic position 296a. Udonelloidea n.class 296a. - trichiura in man 426b. Umbellularia, Sphaeronema n.g., n.sp. in 261b. ___ _ in Britain 217a. Umbrina roncador, Arhythmorhynchus n.sp. in ----, pathogenicity 270d. - - in West Africa 289a. Unilaterilecithum beloussi n.g., n.sp. in Pericrocotus roseus 1001. - - pig, experimental 849b. U.S.S.R., see Russia. Tricoma suilla n.sp. 868a. Trieste, helminths in Rattus norvegicus 139b. U.S.A., see also Alaska, North America. - -, Aelurostrongylus abstrusus 888a. Trilobus höhnelensis n.sp. 342a. telekiensis n.sp. 342a. -, ancylostomiasis 320a. Trilocularia acanthiae-vulgaris in Acanthias vul----, cestodes in shrews 21a. garis 73a. ---, Dicrocoelium dendriticum 20n, 777b. --- , Dirofilaria immitis 251d. - for Urogonoporus armatus Lühe 73a. Tringa glareola, Echinocotyle n.sp. in 688a. — — , Dracunculus insignis 433a. - hypoleucos, Aploparaksis n.sp. in 688a. - -, Choanotaenia n.sp. in 688a. -, Echinocotyle n.sp. in 688a. - - , Fasciola hepatica 772a. - incana, Diacetabulum n.g., n.sp. in 1000. -, Endocotyle n.g., n.sp. in 1000. - - Gongylonema pulchrum 261j. -, Pseudomaritrema n.g., n.sp. in 1000. - -- , helminths in cat 251b, 428n. Trinidad, ancylostomiasis 657a. ---, -- cattle 189a, 777c. ---, -- dog 28b, 95c, 428n. ---, -- fish 843a. Triodontophorus in horse in Venezuela, morphology 630a. Tropicorbis spp., susceptibility to Schistosoma mansoni 13f. --, -- man 20p, 936b. ---, - & meat inspection 222a. ---, - in Ondatra zibethica 201. - centimetralis, effect of desiccation 783k. Turbatrix, see also Anguillula. --, -- Ovis canadensis 132g. --, -- pig 486a. --, -- sheep 10e. - aceti, anthelmintics tested against 155f. ----, metabolism 455a. - -, technique for counting & culturing 97f. - -, Heterodera cruciferae 760a. — —, — — culturing 230bl.
Turdoides somervillei, Anonchotaenia n.sp. in - -, hydatid 602a. ---, lungworms 772b. - -- , Meloidogyne hapla 479e. 688a. -, Hymenolepis n.sp. in 688a. - -, microfilariae in birds 390a. Turdus dauma, Brachylecithum n.sp. in 1001. --, -- rodents 20u. Turkey, Ancylostoma duodenale 694a. ---- , Monogenea 541a, 713g. -, hydatid 694b. - -, nematodes in pig 206a. -, lungworms in sheep 288a. -, Meloidogyne 623a. --- , Neurofilaria n.g., n.sp. 81b. -, Pratylenchus zeae 760d. -, Opisthorchis felineus 963b. schistosome dermatitis 122a, 230df, , Setaria equina 963d. 338b, 646a, 746a. Turkeys, 2-amino-5 nitrothiazole 10b. -, soil nematodes attacking plants 125a. -, "blackhead" in 324a. -, Sparganum mansoni 20c. -, Heterakis gallinae & "blackhead" in 10c. - -, Trichinella spiralis 230u. -, iodine 492a. --, trichinelliasis 777a. -, phenothiazine 312a. - - Tylenchulus semi-penetrans 133a. Plagiorchis (Multiglandularis) megalorchis in Upupa epops, Maritrema n.sp. in 1000. -, Platynotrema n.sp. in 688b. Tylenchidae, biology & systematics 1006. Uranoscopus oligolepis, Symbothriorhynchus n.g., Tylenchorhynchus tesselatus n.sp. on grass 97d. n.sp. in 49a. Tylenchulus semi-penetrans in citrus 30a. Urocleidus reviewed 132h. ---, effect on growth 9a. - doloresas n.sp. in Chaenobryttus coronarius 55b.

Urogommus, key to spp. 320e. - witenbergiella n.sp. for Leucochloridium macrostomum of Witenberg, 1925 320e. Irogonoporus armatus Lühe to Trilocularia acanthiae-vulgaris 73a. Irotocus, key to spp. 320e. Uruguay, hydatid 628a, 631a.

Valeriana officinalis, Heterodera marioni in 119a. Vanellus vanellus, Capillaria vanelli in 58k. Venezuela, filariasis bancrofti 816a.

-, Meloidogyne 480a. -, nematodes in cattle 813b. —, Paragordius varius 565a. —, Triodontophorus 630a. Vertebrates, microfilariae in 50b. Viannaiea n.tribe 1002.

Viannaiinae emended 1002. Viannellea n.tribe 1002. Vigisolepis spinulosa, morphology 329a. Vine, Meloidogyne incognita in 401a.

Vipera lebetina, Ratzia joyeuxi in 12a, 58c. Vitta swifti n.sp. in Micropus affinis 688a.

Vogeloides Skryabin for Osleroides of Orlov, Davtian & Liubimov 132k. Vogeloidinae redefined 132k. Vulpecula marina, Pithophorus n.sp. in 49a.

Vulpes macrotis, Mesogyna n.g., n.sp. in 529e. vulpes, Multiceps endothoracicus in 675c.

Wallagonia attu, Haplocleidus n.sp. on 912b. -, Thaparocleidus n.g., n.sp. on 912c. Wenyonidae n.fam. 305. West Africa, see also Africa, Angola, French West

Africa, Gold Coast, Nigeria, Sierra Leone.

-, Trichuris trichiura 289a Wheat, Anguillulina tritici in 700a.

-, Anguina tritici with Corynebacterium tritici

Wuchereria, see also Filaria, Filariasis bancrofti, Microfilaria.

bancrofti, cuticular bosses 152b. development of microfilaria 674b.

- & elephantiasis in man 728a.

Wuchereria bancrofti in man, arsenamide 54a. --- in India 586t. _ _ _ Madagascar 367d. - -- , mature female in artery 2b. - -- , post-microfilarial stages 2b. ———, prenatal infection 800g.
——— in Tahiti, periodicity 318a. -, non-periodic variety 152a. --, periodicity 367b. - in Samoa, transmission 176b. - - transmitted by Culex fatigans 800f. - - & W. pacifica differentiated 379b. — malayi in Japan, first record 409a. — transmitted by Aëdes togoi 409a. - pacifica & W. bancrofti differentiated 379b.

Xantusia riversiana, Pseudoalaeuris n.sp. in 428b. -, Thelandros n.sp. in 428b. vigilis, Thelandros n.sp. in 428b. Xenopus laevis, metacercariae in 488b. Xesurus punctatus, Ichthyotrema n.g., n.sp. in 588a.

Yemen, schistosomiasis 20b. Yugoslavia, Cyathocephalus truncatus 535a. -, Cysticercus bovis 850e.

-, helminth ova in sewage & swimming pool 668a.

-, helminths in cattle 535b.

- man 909a.

- Rattus norvegicus 808a. -, hydatidosis 850d.

Zenopsis nebulosa, Eubothrioides n.g., n.sp. in 49a. Zoarces viviparus, Ascarophis n.sp. in 962a. Zoniolaiminae n.subf. 1002. Zonitoides arboreus, 1st intermediary for Conspicuum icteridorum 20bd. Zygocotyle lunatum cercaria, mucoid glands 230cg.

CORRIGENDA

0			3
36	ü	4	u
7	1	~	

119 (Journal title) For "Nachrichtenblatt des Deutschen Pflanzenschutzdienstes" read "Nachrichtenblatt für den Deutschen Pflanzenschutzdienst"

230br (Abstract) Line 1 For "Echistoma" read "Echiostoma" 281f (Abstract) Line 1 For "Blatella" read "Blattella"

331d (Title) For "GROSSE, J. E." read "CROSSE, J. E."

428h (Abstract) Line 2 For "Blatella" read "Blattella"

510a (Title) For "beet eelworm" read "beat eelworm"

Page 136 (Addendum) Line 3 For "Bochmeria" read "Boehmeria"